

# *Trypanosoma irwini* n. sp (Sarcocystidophora: Trypanosomatidae) from the koala (*Phascolarctos cinereus*)

L. M. McINNES<sup>1\*</sup>, A. GILLETT<sup>2</sup>, U. M. RYAN<sup>1</sup>, J. AUSTEN<sup>1</sup>, R. S. F. CAMPBELL<sup>3</sup>, J. HANGER<sup>2</sup> and S. A. REID<sup>1</sup>

<sup>1</sup>Division of Health Sciences, School of Veterinary and Biomedical Sciences, Murdoch University, Murdoch, Perth 6150, Western Australia

<sup>2</sup>The Australian Wildlife Hospital, Beerwah, Queensland, Australia

<sup>3</sup>School of Veterinary Science, James Cook University, Townsville, Queensland 4811, Australia

(Received 27 January 2009; revised 18 March 2009; accepted 18 March 2009)

## SUMMARY

The morphology and genetic characterization of a new species of trypanosome infecting koalas (*Phascolarctos cinereus*) are described. Morphological analysis of bloodstream forms and phylogenetic analysis at the 18S rDNA and gGAPDH loci demonstrated this trypanosome species to be genetically distinct and most similar to *Trypanosoma bennetti*, an avian trypanosome with a genetic distance of 0.9% at the 18S rDNA and 10.7% at the gGAPDH locus. The trypanosome was detected by 18S rDNA PCR in the blood samples of 26 out of 68 (38.2%) koalas studied. The aetiological role of trypanosomes in koala disease is currently poorly defined, although infection with these parasites has been associated with severe clinical signs in a number of koalas. Based on biological and genetic characterization data, this trypanosome species infecting koalas is proposed to be a new species *Trypanosoma irwini* n. sp.

Key words: *Trypanosoma* spp., koala, 18S rDNA, gGAPDH, phylogeny.

## INTRODUCTION

Trypanosomes are ubiquitous blood parasites that infect all classes of vertebrates and are the aetiological agents of severe diseases in both animals and humans. Little is known about the prevalence and pathogenesis of trypanosomes in Australian marsupials. To date 10 trypanosome species/genotypes have been identified in marsupials; *Trypanosoma thylacis* in the northern brown bandicoot (*Isodon macrourus*) (Mackerras, 1959; Mackerras and Mackerras, 1960), *T. binneyi* from the platypus (*Ornithorhynchus anatinus*) (McMillan and Bancroft, 1974), and more recently novel *Trypanosoma* spp. have been identified from the eastern barred bandicoot (*Perameles gunnii*) (Bettioli *et al.* 1996), the southern brown bandicoot (*Isodon obesulus*) (Bettioli *et al.* 1998), the woylie (*Bettongia penicillata*) (Smith *et al.* 2008), the chuditch (*Dasyurus geoffroii*) (Smith *et al.* 2008), the eastern grey kangaroo (*Macropus giganteus*), the common wombat (*Vombatus ursinus*) (Noyes *et al.* 1999) and the swamp wallaby (*Wallabia bicolor*) (Hamilton *et al.* 2005). A new species of trypanosome (*T. copemani* n. sp.) has also recently been described from Gilbert's potoroo (*Potorous gilbertii*) and quokka (*Setonix brachyurus*) (Austen *et al.* 2009).

The koala (*Phascolarctos cinereus*) has a fragmented distribution in the eastern states of Australia ranging from north-east Queensland (Qld) to the Eyre Peninsula in South Australia. A variety of factors have contributed to dramatic population declines over much of the koala's geographical range, including habitat loss and fragmentation, urbanization pressures such as increased incidents of motor vehicle strike and dog attacks and a high prevalence of disease. The koala is currently classified as vulnerable to extinction in N.S.W. and Qld under the N.S.W. Threatened Species Conservation Act 1995 and the Qld Nature Conservation (Wildlife) Regulation Act 2006.

Over the past 2 years a number of koalas have presented to the Australian Wildlife Hospital in Beerwah, Qld with serious illness associated with regenerative anaemia and trypanosome infections. The present study describes the morphological, pathological and genetic characterization of a trypanosome identified in the blood of one of these sick koalas as well as a PCR survey of other koalas presented to the hospital for unrelated conditions. We consider that the trypanosome infecting koalas is a new species and propose the name *Trypanosoma irwini* n. sp.

## MATERIALS AND METHODS

### Sources of isolates and sample collection

The Australian Wildlife Hospital at Beerwah, Qld provides veterinary care for sick and injured

\* Corresponding author: Division of Health Sciences, School of Veterinary and Biomedical Sciences, Murdoch University, Murdoch, Perth 6150, Western Australia. Tel: +61 893 602495. Fax: +61 893 104144. E-mail: L.McInnes@murdoch.edu.au

wildlife, including approximately 600 koalas per annum. These koalas are generally presented to the hospital because they have been injured or are suffering from overt disease. Most koalas presented to the hospital originated from south-east Qld or northern N.S.W.

A total of 68 blood samples were collected during routine veterinary examinations of koalas admitted to the Australian Wildlife Hospital between October 2006 and September 2008. Details of the geographical origin of each of the koalas as well as sex, age and reason for admission were recorded. For clinical examination and blood collection, the koalas were anaesthetized with an intramuscular administration of alfaxalone (Alfaxan<sup>®</sup> CD RTU, Jurox Australia) at an approximate dose rate of 3 mg/kg. Anaesthesia was maintained with a combination of oxygen and isoflurane delivered by either mask or endotracheal intubation. Blood samples of approximately 2 ml were collected by venepuncture of the cephalic vein. The blood was mixed with EDTA in a Vacutainer<sup>®</sup> tube (Becton-Dickinson, New Jersey, USA) and stored at 4 °C. The sentinel koala, a seriously ill adult male koala from Lismore N.S.W., was euthanased and a post-mortem examination was conducted. Tissues were fixed in 10% neutral buffered formalin, stained with haematoxylin and eosin and processed for histology using standard methods. All veterinary procedures were performed by registered wildlife veterinarians in accordance with standard veterinary practice.

#### Morphological measurements

Thin-blood smears were prepared using blood from the sentinel koala treated at the Australian Wildlife Hospital, Beerwah, Qld, stained with Diff-Quick stain (Dade Behring Diagnostics Australia Pty Ltd, Lane Cove, N.S.W.) and examined microscopically. Each slide was immersed in methanol for 30 sec, followed by immersion in Diff-Quick Solution I for 30 sec before counterstaining with Diff-Quick Solution II for 30 sec. The slides were then rinsed in tap water, air-dried and a cover-slip mounted using DePeX mounting medium (Merck Pty. Limited, Kilsyth, Vic., Australia).

Digital light micrograph images of trypanosomes in blood films were taken at  $\times 1000$  magnification and measurements of key morphological features (see Table 2) were performed using Image Pro Express version 5.1 (Media Cybernetics Inc., USA).

#### In vitro culture

Attempts were made to culture trypanosomes from koala blood by adding 20  $\mu$ l of fresh blood into 1.8 ml cryopreservation vials containing 1 ml of Modified

Sloppy Evans Medium (MSEM) (Noyes *et al.* 1999). Cultures were incubated in the dark at room temperature for approximately 10–14 days. Microscopic examination of wet-smear preparations of the medium from each culture was performed weekly after the initial 10 to 14-day incubation to detect motile trypanosomes at 200 $\times$  and 400 $\times$  magnification.

#### DNA extraction

Blood samples were preserved in EDTA vials and whole genomic DNA extracted using a MasterPure<sup>™</sup> DNA Purification Kit (EPICENTRE<sup>®</sup> Biotechnologies, Madison, Wisconsin, USA). DNA was eluted in 50  $\mu$ l and stored at  $-20$  °C until use. Additional DNA samples were kindly provided by a number of researchers for sequencing and phylogenetic analysis. They included: *T. cyclops* (LV492), *T. dionsii* (P3), *T. bennetti* (KT-2), *T. corvi* (LSHTM), *T. cruzi marinkellei* (B3), *T. sp.* (KG1), *T. avium* (sp30), *T. avium* (sp40), *T. avium* (SIM3), *T. avium* (*Buzzard*) and *T. avium* (A1073).

#### 18S rDNA amplification and sequencing

Two approximately 900 bp fragments of the 18S rRNA gene were amplified using 2 nested PCRs. The fragments partially overlapped and covered an approximately 1.5 kb fragment of the 18S rRNA gene. Both nested PCRs utilized the same external primers, SLF (5' gctgtttcaaggacttagc 3') and S762 (5' gacttttgcctctctaag 3'). One nested PCR used internal primers S825F (5' accgtttcggctttgttg 3') with S662 (5' gactacaatgctctctaag 3') (~904 bp product) and the other nested PCR used primers S825 (5' accgtttcggctttgttg 3') and SLIR (5' acattgtagtgcgcgtg 3') (~959 bp product). Primers S662, S762R, S825 were sourced from Maslov *et al.* (1996) and SLF and SLIR were designed during the present project. The PCR reactions were performed using 1  $\mu$ l of DNA in a 25  $\mu$ l reaction containing 1 $\times$  PCR buffer, 2 mM MgCl<sub>2</sub>, 0.4 mM dNTPs, 0.8  $\mu$ M of each primer and 0.04 U/ $\mu$ l Tth+ Taq DNA polymerase (Fisher Biotec Australia, Wembley, W.A., Australia). The PCR conditions consisted of a pre-PCR step with 95 °C for 5 min, 50 °C for 2 min and an extension of 72 °C for 4 min followed by 35 cycles of 94 °C for 30 sec, 52 °C for 30 sec and an extension of 72 °C for 2 min and 20 sec after which a final extension of 72 °C for 7 min. All positive PCR products were purified using a MO BIO UltraClean<sup>™</sup> 15 DNA Purification Kit (MO BIO Laboratories Inc. West Carlsbad, California, USA) and sequenced using an ABI Prism<sup>™</sup> Terminator Cycle Sequencing kit (Applied Biosystems, Foster City, California, USA) on an Applied Biosystem 3730 DNA Analyzer.

Table 1. Koalas examined and found positive for trypanosome infections in this study

Blood collection date	Sex	Age (years)	18S PCR	gGAPDH PCR	Location
30/10/06	M	7–8	+	+	Lismore Shire, N. S. W.
17/11/06	F	4	+	+	Lismore Shire, N. S. W.
04/12/06	M	5	+	+	Lismore Shire, N. S. W.
15/06/07	M	5	+		Caboolture Shire, Qld.
24/06/07	M	5	+		Noosa Shire, Qld.
03/09/07	F	1–2	+	+	Byron Shire, N. S. W.
28/09/07	M	3	+		Caboolture Shire, Qld.
16/10/07	M	2–3	+	+	Lismore Shire, N. S. W.
18/10/07	M	4	+		Lismore Shire, N. S. W.
24/10/07	F	3	+		Gold Coast Qld.
25/10/07	F	3	+		Caboolture Shire, Qld.
26/10/07	M	5	+		Caboolture Shire, Qld.
01/11/07	M	10	+	+	Gold Coast Qld.
06/12/07	M	5	+	+	Lismore Shire, N. S. W.
06/12/07	M	4	+	+	Lismore Shire, N. S. W.
28/12/07	F	7	+	+	Lismore Shire, N. S. W.
28/12/07	M	2	+	+	Lismore Shire, N. S. W.
28/12/07	M	9	+	+	Byron Shire, N. S. W.
28/12/07	M	<1	+	+	Lismore Shire, N. S. W.
13/02/08	F	8	+	+	Byron Shire, N. S. W.
08/05/08	F	9	+		Byron Shire, N. S. W.
08/05/08	M	6	+		Lismore Shire, N. S. W.
16/09/08	M	9	+		Redland Shire, Qld.
24/09/08	F	2	+	+	Pine Rivers Shire, Qld.
24/09/08	F	4	+	+	Redland Shire, Qld.
24/09/08	F	4	+	+	Caboolture Shire, Qld.

#### *gGAPDH amplification and sequencing*

An approximately 880 bp fragment of the glycosomal glyceraldehyde phosphate dehydrogenase (gGAPDH) gene was PCR amplified using a hemi-nested PCR. The primary PCR used primers GAPDHF (5' ctymtcggnamkgagatygayg 3') and GAPDHR (5' grtksgartadccccactcg 3') (designed during this study), followed by a secondary reaction using forward primer GAPDHF (5' ctymtcggnamkgagatygayg 3') and reverse primer G4a (5' gtttygcagsgtcgcttg 3') sourced from Hamilton *et al.* (2004). Clustal W alignments were used to design the GAPDHF and GAPDHR primers to be able to amplify *Leishmania*, *Trypanosoma brucei*, *Trypanosoma vivax* and *Trypanoplasma borrelli* gGAPDH sequences but not bovine and mouse gGAPDH sequences. The PCR reactions were performed using 1 µl of DNA in a 25 µl reaction containing 1 × PCR buffer, 2 mM MgCl<sub>2</sub>, 0.4 mM dNTPs, 0.8 µM of each primer, and 0.04 U/µl Tth + Taq DNA polymerase (Fisher Biotec Australia). The PCR conditions consisted of a pre-PCR step with 95 °C for 5 min, 50 °C for 2 min and an extension of 72 °C for 4 min followed by 35 cycles of 94 °C for 30 sec, 50 °C for 30 sec (primary PCR) or 52 °C for 30 sec (secondary PCR) with an extension of 72 °C for 2 min and 20 sec and finishing with an extension of 72 °C for 7 min. DNA sequencing was conducted as described above.

#### *Phylogenetic analysis*

The details of the 18S PCR-positive koalas are detailed in Table 1. Nucleotide sequences generated for koala-derived trypanosomes for the 18S rDNA and gGAPDH loci were combined with sequences from a number of related trypanosomatids from GenBank and phylogeny inferred from Maximum Likelihood analysis by the program PhyML (Dereeper *et al.* 2008). The reliability of the inferred trees was assessed by the approximate likelihood ratio test (aLRT) (Anisimova and Gascuel, 2006), a statistical test of branch support and an alternative to non-parametric bootstrap branch support estimation. Novel nucleotide sequence for *T. irwini* at both the 18S rDNA and gGAPDH loci were submitted to GenBank under the following Accession numbers (FJ649479 and FJ649485). In addition, novel gGAPDH sequences for *T. bennetti* (KT-2) (FJ649486), *T. cyclops* (LV492) (FJ649493), *T. dionsii* (P3) (FJ649494), *T. corvi* (LSHTM) (FJ649496), *T. avium* (sp30) (FJ649490), *T. avium* (sp40) (FJ649489), *T. avium* (Buzzard) (FJ649491), *T. avium* (A1073) (FJ649488), *T. avium* (SIM3) (FJ649487), *T. cruzi marinkellei* (B3) (FJ649495) and *T. sp.* (KG1) (FJ649492) and novel 18S rDNA sequences for *T. avium* (sp30) (FJ649482), *T. avium* (sp40) (FJ649483), *T. avium* (Buzzard) (FJ649481), *T. avium* (A1073) (FJ649480) and *T. cruzi*




Fig. 1. Bloodstream trypomastigote in a Giemsa-stained blood smear from koala (*Phascolarctos cinereus*) from Lismore, N.S.W. (sentinel koala). Scale bar represents 10  $\mu$ m.

*marinkellei* (B3) (FJ649484) generated during the present study for the phylogenetic analysis have also been submitted to GenBank.

Parsimony analysis and measurement of genetic distance were conducted using Mega version 4 (Mega4: Molecular Evolutionary Genetics Analysis software, Arizona State University, Tempe, Arizona, USA) (Tamura *et al.* 2007).

## RESULTS

### Species description

*Trypanosoma irwini* n. sp. (Fig. 1)

**Diagnosis:** Parasites observed in blood smears were long, thin organisms with a single flagellum, an undulating membrane, a kinetoplast and a nucleus, characteristics consistent with the trypomastigote stage of a trypanosomatid.

### Taxonomic summary

**Vertebrate host:** Koala (*Phascolarctos cinereus*)

**Invertebrate host:** Unknown

**Type location:** Lismore, N.S.W., Australia

**Additional locations:** A range of locations in Qld, Australia

**Site of infection:** Blood

**Pre-patent and patent periods:** Unknown

**Etymology:** This species is named *Trypanosoma irwini* in honour of the wildlife conservationist Steve Irwin.

### Parasite measurements

The morphology of the trypanosomes detected in blood smears from the sentinel koala was consistent with trypomastigote life-cycle stages of other described *Trypanosoma* spp. according to Hoare (1972). The trypomastigote-like stages had a prominent kinetoplast, undulating membrane, a pointed posterior end and a long free flagellum (Fig. 1). A summary of

Table 2. The mean dimensions and standard errors (S.E.) of morphological features of novel *Trypanosoma* sp. isolated from bloodstream forms from the koala (*Phascolarctos cinereus*) determined from the sentinel koala case

(Measurements determined from 9 trypomastigotes.)

Feature*	Observed range ( $\mu$ m)	Mean $\pm$ S.E. ( $\mu$ m)
Total length	32.1–38.7	36.3 $\pm$ 0.7
Breadth	1.9–4.5	3.0 $\pm$ 0.3
PK	3.0–5.2	3.6 $\pm$ 0.3
KN	6.0–12.1	10.3 $\pm$ 0.6
NA	8.4–16.2	11.9 $\pm$ 0.8
FF	8.6–12.8	10.3 $\pm$ 0.9

- \* Total length: body length measured along mid-line including free flagellum (total length).  
 Breadth: maximum breadth measured at nucleus level (including undulating membrane).  
 PK: distance between the posterior end and the kinetoplast.  
 KN: distance between the kinetoplast and posterior edge of the nucleus.  
 NA: distance between the anterior edge of the nucleus and the anterior end of the body.  
 FF: length of the free flagellum.

the morphological features of the bloodstream forms of *T. irwini* observed in 9 trypomastigotes is presented in Table 2.

### In vitro culture

Numerous attempts to culture the parasite were unsuccessful.

### Clinical and pathological findings in the sentinel koala

Of the 26 *T. irwini* PCR positives identified, 9 koalas sampled appeared to be free of any disease at the time of sampling. Clinical and pathological findings of the sentinel koala are outlined below. On initial presentation to the hospital, the sentinel koala displayed depression, pale mucous membranes, unilateral chronic keratoconjunctivitis and generalized weakness. Significant clinical pathological findings included profound and regenerative anaemia (packed cell volume (PCV) of 9% when normal PCV is 30–40% and haemoglobin 32 g/l, where the normal range is 88–140 g/l). Examination of a blood film revealed a marked regenerative response to the anaemia and low numbers of parasites morphologically consistent with the trypomastigote stages of a trypanosomatid organism (Fig. 1). Cytological examination of bone marrow revealed erythroid hyperplasia. The koala was euthanized after 1 month due to hepatic and renal failure. A necropsy examination revealed osteochondromas affecting a

number of ribs, haemosiderosis and extramedullary haematopoiesis of the spleen, atrophy of lymphoid tissues, bone marrow erythroid and myeloid hyperplasia, severe extensive peri-acinar necrosis of the liver, interstitial nephritis and oxalate nephrosis of the kidneys and interstitial pneumonia.

#### Genetic characterization

Of a total of 68 koalas screened for the presence of trypanosomes, using a PCR to amplify the 18S rDNA locus, 26 were determined to be infected with a *Trypanosoma* sp. (38.2%) (Table 1). Eleven of the trypanosome PCR-positive koalas were from Lismore Shire, N.S.W., 4 were from Byron Shire in N.S.W., 11 from various locations (over 5 shires) in Qld. Clustal W alignment of the partial 18S rDNA sequences amplified from all 26 koalas showed the DNA sequences to be 100% identical. For the purpose of phylogenetics analysis, some of the 26 18S rDNA-positive samples were also PCR analysed at the gGAPDH locus and 16 were successfully amplified at the gGAPDH locus and the resultant DNA sequences were found to be 100% identical. Representative DNA sequences for the 18S rDNA and gGAPDH loci of the novel trypanosome were deposited in the GenBank database as described above.

Maximum likelihood and parsimony analysis produced similar trees (data not shown). Phylogenetic analysis at both the 18S rDNA and gGAPDH loci revealed that *T. irwini* was genetically distinct and grouped with an avian trypanosome *Trypanosoma bennetti* (Figs. 2 and 3, PhyML trees shown). At the 18S rDNA locus, for which more sequence information was available from GenBank, *T. irwini* also grouped with *Trypanosoma minasense*, a trypanosome of neotropical primates and an isolate from an Australian marsupial, the chuditch (*T. sp.* CHU1). A composite tree of 18S rDNA and gGAPDH sequences also grouped *T. irwini* most closely with *T. bennetti* (Fig. 4, PhyML tree shown).

At the 18S rDNA locus, *T. irwini* had a 0.9% and 1.7% genetic distance from *T. bennetti* and *T. minasense* respectively. At the gGAPDH gene the percentage genetic distance between *T. irwini* and *T. bennetti* was 10.7% (gGAPDH sequence for *T. minasense* was unavailable).

#### DISCUSSION

This is the first description of a *Trypanosoma* sp. isolated from koalas in Australia. The observation that the 26 trypanosome-positive koalas were geographically dispersed suggests that *T. irwini* appears to have a wide distribution in N.S.W. and Qld. The pathogenicity of *T. irwini* in koalas is still unclear. Some of the clinical and pathological findings in the sentinel koala, particularly those

suggestive of extravascular haemolysis and generalized immune reaction, are entirely consistent with trypanosomiasis. However, there were other lesions in this koala that are unlikely to be aetiologically related to the trypanosome infection: the presence of various and diverse pathology precludes a definitive diagnosis of trypanosomiasis as the primary cause of illness in this case. Interestingly, approximately one third (9/26) of the *T. irwini*-positive koalas sampled for this study did not exhibit any clinical disease referable to that infection at the time they were sampled (data not shown). These koalas were admitted to the hospital as the result of traumatic injuries (by car or dog).

Immunosuppression and immunodeficiency syndromes are commonly associated with pathogenic retrovirus infection and immunosuppression has also been shown to be associated with trypanosomiasis as noted in *Trypanosoma evansi* infections (Dargantes *et al.* 2005). Infection with the koala retrovirus (KoRV) is widespread in both captive and wild koalas and associated disease incidence is high (Hanger *et al.* 2000, 2003; Tarlinton *et al.* 2005). It is possible that a trypanosome infection in a healthy koala remains innocuous but when the immune status of the koala changes due to infection such as with KoRV, clinical signs of trypanosomiasis manifest. The level of retroviral RNA in serum was not measured in the euthanased koala, and therefore it is not possible to comment on the likely involvement of the KoRV in the disease observed in that individual. However, the fact that *Trypanosoma irwini* n. sp. was also identified in koalas without any disease referable to this infection suggests that it may be non-pathogenic or have low pathogenicity in many individuals, presumably those with intact immune systems.

Although it is possible that *T. irwini* n. sp. co-evolved with its host and may have limited pathogenicity in many individuals, its pathogenic potential (and therefore consequences for conservation) should not be ignored given the currently incomplete understanding of KoRV-associated disease, particularly the immunodeficiency syndrome. Further work is required to investigate the prevalence and pathogenic potential of *T. irwini* in koalas, particularly in those co-infected with KoRV.

Current phylogenetic evidence suggests that *T. irwini* is genetically most closely related to the avian trypanosome *T. bennetti*, which was isolated from an American kestrel (*Falco sparverius*) (Kirkpatrick and Terway-Thompson, 1986) and *T. minasense*, which infects over 32 species of neotropical primates (Ziccardi *et al.* 1996). Most of the morphological measurements of the bloodstream form of *T. irwini* are within the range of those published for cultured *T. bennetti* and *T. minasense* with the exception of the measurement from the posterior end to the kinetoplast (PK). *Trypanosoma*




Fig. 2. Phylogenetic analysis of the relationships of *Trypanosoma* spp. and novel trypanosomatid *Trypanosoma irwini* isolates (26 in total) from the koala using Maximum Likelihood analysis performed by PhyML 3.0 based on partial 18S rRNA gene sequences (~1500 bp). Branch support values (> 60%) determined by aLRT. **a**) Demonstrates the phylogenetic position of *T. sp.* isolate Chuditch (CHU1) (~600 bp sequences). **b**) Phylogenetic position of *T. copemani* isolates (Q10, Q3, P94 and P63) and *T. sp.* TRY1 and TRY2 (~600 bp sequences). Scale bar represents the number of substitutions per site.

*minasense* is usually reported to have a posterior to kinetoplast measurement of 5.7–15 µm (Hoare, 1972; Sato *et al.* 2008) and *T. bennetti* 6.7–12.3 µm (Kirkpatrick and Terway-Thompson, 1986). In

contrast, the PK for *T. irwini* is considerably shorter at 3–5.2 µm. The utility of delineating trypanosome species based on morphometrics is however questionable as pleomorphism of bloodstream




Fig. 3. Phylogenetic analysis of the relationships of *Trypanosoma* spp. and the novel trypanosomatid *Trypanosoma irwini* isolates (16 in total) from the koala using Maximum Likelihood analysis performed by PhyML 3.0 based on partial gGAPDH sequences (~900 bp). Branch support values (>60%) determined by aLRT. Scale bar represents the number of substitutions per site.

trypanosomatid forms of numerous *T. spp.* has been documented. Pleomorphism of bloodstream forms of *T. minasense* has been reported in different experimentally infected hosts, namely squirrel monkeys and marmosets, where the level of variation did not conform to previously measured *T. minasense* blood forms (Ziccardi and Lourenco-de-Oliveira, 1999). Pleomorphism of bloodstream forms of *T. bennetti*

has also been reported with an increase in *in vitro* culturing temperature producing a posterior migration of the kinetoplast, affecting the PK values (Kirkpatrick and Terway-Thompson, 1986).

There are no established criteria for delimiting species within the genus *Trypanosoma*. However, it has been proposed that the genetic distance at the 18S rDNA locus of 0.7% and 3.4% is sufficient to




Fig. 4. Phylogenetic analysis of the relationships of *Trypanosoma* spp. and the novel trypanosomatid *Trypanosoma irwini* isolates (16 in total) from the koala based on composite gGAPDH and 18S rDNA partial sequences (~900 and ~1500 bp respectively) using Maximum Likelihood analysis performed by PhyML 3.0. Branch support values (>60%) determined by aLRT. Scale bar represents the number of substitutions per site.

classify distinct species for *Theileria* sp. and *Babesia* sp. respectively (Schnittger *et al.* 2003). In the present study, molecular and phylogenetic analysis at 2 unlinked loci indicated that *T. irwini* warrants separate species status. The genetic distance between *T. irwini* and its closest relative *T. bennetti* at the 18S rDNA locus is 0.9%, whereas the genetic difference between 2 established species, *T. avium*

isolates and *T. grayi* is 0.6–0.8%. At the gGAPDH locus, established species *T. conorhini*, *T. vespertilionis*, *T. mega* and *T. rotarium* have between 4.4% and 5.3% genetic distances, whilst intraspecies genetic distance within *T. avium* is 0.9–2%. The genetic distance between *T. bennetti* and *T. irwini* at the gGAPDH locus is 10.7%, which is equal to or greater than genetic distances between established species.



The 3 phylogenetic trees produced in order to determine the evolutionary position of *T. irwini* amongst the trypanosomes largely produced the same topology of other researchers with a few exceptions. The commonly reported aquatic, avian, *T. lewisi*, *T. cruzi*, *T. brucei* and *T. theileri* clades reported by Hamilton *et al.* (2007) were generated in the 18S rDNA, gGAPDH and concatenated analyses with the exception of a separation of the avian clade in the 18S rDNA tree. This split in the avian clade has also been produced in 18S rDNA ML analysis by other researchers (Hamilton *et al.* 2004, 2005; Votykka *et al.* 2004). It has been suggested, however, that the 18S rDNA locus is an unreliable marker for inferring deep level trypanosome phylogenetic analysis (Hamilton *et al.* 2004) due to the fact that the 18S rRNA genes of taxa in the *Trypanosoma* genus are evolving at least 4 significantly different rates (Stevens and Rambaut, 2001). This is primarily why the gGAPDH locus was also analysed as it is increasingly used in *Trypanosoma* phylogeny and is under different evolutionary constraints to the 18S rRNA gene (Hamilton *et al.* 2004). The topology of the gGAPDH and concatenated 18S rDNA and gGAPDH trees were nearly identical and very similar to equivalent trees presented by Hamilton *et al.* (2007) and Viola *et al.* (2009). The variation between the gGAPDH and concatenated tree was primarily due to lack of sequence data for the gGAPDH locus as the '*T. irwini*' clade only comprised *T. irwini* and *T. bennetti* due to *T. minasense* gGAPDH sequence being unavailable. In addition, the gGAPDH analysis of the '*T. irwini*' clade showed low aLRT branch support (37) for association with the '*T. theileri*' clade, whilst in the concatenated analysis the '*T. irwini*' clade is related to the 'lizard' clade with moderate branch support (64). The addition of the *T. minasense* gGAPDH sequence will perhaps help to resolve the potential relationship with the 'lizard' clade in the future.

The phylogenetic relationships of *T. irwini* may also be further clarified once more marsupial-derived trypanosome species are characterized. The relationship between *T. irwini* and *T. thylacis* from the northern brown bandicoot (*Isoodon macrourus*) (Mackerras, 1959; Mackerras and Mackerras, 1960), the *Trypanosoma* spp. from the southern brown bandicoots (*Isoodon obesulus*) (Bettiol *et al.* 1998) and eastern barred bandicoots (*Perameles gunnii*) (Bettiol *et al.* 1996) cannot be evaluated because there are no genetic sequences available for these trypanosomes. Despite the wombat being the closest living relative to the koala (Osborne *et al.* 2002), the *T. copemani* genotypes (AAI, AAP and H26) isolated from wombats are genetically distinct (genetic distance 5.6% at 18S rDNA and 20.4% at gGAPDH) from *T. irwini*. This is also true for quokka and Gilbert's potoroo isolates of *T. copemani* (genetic distance 5.3% at 18S rDNA) and woylie-derived

*Trypanosoma* isolates (TRY1 and TRY2) (genetic distance 11.4% and 10.6% at 18S rDNA respectively) which are phylogenetically associated with *T. copemani* (Austen *et al.* 2009). Of all of the marsupial trypanosome genotypes identified thus far, the chuditch isolate (CHU1) was the closest marsupial-derived genotype to *T. irwini* with a genetic distance of 2.8% at the 18S rDNA locus, creating a clade with *T. minasense*, *T. bennetti* and *T. irwini*. The relationship of the chuditch isolate could not be examined in detail, as the available 18S rDNA sequence was short (only 563 bp) and to date no gGAPDH sequence has been generated.

The '*T. irwini*' clade formed is a perplexing mix of isolates with vast differences in geography and hosts, with *T. bennetti* isolated from a bird in North America (also found in a PCR survey in African songbirds (Sehgal *et al.* 2001), *T. minasense* from neotropical primates in South America, the chuditch (CHU1) or western quoll, a carnivorous Australian marsupial and *T. irwini* from the koala. The clade formed in the analysis could be evidence of host switching as has been suggested for *T. binneyi*, the platypus trypanosome and *T. therezieni*, a trypanosome of Madagascan chameleons that both fall in the 'aquatic' clade of trypanosomes from fish and amphibians. The phylogenetic placement of these trypanosomes indicates that host ecology may be more influential than host phylogeny in the evolution of some trypanosomes (Stevens *et al.* 2001). Host switching could also be evidence of trypanosome and vector co-evolution rather than host co-evolution. It should be noted that the '*T. irwini*' clade presents the first evidence of possible mammalian-avian host switching in the trypanosomes. The vast differences in host phylogeny, host ecology and lack of knowledge about vectors of the trypanosomes of the '*T. irwini*' clade do, however, prevent formation of a plausible hypothesis regarding what factors other than genetics unite the trypanosomes of this clade.

Despite numerous attempts with fresh blood samples from trypanosome-infected koalas, establishment of *in vitro* cultures in Modified Sloppy Evans Media (MSEM) was not successful. With the benefit of the phylogenetic analysis, culturing attempts in the future could focus on testing media used in the *in vitro* culturing of the genetically most similar trypanosomes *T. bennetti* (Kirkpatrick and Terway-Thompson, 1986) and *T. minasense* (Ziccardi *et al.* 1996). A feeder cell line system (possibly even host derived) may also need to be developed as used for the successful isolation and propagation of numerous trypanosomes including *Trypanosoma* isolate KG1 from a tick (Thekisoe *et al.* 2007) and *Trypanosoma* isolate TSD1 isolated from a Sika deer (Hatama *et al.* 2007).

Determining the vector of *T. irwini* is important to understanding the transmission and phylogenetic relationships of this parasite. However, the vectors

of *T. bennetti* and *T. minasense* are currently unknown and can therefore not offer any suggestions for potential vectors of *T. irwini*. Trypanosome vectors are commonly haematophagous arthropods such as flies, fleas and bugs. Ectoparasites, which have been identified on koalas, include ticks (*Ixodes* spp. and *Haemaphysalis* spp.), fleas (e.g. *Ctenophthalmus felis*), mites (*Austrochirus perkinsi*, *Sarcoptes scabiei*, *Demodex* spp. and *Notoedres cati*) (Jackson *et al.* 2003) and mosquitoes. Mosquitoes and flies are often observed worrying the noses and the eartips of koalas (Gillett, A., personal communication). The wombat flea (*Lycopsylla nova*) has been suggested as a possible vector for the wombat isolate H26 (Noyes *et al.* 1999). Ticks are also a possible vector for *T. irwini* because they have been shown to be naturally infected with trypanosomes in a number of studies (el Kady, 1998; Latif *et al.* 2004; Thekisoe *et al.* 2007).

In conclusion, the genetic analysis of 2 unlinked loci confirm that the trypanosome identified to be infecting and potentially causing disease in the koala is a new species; *Trypanosoma irwini* and the first to be described in the koala.

The authors thank the staff and carers at, and connected with, the Australia Wildlife Hospital, Beerwah, Qld. Additional DNA samples were gratefully received for 18S rDNA and gGAPDH phylogenetic tree construction: *T. cyclops* (LV492) and *T. dionsii* (P3) from Wendy Gibson, *T. bennetti* (KT-2; ATCC 50102), *T. corvi* (LSHTM), *T. avium* (Buzzard), *T. avium* (sp30), *T. avium* (sp40), and *T. avium* (A1073) from Julius Lukes and Jan Votypka, *T. cruzi marinkellei* (B3) from Christian Barnabe and *T. sp.* (KG1) was obtained from Noburu Inoue and Oriel Thekisoe. The authors would also like to thank Patrick Hamilton for invaluable phylogenetic advice.

## REFERENCES

- Anisimova, M. and Gascuel, O.** (2006). Approximate likelihood-ratio test for branches: A fast, accurate, and powerful alternative. *Systematic Biology* **55**, 539–552.
- Austen, J., Jefferies, R., Friend, T., Adams, P., Ryan, U. and Reid, S.** (2009). Morphological and molecular characterization of *Trypanosoma copemani* n.sp. (Trypanosomatidae) isolated from Gilbert's potoroo (*Potorous gilbertii*) and quokka (*Setonix brachyurus*). *Parasitology* **136**, (in the Press).
- Bettiol, S. S., Goldsmid, J. M., Le, D. D. and Driessen, M.** (1996). The first record of a member of the genus *Hepatozoon* in the eastern barred bandicoot (*Perameles gunnii*) in Tasmania. *Journal of Parasitology* **82**, 829–830.
- Bettiol, S. S., Jakes, K., Le, D. D., Goldsmid, J. M. and Hocking, G.** (1998). First record of trypanosomes in Tasmanian bandicoots. *Journal of Parasitology* **84**, 538–541.
- Dargantes, A. P., Campbell, R. S. F., Copeman, D. B. and Reid, S. A.** (2005). Experimental *Trypanosoma evansi* infection in the goat. II. Pathology. *Journal of Comparative Pathology* **133**, 267–276.
- Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., Dufayard, J. F., Guindon, S., Lefort, V., Lescot, M., Claverie, J. M. and Gascuel, O.** (2008). Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Research* **36**, W465–469.
- El Kady, G. A.** (1998). Protozoal parasites in tick species infesting camels in Sinai Peninsula. *Journal of the Egyptian Society of Parasitology* **28**, 765–776.
- Hamilton, P. B., Gibson, W. C. and Stevens, J. R.** (2007). Patterns of co-evolution between trypanosomes and their hosts deduced from ribosomal RNA and protein-coding gene phylogenies. *Molecular and Phylogenetic Evolution* **44**, 15–25.
- Hamilton, P. B., Stevens, J. R., Gaunt, M. W., Gidley, J. and Gibson, W. C.** (2004). Trypanosomes are monophyletic: evidence from genes for glyceraldehyde phosphate dehydrogenase and small subunit ribosomal RNA. *International Journal for Parasitology* **34**, 1393–1404.
- Hamilton, P. B., Stevens, J. R., Gidley, J., Holz, P. and Gibson, W. C.** (2005). A new lineage of trypanosomes from Australian vertebrates and terrestrial bloodsucking leeches (Haemadipsidae). *International Journal for Parasitology* **35**, 431–443.
- Hanger, J. J., Bromham, L. D., McKee, J. J., O'Brien, T. M. and Robinson, W. F.** (2000). The nucleotide sequence of koala (*Phascolarctos cinereus*) retrovirus: a novel type C endogenous virus related to Gibbon ape leukemia virus. *Journal of Virology* **74**, 4264–4272.
- Hanger, J. J., McKee, J. J., Tarlington, R. and Yates, A.** (2003). Cancer and haematological disease in koalas: A clinical and virological update. In *Annual Conference of the Australian Veterinary Association*, pp. 19–30. Australian Association of Veterinary Conservation Biology Section, Cairns, Australia.
- Hatama, S., Shibahara, T., Suzuki, M., Kadota, K., Uchida, I. and Kanno, T.** (2007). Isolation of a *Megatrypanum* trypanosome from sika deer (*Cervus nippon yesoensis*) in Japan. *Veterinary Parasitology* **149**, 56–64.
- Hoare, C. A.** (1972). *The Trypanosomes of Mammals. A Zoological Monograph*. Blackwell Scientific Publications, Oxford, UK.
- Jackson, S., Reid, K., Spittal, D. and Romer, L.** (2003). Koalas. In *Australian Mammals: Biology and Captive Management* (ed. Jackson, S.), pp. 145–181. CSIRO Publishing, Collingwood, Australia.
- Kirkpatrick, C. E. and Terway-Thompson, C. A.** (1986). Biochemical characterization of some raptor trypanosomes. II. Enzyme studies, with a description of *Trypanosoma bennetti* n. sp. *Canadian Journal of Zoology* **64**, 195–203.
- Latif, A. A., Bakheit, M. A., Mohamed, A. E. and Zwegarth, E.** (2004). High infection rates of the tick *Hyalomma anatolicum anatolicum* with *Trypanosoma theileri*. *Onderstepoort Journal of Veterinary Research* **71**, 251–256.
- Mackerras, M.** (1959). The haematozoa of Australian mammals. *Australian Journal of Zoology* **7**, 105–135.
- Mackerras, M. and Mackerras, I.** (1960). The haematozoa of Australian birds. *Australian Journal of Zoology* **8**, 226–260.
- Maslov, D. A., Lukes, J., Jirku, M. and Simpson, L.** (1996). Phylogeny of trypanosomes as inferred from the

- small and large subunit rRNAs: Implications for the evolution of parasitism in the trypanosomatid protozoa. *Molecular and Biochemical Parasitology* **75**, 197–205.
- McMillan, B. and Bancroft, B.** (1974). On the morphology of *Trypanosoma binneyi* Mackerras, 1959 from the platypus *Ornithorhynchus anatinus*. *International Journal for Parasitology* **4**, 441–442.
- Noyes, H. A., Stevens, J. R., Teixeira, M., Phelan, J. and Holz, P.** (1999). A nested PCR for the ssrRNA gene detects *Trypanosoma binneyi* in the platypus and *Trypanosoma* sp. in wombats and kangaroos in Australia. *International Journal for Parasitology* **29**, 331–339.
- Osborne, M. J., Christidis, L. and Norman, J. A.** (2002). Molecular phylogenetics of the Diprotodontia (kangaroos, wombats, koala, possums, and allies). *Molecular Phylogenetics and Evolution* **25**, 219–228.
- Sato, H., Leo, N., Katakai, Y., Takano, J. I., Akari, H., Nakamura, S. I. and Une, Y.** (2008). Prevalence and molecular phylogenetic characterization of *Trypanosoma (Megatrypanum) minasense* in the peripheral blood of small neotropical primates after a quarantine period. *Journal of Parasitology* **94**, 1128–1138.
- Schnittger, L., Yin, H., Gubbels, M. J., Beyer, D., Niemann, S., Jongejan, F. and Ahmed, J. S.** (2003). Phylogeny of sheep and goat *Theileria* and *Babesia* parasites. *Parasitology Research* **91**, 398–406.
- Sehgal, R. N., Jones, H. I. and Smith, T. B.** (2001). Host specificity and incidence of *Trypanosoma* in some African rainforest birds: a molecular approach. *Molecular Ecology* **10**, 2319–2327.
- Smith, A., Clark, P., Averis, S., Lymbery, A. J., Wayne, A. F., Morris, K. D. and Thompson, R. C.** (2008). Trypanosomes in a declining species of threatened Australian marsupial, the brush-tailed bettong *Bettongia penicillata* (Marsupialia: Potoroidae). *Parasitology* **135**, 1329–1335.
- Stevens, J. and Rambaut, A.** (2001). Evolutionary rate differences in trypanosomes. *Infection, Genetics and Evolution* **1**, 143–150.
- Stevens, J. R., Noyes, H. A., Schofield, C. J. and Gibson, W.** (2001). The molecular evolution of Trypanosomatidae. *Advances in Parasitology* **48**, 1–56.
- Tamura, K., Dudley, J., Nei, M. and Kumar, S.** (2007). MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* **24**, 1596–1599.
- Tarlinton, R., Meers, J., Hanger, J. and Young, P.** (2005). Real-time reverse transcriptase PCR for the endogenous koala retrovirus reveals an association between plasma viral load and neoplastic disease in koalas. *Journal of General Virology* **86**, 783–787.
- Thekiso, O. M., Honda, T., Fujita, H., Battsetseg, B., Hatta, T., Fujisaki, K., Sugimoto, C. and Inoue, N.** (2007). A trypanosome species isolated from naturally infected *Haemaphysalis hystricis* ticks in Kagoshima Prefecture, Japan. *Parasitology* **134**, 967–974.
- Viola, L. B., Almeida, R. S., Ferreira, R. C., Campaner, M., Takata, C. S., Rodrigues, A. C., Paiva, F., Camargo, E. P. and Teixeira, M. M.** (2009). Evolutionary history of trypanosomes from South American caiman (*Caiman yacare*) and African crocodiles inferred by phylogenetic analyses using SSU rDNA and gGAPDH genes. *Parasitology* **136**, 55–65.
- Votypka, J., Lukes, J. and Obornik, M.** (2004). Phylogenetic relationship of *Trypanosoma corvi* with other avian trypanosomes. *Acta Protozoologica* **43**, 225–231.
- Ziccardi, M. and Lourenco-De-Oliveira, R.** (1999). Polymorphism in trypomastigotes of *Trypanosoma (Megatrypanum) minasense* in the blood of experimentally infected squirrel monkey and marmosets. *Memorias do Instituto Oswaldo Cruz* **94**, 649–653.
- Ziccardi, M., Lourenco De Oliveira, R. and Nogueira, R.** (1996). The haemoculture of *Trypanosoma minasense* Chagas, 1908. *Memorias do Instituto Oswaldo Cruz* **91**, 501–505.