



Murdoch
UNIVERSITY

MURDOCH RESEARCH REPOSITORY

<http://researchrepository.murdoch.edu.au/18322/>



Coventry, Jocelyn (2013) *Review of aspects of breeding herd performance from beef cattle projects on the arid rangelands of the Alice Springs District*. Masters by Research thesis, Murdoch University is licensed under a [Creative Commons Attribution-NonCommercial 3.0 Unported License](https://creativecommons.org/licenses/by-nc/3.0/).

Permissions beyond the scope of this license may be available at jocelyn.coventry@nt.gov.au.

**Review of Aspects of Breeding Herd Performance
from Beef Cattle Projects on the Arid Rangelands
of the Alice Springs District**

by
Jocelyn Coventry BSc, BVMS, MACVSc

This thesis is presented for the degree of
Master of Philosophy,
of Murdoch University, WA
in the year of 2013

DECLARATION

I declare that this thesis is my own account of my research.

The main content is based on review of data from three primary beef cattle projects.

Published and unpublished reports from these primary beef cattle projects have been cited where used for summaries and analyses.

.....

(Jocelyn Coventry

date)

ABSTRACT

To effectively record the variability that is a feature of beef cattle herds on arid rangelands, sustained periods of objective data collection are required. This has been infrequently achieved on pastoral properties in the Alice Springs district of the Northern Territory. In this thesis, historical data are analysed from three primary beef cattle projects with breeding herds north of Alice Springs. Each of these projects involved data and sample collections during a 4- to 5-year period. Publications from each project were reviewed to identify summaries on breeding herd performance and gaps in data analyses. Where gaps were identified and data were available, new summaries and analyses were undertaken. The results of those analyses enabled comparison with existing knowledge about beef cattle production on arid rangelands, and provided new information on reproductive performance, mortality, nutritional deficiencies, infestations and infections in continuously-mated herds of the Alice Springs district. Information on reproductive performance and mortality was provided for different age and gender groups respectively. The aged cow group had the highest percentage of conception losses before branding (10.3%). Heifers had the highest peri-/ post-natal conception losses (4.5%), with suspected dystocia and mortality in up to 20% of some heifer groups. For lactating individuals in 'forward store' body condition at muster, the pregnancy percentage of heifers (45%) was similar to that of other cow age-groups (48–53%). Extrapolated (unconfirmed) annual mortality was 6.5% in herd bulls and up to 3.1% in cows.

Information on specific cattle health issues was provided about potential dietary deficiencies in phosphorus (P), vitamin A and nitrogen (N), as well as about intestinal parasite infestations and infection with bovine viral diarrhoea virus (BVDV).

Phosphorus deficiency, as defined by serum P (≤ 1.1 mmol/L) and faecal P ($\leq 0.19\%$),

was associated with low rainfall, dry grass, absence of pasture germination, landsystems with low soil P, and lactating or aged cows. Sub-optimal levels of serum vitamin A (< 0.26 mg/L) were associated with dry seasonal conditions, low body condition and aged cows. A fall in cattle nutrition based on faecal N below 1.23% was related to the presence of lactation and pasture germination. Clinical intestinal nematodiasis (faecal egg counts over 700 eggs per gram) in weaned calves was correlated with a large rainfall event after a long dry period. In one herd endemically infected with BVDV, individual cows became seronegative during a 3-year period.

Results of the thesis study provided benchmarks to establish and validate measures of cattle health and reproductive performance in the Alice Springs district. Regional reference ranges were established for albumin (cows: 32–43 g/L) and haemoglobin (cows: 125–153 g/L). Averages over 3- to 5-year periods were provided for annual branding (range: 64–87%) and weaning (range: 79–84%) percentages. The highest mean annual number of branded calves per herd bull was calculated at the lowest examined bull percentage (24.6 calves at 4.1% bulls). The average weight of weaned calves per 100 kg of mated cows was established under varied herd and seasonal conditions (range of averages: 37–64 kg).

The challenges of collecting and analysing data from extensively-managed cattle herds can necessitate the use of default measures, extrapolations and estimations. In this thesis, the use of default measures helped to define conception losses, measures of body mass-at-conception, annual mortality, arid rangeland pasture conditions for ‘optimal’ cattle nutrition, variable seasonal conditions, and diseased sub-groups.

In summary, this study contributes to the knowledge about cattle grazing on arid rangelands. The key contributions are in the analyses that establish and validate benchmarks, as well as in the discussions on reproductive performance and ‘best practice’ management.

TABLE OF CONTENTS

DECLARATION	i
ABSTRACT	iii
INDEX OF TABLES	vii
INDEX OF FIGURES	viii
INDEX OF APPENDICES	ix
GENERAL ACKNOWLEDGMENTS	xi
ABBREVIATIONS	xiii
CHAPTER 1. INTRODUCTION	1
SECTION 1.1. BACKGROUND TO THE PRESENT STUDY	1
CHAPTER 2. REVIEW	3
SECTION 2.1. CATTLE PRODUCTION - ALICE SPRINGS DISTRICT AND COMPARABLE ARID AREAS.....	3
2.1.1. Natural Resources	3
2.1.2. Cattle Industry.....	9
2.1.3. General Herd.....	12
2.1.4. Herd Bulls	30
2.1.5. Cows	34
2.1.6. Calf Production	38
SECTION 2.2. LABORATORY TESTING.....	44
SECTION 2.3. SUMMARY	46
CHAPTER 3. MATERIALS AND METHODS	47
SECTION 3.1. INTRODUCTION	47
SECTION 3.2. SITES	47
3.2.1. Location	47
3.2.2. Cattle.....	51
SECTION 3.3. SAMPLES AND DATA FOR STUDY	51
3.3.1. Collection at Sites	51
3.3.2. Management of Samples and Data	53
3.3.3. Interpretation and Analysis of Data	54
SECTION 3.4. SUMMARY	76
CHAPTER 4. HERD BULL RESULTS	77
SECTION 4.1. INTRODUCTION	77
SECTION 4.2. HERD BULL DYNAMICS	77
4.2.1. Herd Bulls in Paddock	77
4.2.2. Recruitment of Herd Bulls.....	79
4.2.3. Removal of Herd Bulls	80
SECTION 4.3. SUMMARY	82

CHAPTER 5. COW RESULTS.....	83
SECTION 5.1.INTRODUCTION	83
SECTION 5.2.REPRODUCTIVE PERFORMANCE.....	83
5.2.1. Conception	83
5.2.2. Lactation	89
5.2.3. Calf Production	93
SECTION 5.3.HERD DYNAMICS	96
5.3.1. Removal of Cows	96
5.3.2. Recruitment of Cows	100
SECTION 5.4.SUMMARY	103
CHAPTER 6. LABORATORY TEST RESULTS.....	107
SECTION 6.1.INTRODUCTION	107
SECTION 6.2.BACTERIOLOGY	107
6.2.1. Preputial Scrapings	107
SECTION 6.3.SEROLOGY	107
6.3.1. Serum Antibody to Leptospires	107
6.3.2. Serum Antibody to Viruses	109
SECTION 6.4.BIOCHEMISTRY AND HAEMATOLOGY	110
6.4.1. Optimal Biochemical and Haematological Values.....	110
6.4.2. Variation in Biochemical Values.....	111
SECTION 6.5.FAECAL CHEMISTRY	112
6.5.1. Variation in Faecal Chemistry Values	112
SECTION 6.6.FAECAL PARASITOLOGY	116
6.6.1. Identity of Intestinal Parasites	116
6.6.2. Variation in Faecal Egg Counts and Oocyst Counts	116
SECTION 6.7.SUMMARY	118
CHAPTER 7. DISCUSSION.....	123
SECTION 7.1.INTRODUCTION	123
SECTION 7.2.REVIEW OF HERD BULL RESULTS	123
SECTION 7.3.REVIEW OF COW RESULTS.....	128
SECTION 7.4.REVIEW OF LABORATORY TEST RESULTS	156
7.4.1. Bacteriology.....	156
7.4.2. Serology.....	160
7.4.3. Biochemistry and Haematology	170
7.4.4. Faecal Chemistry	177
7.4.5. Faecal Parasitology.....	183
SECTION 7.5.SUMMARY	190
APPENDICES	193
REFERENCES	243

INDEX OF TABLES

Table 3-1.	Location and primary beef cattle projects for sites.	49
Table 3-2.	Summary of rainfall and temperatures at sites.	50
Table 3-3.	Factors limiting data on herd bulls and cows at sites A, B and C.	56
Table 3-4.	Management techniques for data on herd bulls at sites A, B and C.	57
Table 3-5.	Management techniques for data on cows at sites A, B and C.	59
Table 4-1.	Bull percentage: financial year range for sites A, B and C.	77
Table 4-2.	Mean number of branded calves per herd bull: financial year range for sites A and B.	78
Table 4-3.	Recruited percentage of herd bulls, averaged per opening conditions of financial years at sites A and B.	79
Table 4-4.	Removed percentage of herd bulls, averaged per opening conditions of financial years at sites A and B.	81
Table 4-5.	Recorded mortality percentage of herd bulls, averaged per opening conditions of financial years at site B.	81
Table 5-1.	Annual conception percentage per cow age-group: first two years at site B.	84
Table 5-2.	Mean monthly conception percentage after muster and rainfall events at sites A and B.	86
Table 5-3.	Weight-at-conception per cow age-category: financial year range for sites A, B and C.	87
Table 5-4.	Body condition score-at-conception per cow age-category: median and range for sites B and C.	87
Table 5-5.	Lactation percentage averaged per opening conditions of financial years at sites B and C.	89
Table 5-6.	Lactation percentage per 6-months cumulative rainfall at sites B and C.	90
Table 5-7.	Lactation percentage per body condition at sites B and C.	90
Table 5-8.	Lactation percentage per cow age-group at sites B and C.	91
Table 5-9.	Branding percentage: average and financial year range for sites A and B.	93
Table 5-10.	Weaning percentage: average and calendar year range for sites B and C.	94
Table 5-11.	Mean weight of weaner-age calves: range at muster for sites A, B and C.	95
Table 5-12.	Average weight of weaned calves per head of mated cows and per 100 kg of mated cows: calendar year range for sites B and C.	96
Table 5-13.	Reasons for removal: numbers of cows that were removed for sole and composite reasons at site B.	97
Table 5-14.	Removed percentage of cows, averaged per opening conditions of financial years at site B.	98
Table 5-15.	Recorded mortality percentage of cows, averaged per opening conditions of financial years at site B.	99

Table 5-16.	Summary of the last records for a group of 56 persistently absent cows from site B.	99
Table 5-17.	Recruited percentage of cows, averaged per opening conditions of financial years at site B.	100
Table 5-18.	Lactation percentage and retained percentage: heifers compared with young cows at site B.	102
Table 7-1.	Regional reference ranges for biochemistry and haematology of cows on arid rangelands at site B.	171

INDEX OF FIGURES

Figure 2-1.	Location of the Alice Springs district in the Northern Territory, as shown by region 711 in an Australian agricultural and grazing industries survey (AAGIS) (Hooper 2001).	3
Figure 3-1.	Location of sites with respect to climatic classifications for the Northern Territory.	48
Figure 5-1, parts a. and b.	Monthly rainfall and average monthly minimum temperature, correlated with monthly conception percentage for site B.	85
Figure 5-2, parts a. and b.	Body condition score-at-conception, by percentage of cow age-category for sites B and C.	88
Figure 5-3, parts a. and b.	Mean weight of weaned calves at musters for sites B and C.	95

INDEX OF APPENDICES

Appendix 1. Paddock infrastructure, landsystems and rangeland pasture-types at sites.	193
Appendix 2. Monthly rainfall totals for sites.	196
Appendix 3. Cattle herd structure and management at sites.	198
Appendix 4. Number of cows, herd bulls and calves related to annual calf production at sites.	201
Appendix 5. Range of cattle numbers and stocking densities at sites.	203
Appendix 6. Gantt charts: number of cows in paddock at sites.	205
Appendix 7. Number of mustered cows, including those removed (by culling or mortality) and those recruited for sites.	207
Appendix 8. Type and delegation of primary beef cattle project activity at sites.	210
Appendix 9. Muster dates at sites.	212
Appendix 10. Categories of cattle and environmental variables in study.	214
Appendix 11. Comparing mean faecal counts of sub-groups for stratification of representative groups at site B.	216
Appendix 12, parts a. and b. Cattle data and samples at sites.	217
Appendix 13. Pregnancy percentage at muster, within categories of body condition for lactating and non-lactating cows, per cow age-group at sites B and C.	225
Appendix 14. Percentage of confirmed pregnancies with lactation failure, per stage of reproductive loss and cow age-group at site B.	227
Appendix 15. Percentage of herd bulls with a positive culture of preputial scrapings for <i>Campylobacter fetus</i> subsp. <i>venerealis</i> and <i>Trichostrongylus axei</i> , per bull age-category and year at site B.	228
Appendix 16. Percentage of herd bulls and cows with MAT titres to <i>L. sv. hardjo</i> , <i>L. sv. pomona</i> and <i>L. sv. tarassovi</i> , per age-categories of bulls and cows at site B.	229
Appendix 17. Percentage of herd bulls and cows with MAT titres to <i>L. sv. hardjo</i> , <i>L. sv. pomona</i> and <i>L. sv. tarassovi</i> , per 2-months cumulative rainfall at site B.	230
Appendix 18. Lactation and pregnancy percentages with positive titres to <i>L. sv. hardjo</i> and <i>L. sv. tarassovi</i> : odds ratios and χ^2 values, per cow age-category at site B.	231
Appendix 19. Percentage of herd bulls and cows with AGID level of antibody to BVDV and SN titres to BHV1, per age-categories of bulls and cows at site B.	232
Appendix 20. Lactation and pregnancy percentages with positive titres to BVDV and BHV1: odds ratios and χ^2 values, per cow age-category at site B.	233
Appendix 21. Biochemistry and haematology under 'optimal' seasonal conditions: means and medians for cow and herd bull groups at site B.	234
Appendix 22. Biochemical indicators of nutrition, per categories of financial years: 25 th percentile and median value of lowest ranked year, compared with reference range for cow and herd bull groups at sites B and C.	238

Appendix 23. Faecal chemistry values, per categories of 2-months cumulative rainfall:
median and quartile range for cow and steer groups at sites B and C. . 239

Appendix 24. Genera of intestinal nematode eggs and larvae in faecal samples at site B.
..... 241

Appendix 25. Faecal egg and oocyst counts, per categories of financial years: key data
distribution ranges for cow, herd bull and weaned-calf groups at site B.
..... 242

GENERAL ACKNOWLEDGMENTS

I would like to acknowledge and thank the people who have inspired this thesis.

The owners and managers of 'Hamilton Downs', 'Mount Riddock' and 'Narwietooma' pastoral stations in the Alice Springs district are acknowledged for their generosity and dedication, which enabled collection of research data for three primary beef cattle projects. Special mention is made of the Cadzow Family and their staff on 'Mount Riddock' station.

Present and former colleagues of the (now) Northern Territory Department of Primary Industry and Fisheries, Alice Springs, are thanked for their key contributions for the undertaking and reporting on two of the primary beef cattle projects (1991-1996, 1998-2001). Special mention is made of the contributions by Dr. Taffy (Owen) Williams, Mr. Bruce Tye, Mr. Keith Hill and Mr. Andrew Phillips.

Former CSIRO researchers, Dr. William Low and Dr. Jeff Wood, are thanked for the raw data and text in an unpublished manuscript from one primary beef cattle project (1971-1975). Dr. William Low and former staff of (now) Queensland Department of Agriculture, Fisheries and Forestry, Mr. John Bertram and Dr. Richard Holroyd, are all thanked for their constructive critiques for this thesis.

Finally but foremost, Prof. Ian Robertson and Dr. Eric Taylor, current and former Murdoch University academics, respectively, of the College of Veterinary Medicine in the School of Veterinary and Life Sciences, are thanked for their supervision and support. Together with the legacy of support from the late Dr. Diana Pinch and the study award from the (now) Northern Territory Department of Primary Industry and Fisheries, their time and patience has enabled completion of the thesis.

ABBREVIATIONS

Term	Abbreviation
Australian Bureau of Agricultural and Resource Economics and Sciences	ABARES
agar gel immunodiffusion	AGID
analysis of variance	ANOVA
animal (cattle) equivalents	AE
artificial insemination	AI
aspartate-A-transferase	AST
Australian agricultural and grazing industries survey	AAGIS
blood glutathione peroxidase	GSH-Px
body condition score	BCS
bovine herpes virus type 1	BHV1
bovine viral diarrhoea virus	BVDV
BVDV disease complex, including mucosal disease	BVD/MD
Brucellosis and Tuberculosis Eradication Campaign	BTEC
bull breeding soundness examination	BBSE
calcium	Ca
carbon 3 plant	C3 plant
chloride	Cl
copper	Cu
creatine phosphokinase	CPK
crude protein	CP
deoxyribonucleic acid	DNA
dry matter digestibility	DMD
eggs per gram (of faeces)	epg
enzyme-linked immunosorbent assay	ELISA
estimated breeding value	EBV
Faecal Near Infra-red Spectroscopy	F-NIRS
faecal percentage of dry matter	Faecal %DM
faecal percentage of nitrogen	Faecal %N
faecal percentage of phosphorus	Faecal %P
gamma glutamyl transferase	GGT
haemoglobin	Hb
infectious balanoposthitis	IBP
infectious bovine rhinotracheitis	IBR
infectious pustular vulvovaginitis	IPV
iron	Fe
magnesium	Mg
manganese	Mn
micro-agglutination test	MAT
nitrogen	N
Northern Territory Government	NTG
oocysts per gram (of faeces)	opg
persistently infected cattle (with BVDV)	PI cattle
phosphorus	P
polymerase chain reaction (amplification assay)	PCR
potassium	K

Term	Abbreviation
Rangeland Condition Assessment	RCA
red blood cells	RBC
selenium	Se
serum neutralisation	SN
sodium	Na
sulphur	S
United States of America	USA
zinc	Zn
95% confidence interval	95% CI