



**Murdoch**  
UNIVERSITY

**MURDOCH RESEARCH REPOSITORY**

*This is the author's final version of the work, as accepted for publication following peer review but without the publisher's layout or pagination.*

*The definitive version is available at*

<http://dx.doi.org/10.1007/s10340-014-0566-6>

**Barton, A.F.M., Clarke, B.R., Dell, B. and Knight, A.R. (2014)  
Post-emergent herbicidal activity of cineole derivatives. *Journal of Pest Science*, 87 (3). pp. 531-541.**

<http://researchrepository.murdoch.edu.au/21001/>

Copyright: © 2014 Springer-Verlag Berlin Heidelberg.

It is posted here for your personal use. No further distribution is permitted.

## Post-emergent Herbicidal Activity of Cineole Derivatives

Allan F. M. Barton<sup>a</sup>, Brenton R. Clarke<sup>b</sup>, Bernard Dell<sup>a</sup>, and Allan R. Knight<sup>\*a</sup>

<sup>a</sup>School of Veterinary and Life Sciences, Murdoch University, South Street, Murdoch, Western Australia, 6150.

<sup>b</sup>School of Engineering and Information Technology, Murdoch University, South Street, Murdoch, Western Australia, 6150;

\*Author to whom correspondence should be addressed (telephone: 61 08 9360 6871; fax 61 08 9360 6452; e-mail [a.knight@murdoch.edu.au](mailto:a.knight@murdoch.edu.au))

### Abstract

Essential oils are being investigated as potential herbicides or to provide leads to new environmentally and socially acceptable herbicides. Novel hydroxy and ester derivatives of 1,8-cineole and 1,4-cineole were synthesised, by chemical and biological methods, and have shown pre-emergence herbicidal activity against annual ryegrass and radish. Effects on post-emergence activity of these derivatives, as well as 1,8-cineole, eucalyptus oil and the carboxylic acids from which the esters were derived, against annual ryegrass and radish, are reported here. Results suggest that reduced root and shoot growth observed in pre-emergence herbicidal bioassays were due to post-emergence activity rather than delayed germination. All tested substances had a dose-dependent, post-emergence herbicidal activity against annual ryegrass and radish with many derivatives showing improved activity relative to 1,8-cineole and high-cineole eucalyptus oil. However, results do not support the postulate that cineole esters would be more active than their respective carboxylic acid and the hydroxy

26 cineole. Phytotoxicity of ester derivatives may be due to metabolic cleavage of the  
27 esters to the hydroxy cineole and carboxylic acid within the plant.

28

29 **Keywords:** Eucalyptus oil, cineole, hydroxy cineole, monoterpenes, phytotoxicity.

30

## 31 **Introduction**

32 The decrease in crop yields in agricultural systems and reduction in biodiversity in  
33 natural areas of vegetation due to weeds requires a range of strategies to reduce their  
34 impacts. Prior to the mid-twentieth century, strategies to manage weeds were  
35 primarily non-chemical, categorised as either mechanical or cultural (Kohli et al.  
36 2006). Mechanical methods such as harrowing and inter-row hoeing together with  
37 cultural methods like fertilizer placement, seed vigour, seeding rate and competitive  
38 varieties provide encouraging results but long-term solutions require consideration at  
39 the cropping system level. In low-external input or ecologically sustainable farming  
40 practices weed control may be most effectively approached by examining interactions  
41 among system components and agricultural practices occurring in the current crop,  
42 the subsequent crop and between crops (Bàrberi 2002; Melander et al. 2005). For  
43 example, the slower release of nutrients from organic fertilisers compared to synthetic  
44 fertilisers delays weed emergence in turn leading to competition between the crop and  
45 weeds occurring later and with the potential for effects to carry over to the next  
46 growing season and likely causing changes in the weed community (Bastiaans and  
47 Drenth 1999; Liebman 2000; McCloskey et al. 1996). These approaches are  
48 ecologically sustainable but can be labour intensive and may not provide the level of  
49 weed control needed to give sufficient crop yields to feed the world's population  
50 (Gianessi 2009). As well, as understanding of ecosystems has developed, biological  
51 control agents have been used but they are usually targeted at a particular species,  
52 they can be slow to give an acceptable level of control and usually they are not  
53 applicable to cropping systems.

54 Thus for cropping systems it may be necessary to use chemical weed control  
55 to reduce the damage to an acceptable level but herbicide applications have

56 drawbacks including loss of non-target organisms from microbes to vertebrates due to  
57 toxic soil and water residues which can cause indirect ecological problems (Lewis et  
58 al. 2009; Wang et al. 2010; Willemsen and Hailey 2001), possible impacts on human  
59 health from residues in foods exposed to these chemicals (Chade et al. 2006; Fantke et  
60 al. 2012; Rouimi et al. 2012) and development of herbicide resistance in weed species  
61 (Heap 2007).

62 Plant-sourced compounds that exhibit phytotoxicity, and so may act as  
63 herbicides or provide leads to novel herbicides, have the potential to avoid some of  
64 these problems. Their half-lives are likely to be relatively short thus reducing  
65 problems of residues in food and water, and they may have novel modes of action  
66 thus potentially addressing problems of resistance to current herbicides (Duke et al.  
67 2002).

68 Essential oils and their constituents, including monoterpenes, are plant  
69 secondary metabolites being increasingly studied for their allelopathic, herbicidal,  
70 insecticidal, acaracidal and other biological activities. The varied ecological roles of  
71 monoterpenes include reduced susceptibility to insect herbivory, attractants for  
72 pollinators (Ibanez et al. 2010) and suppression of germination of competing plants.  
73 There are many reports of the insecticidal activity of essential oils and their  
74 monoterpene constituents, including 1,8-cineole. Edwards et al. (1993) and Matsuki  
75 et al. (2011) observed that increased 1,8-cineole content in leaves reduced herbivory  
76 by Christmas beetles. 1,8-Cineole was found to be highly toxic to the grain beetles  
77 *Prostephanus truncates*, *Sitophilus granarius*, *S. zeamais* and *Tribolium castaneum*  
78 with 100% mortality at 0.5  $\mu$ L 1,8-cineole per kilogram of grain after 24 hours  
79 (Obeng-Ofori et al. 1997). Polatoğlu (2013) also reported toxicity of the essentials  
80 oils of *Achillea* species, which have high 1,8-cineole content, to *S. granarius*. 1,8-

81 Cineole and other monoterpenes have shown toxicity and feeding and oviposition  
82 deterrence to the moth species *Helicoverpa armigera* (Hübner), *Spodoptera litura* (F.)  
83 and *Chilo partellus* Swinhoe (Koul et al. 2013). The fumigant activity of eucalyptus  
84 oils from a number of *Eucalypt* species on adult *Aedes aegypti*, the yellow fever  
85 mosquito, correlates with 1,8-cineole content of the oils (Lucia et al 2009) and  
86 Lampman et al. (2000) showed mosquito larvicidal activity for 1,8-cineole.  
87 Acaricidal activity of essential oils containing 1,8-cineole has been demonstrated with  
88 *Rosmarinus officinalis* L., *Salvia officinalis* and Myrtaceae essential oils active  
89 against the two-spotted mite *Tetranychus urticae* (Laborda et al. 2013; Miresmailli et  
90 al. 2006; Roh et al. 2013). As well, a terpene-based solution containing 1,8-cineole  
91 gave a mortality of 96.7% against the western honey bee parasitic mite *Varroa*  
92 *jacobsonii* (Calderone and Spivak 1995).

93 A wide range of classes of volatile monoterpenes inhibit plant growth (Amri et al.  
94 2012; Apsland 1968; Chaimovitsh et al. 2011; Muller and Muller 1964; Vaughn and  
95 Spencer 1993). For example, *Artemisia frigida* has been shown to have inhibiting  
96 effects on plant communities in the steppe of northern China with its leaf volatile  
97 components comprising mainly monoterpenes, including 1,8-cineole (Li et al. 2011).  
98 The high-1,8-cineole essential oil of *Eucalyptus erythrocorys* has demonstrated  
99 herbicidal activity against *Sinapis arvensis* L. and *Phalaris canariensis* L. preventing  
100 seed germination at oil concentration of 1.5  $\mu\text{L mL}^{-1}$  and delaying and decreasing  
101 germination at lower concentrations (Ben Ghnaya et al. 2013).

102 In the field, the volatility of monoterpenes may limit plant uptake and  
103 therefore their effectiveness. There are reports on attempts to address volatility  
104 through microencapsulation or nano-formulation of essential oils for the purposes of  
105 both herbicidal and insecticidal activity. Nanoparticles loaded with garlic essential oil

106 and *Mentha* oil nanoparticles have been shown to maintain activity against insects  
107 over an extended period (Yang et al. 2009; Kumar et al. 2013) and polyurea  
108 microcapsules containing essential oils reduced seed germination relative to controls  
109 although were not as effective as neat oils (Scarfato et al. 2007). However, there are  
110 few reports of attempts to reduce volatility whilst maintaining herbicidal activity by  
111 the synthesis of derivatives of monoterpenes. The derivatives have increased  
112 molecular mass to lower vapour pressure and give a slower evaporation rate compared  
113 to the parent monoterpene. Vaughn and Spencer (1996) prepared benzyl ether  
114 derivatives of a number of monoterpenes for subsequent herbicidal testing but the  
115 only report of the synthesis of ester derivatives of monoterpenes such as 1,8-cineole  
116 for this purpose is for pre-emergence testing (Barton et al. 2010).

117         Whilst it is usually preferable to apply herbicides before crops emerge, weeds  
118 can emerge to compete with crops that are slow to germinate such as chickpeas or  
119 crops may be poor competitors, and so require weed control. Zero or reduced tillage  
120 in broad-acre farming has led to volunteer cereals growing with crops and so causing  
121 reduced yields (Friesen et al. 1990; O'Donovan 1992; Wilson et al. 2010). Weed  
122 control is necessary after crop germination where there are weed species that have  
123 lengthy germination periods, grow strongly in autumn and spring, produce large  
124 quantities of seed, or have a long lasting seed bank.

125         An aim of this work was to assess and compare post-emergent herbicidal  
126 activity of eucalyptus oil, 1,8-cineole, the major component in the leaf oil of many  
127 eucalypts, and hydroxy and ester derivatives of 1,8-cineole and 1,4-cineole. The  
128 study investigated whether observed reductions in root and shoot growth of annual  
129 ryegrass (*Lolium rigidum*) and radish (*Raphanus sativus* var. Long Scarlet) when  
130 seeds were treated pre-emergence with these substances (Barton et al. 2010) was due

131 to delayed germination or reductions in growth. It was also postulated that on uptake  
132 by plants the cineole esters may undergo metabolic cleavage to give the hydroxyl-  
133 cineole and the carboxylic acid and so any herbicidal activity of the esters may, in  
134 fact, be due to the hydroxyl-cineole and its carboxylic acid. Thus the post-emergence  
135 herbicidal activity of the carboxylic acids corresponding to the esters was also  
136 assessed. For the purposes of the work reported here, pre-emergence activity was  
137 defined as herbicidal activity preventing seed germination and post-emergent activity  
138 was defined as activity preventing or reducing further growth after emergence of the  
139 radicle and plumule.

#### 140 **Materials and Methods**

141 **Chemicals.** All chemicals were purchased from standard commercial suppliers.  
142 Eucalyptus oil (96% v/v 1,8-cineole) was obtained from Kalannie Distillers, Kalannie  
143 Western Australia. The oil was from *Eucalyptus kochii* subsp *horistes* and *Eucalyptus*  
144 *kochii* subsp. *Plenissima*.

145 **Synthesis of 1,8-Cineole and 1,4-Cineole Derivatives.** Cineole derivatives were  
146 prepared as described in Barton et al. (2010) and references therein.

147 **Seed Sources.** Annual ryegrass seeds (*Lolium rigidum*) were obtained from the  
148 Wongan Hills Research Station 2EA, Western Australia, in November 2002 and  
149 radish seeds (*Raphanus sativus* var. Long Scarlet) were a commercially available  
150 variety (Mr Fothergill's Seeds Pty Ltd).

151 **Seed Treatment.** Seeds were surface sterilised in 2% sodium hypochlorite solution  
152 for 10 minutes, rinsed 3 times with sterile deionised water and then imbibed for  
153 approximately 15 hours in sterile deionised water.

154 **Post-emergence Bioassays.** Substances assessed for post-emergence activity were  
155 1,8-cineole **1**, eucalyptus oil, 3-oxo-1,8-cineole **2**; the hydroxylated cineole



156 compounds **3**, **4a** and **5a**; 1,8-cineole esters **4b-d**; and 1,4-cineole esters **5b** and **c**  
157 (Figure 1). The cineole esters assessed for post-emergence activity were selected on  
158 the basis of their activity in pre-emergence testing (Barton et al. 2010). Carboxylic  
159 acids assessed in these post-emergence bioassays were those corresponding to the  
160 esters. Seeds were germinated on sterile water agar in Petri dishes before transfer to  
161 the 55 mm Petri dishes that had been prepared with the test compounds. The radish  
162 seeds took 24 hours and the ryegrass seeds took 40 hours to germinate when  
163 incubated at 25 °C. The water agar was prepared by autoclaving (103.4 kPa, 121 °C,  
164 30 minutes) 4.0 g of agar (BBL™ Agar, Grade A) in 500 mL of deionised water  
165 containing calcium (0.05 mol L<sup>-1</sup>) and boron (0.001 mol L<sup>-1</sup>). Petri dishes (55 mm  
166 plastic) (or Pyrex dishes for chloroform solutions) for the post-emergence bioassays  
167 were prepared, under sterile conditions, by pouring the agar into them to a depth of  
168 approximately 2 mm and allowing them to solidify. A solution (1 mL) of the test  
169 compound in the required solvent (Table 1) was introduced into the Petri dish using a  
170 micropipette and the dish left open in a laminar flow cabinet for 3 hours to allow  
171 evaporation of the organic solvent. The concentrations for these post-emergence  
172 bioassays ranged from concentrations where seedlings showed little or no response to  
173 those with complete or nearly complete mortality in pre-emergence bioassays (Barton  
174 et al. 2010) (Table 2).

175 Filter paper bioassays were used for 1,8-cineole and eucalyptus oil. Filter  
176 papers (Whatman number 4) were autoclaved, oven dried and placed into autoclaved  
177 pyrex Petri dishes (55 mm) under sterile conditions. 1,8-Cineole solution or  
178 eucalyptus oil solution (1 mL) was transferred on to the filter paper using a  
179 micropipette, the lid placed on the Petri dish and the dish sealed with plastic food  
180 wrap. The 1,8-cineole and eucalyptus oil were prepared in aqueous solution with 0.34

181 mg mL<sup>-1</sup> of the non-ionic surfactant Tween<sup>®</sup> 80, polyoxyethylene (20) sorbitan  
182 monooleate. The deionised water/Tween<sup>®</sup> 80 solution, containing calcium and boron  
183 as above, was autoclaved prior to preparation of the 1,8-cineole and eucalyptus oil  
184 solutions. All glassware used in the preparation of these solutions was washed with  
185 2% sodium hypochlorite solution and then rinsed with sterile deionised water.  
186 Ten seedlings were placed in each Petri dish and then sealed with plastic food wrap.  
187 The Petri dishes were placed randomly in a tray with Styrofoam supports to angle the  
188 dishes at approximately 70° to the horizontal to encourage straighter root and shoot  
189 growth. Petri dishes with 1,8-cineole and eucalyptus oil were placed flat. The tray  
190 was incubated under light (135 to 195  $\mu\text{E m}^{-2} \text{s}^{-1}$  photosynthetic active radiation) at  
191 25 °C for 48 hours. Prior to measurement of the increase in radish root and shoot  
192 lengths, seedlings were frozen in their Petri dishes and thawed. This softened their  
193 roots and shoots making easier measurement of their lengths. Ryegrass shoots were  
194 too fragile to be frozen and thawed.

195 Two controls, one with and one without solvent, were used for each  
196 experiment. For solvent controls, solvent (1 mL) was pipetted on to the surface of the  
197 agar and the Petri dish left open in a laminar flow cabinet for three hours. The non-  
198 solvent control consisted of the same agar solution in Petri dishes that were similarly  
199 left open in a laminar flow cabinet for three hours.

200 **Experimental Design and Data Analysis.** Five replicates were used at each  
201 concentration and for controls. Petri dishes were placed in a randomised manner in  
202 the support tray. Each experiment was repeated in duplicate with two-tailed t-tests  
203 showing no significant difference between repeats at  $P = 0.05$ . Data for effects of  
204 concentration on increased root and shoot growth were subjected to one way analysis  
205 of variance (ANOVA) using the SPSS 15.0 statistics package (SPSS Inc., 2007).

206 Differences between means were tested using Tukey's HSD test and were considered  
207 to be statistically different at  $P < 0.05$ . Modelling dose response data using non-linear  
208 log-logistic regression analysis to fit it to a sigmoidal curve to determine the  $I_{50}$  (50%  
209 inhibition) values for root growth and shoot growth was carried out as described by  
210 Seefeldt et al. (1995).

## 211 **Results**

212 These post-emergence bioassays showed that for all tested substances the radish and  
213 ryegrass had a dose response with inhibition of root and shoot growth increasing with  
214 concentration (Figures 2, 3, 4 and 5). As well, for both plant species the post-  
215 emergence bioassays confirmed that reduced root and shoot growths observed in pre-  
216 emergence bioassays (Barton et al. 2010) were due to growth inhibition rather than  
217 germination delay.

218 **Acids on Radish.** Acetic acid suppressed radish root growth and shoot growth at and  
219 above  $0.01 \text{ mol L}^{-1}$  (Figure 2 (a)). Radish roots were more sensitive to benzoic acid  
220 than were the shoots with suppression of root growth first observed at  $0.000316 \text{ mol}$   
221  $\text{L}^{-1}$  and for shoot growth at  $0.001 \text{ mol L}^{-1}$  (Figure 2 (c)). Hexanoic acid inhibited  
222 radish root growth at and above  $0.0025 \text{ mol L}^{-1}$  (Figure 2 (e)). At the highest  
223 concentration of  $0.05 \text{ mol L}^{-1}$ , hexanoic acid reduced root growth by approximately  
224 99%. Radish shoot growth was reduced by hexanoic acid at  $0.007 \text{ mol L}^{-1}$  reaching  
225 87% reduction compared to the mean of the control at  $0.05 \text{ mol L}^{-1}$  (Figure 2 (e)).

226 **1,8-Cineole and Eucalyptus Oil on Radish.** Suppression of roots and shoots by 1,8-  
227 cineole **1** was significant at and above  $0.1 \text{ mol L}^{-1}$  (Figure 2 (b)). Eucalyptus oil  
228 suppressed growth of roots at and above  $0.01 \text{ g mL}^{-1}$  and shoots above  $0.0316 \text{ g mL}^{-1}$   
229 (Figure 2 (f)). Roots turned brown and became dehydrated when exposed to 1,8-

230 cineole **1** or eucalyptus oil at their highest concentrations, and no new root growth  
231 occurred.

232 **1,8-Cineole Derivatives on Radish.** 3-Oxo-1,8-cineole **2** suppressed shoot growth at  
233 and above 0.01 mol L<sup>-1</sup> whilst root growth was reduced above 0.025 mol L<sup>-1</sup> (Figure  
234 2 (d)). There was complete inhibition of radish root and shoot growth by 2-*endo*-  
235 hydroxy-1,8-cineole **3** at 0.2 mol L<sup>-1</sup>, the highest concentration tested, as well as  
236 significant (for root  $P = 1.14 \times 10^{-12}$ , for shoot  $P = 3.41 \times 10^{-3}$ ) suppression above  
237 0.05 mol L<sup>-1</sup> (Figure 3 (a)). At the highest concentration, 2-*endo*-hydroxy-1,8-cineole  
238 **3** caused browning at the root tip. 3-*exo*-Hydroxy-1,8-cineole **4a** suppressed ( $P =$   
239  $1.78 \times 10^{-6}$ ) radish root growth at and above 0.01 mol L<sup>-1</sup> whilst it only suppressed  
240 shoot growth at and above 0.1 mol L<sup>-1</sup> (Figure 2 (c)). Radish shoots treated with 3-  
241 *exo*-hydroxy-1,8-cineole **4a** were clearly lighter green than shoots of the control  
242 seedlings. Shoot suppression ( $P = 1.87 \times 10^{-4}$ ) by 3-*exo*-benzoxy-1,8-cineole **4b** was  
243 seen at and above 0.01 mol L<sup>-1</sup> whilst root suppression ( $P = 3.05 \times 10^{-3}$ ) occurred  
244 above 0.0316 mol L<sup>-1</sup> (Figure 2 (e)).

245 **1,4-Cineole Derivatives on Radish.** 2-*exo*-Hydroxy-1,4-cineole **5a** suppressed  
246 further radish root growth at all the concentrations tested, with growth reducing to 2%  
247 of the control mean at 0.1 mol L<sup>-1</sup> (Figure 3 (b)). This compound only depressed  
248 further shoot growth at and above 0.04 mol L<sup>-1</sup> (Figure 3 (b)). The 2-*exo*-hydroxy-  
249 1,4-cineole **5a** caused browning of the radish root tips at 0.1 mol L<sup>-1</sup>. 2-*exo*-Acetoxy-  
250 1,4-cineole **5b** inhibited root and shoot growth only at the highest tested concentration  
251 of 0.1 mol L<sup>-1</sup> (Figure 3 (d)). 2-*exo*-Acetoxy-1,4-cineole, as for 3-*exo*-hydroxy-1,8-  
252 cineole, caused shoots to be paler green than shoots of control seedlings. Although 2-  
253 *exo*-hexoxy-1,4-cineole **5c** suppressed root and shoot growth at all tested

254 concentrations it did not completely inhibit further root or shoot growth even at the  
255 highest tested concentration (Figure 3 (f)).

256 **Acids on Ryegrass.** Acetic acid suppressed post-emergent ryegrass root growth at  
257 and above  $0.0025 \text{ mol L}^{-1}$  with root length decreasing to about 6% of the control  
258 mean at  $0.05 \text{ mol L}^{-1}$  (Figure 4 (a)). Shoot growth was suppressed by acetic acid  
259 above  $0.01 \text{ mol L}^{-1}$  (Figure 4 (a)). *t*-Butylacetylacetic acid suppressed root growth at  
260  $0.001 \text{ mol L}^{-1}$  ( $P = 3.55 \times 10^{-13}$ ) with complete inhibition of growth at the highest  
261 concentration of  $0.0316 \text{ mol L}^{-1}$  (Figure 4 (c)). Shoots were first suppressed at  
262  $0.00316 \text{ mol L}^{-1}$  by benzoic acid (Figure 4 (c)). Hexanoic acid suppressed root and  
263 shoot growth at and above  $0.0025 \text{ mol L}^{-1}$  (Figure 4 (e)). This acid completely  
264 stopped further root growth at  $0.02 \text{ mol L}^{-1}$  and shoot growth at  $0.05 \text{ mol L}^{-1}$  (Figure  
265 4 (e)).

266 **1,8-Cineole and Eucalyptus Oil on Ryegrass.** 1,8-Cineole **1** stopped ryegrass root  
267 and shoot growth above  $0.1 \text{ mol L}^{-1}$  with root suppression first occurring at  $0.0316$   
268  $\text{mol L}^{-1}$  ( $P = 3.55 \times 10^{-13}$ ) and shoot suppression at  $0.1 \text{ mol L}^{-1}$  (Figure 4 (b)).  
269 Eucalyptus oil suppressed ryegrass root and shoot growth above  $0.00316 \text{ g mL}^{-1}$  and  
270 completely inhibited root growth above  $0.01 \text{ g mL}^{-1}$  and shoot growth above  $0.0316 \text{ g}$   
271  $\text{mL}^{-1}$  (Figure 4 (f)).

272 **1,8-Cineole Derivatives on Ryegrass.** 3-Oxo-1,8-cineole **2** reduced ryegrass root  
273 growth at  $0.005 \text{ mol L}^{-1}$  whilst shoot growth was decreased above  $0.0025 \text{ mol L}^{-1}$   
274 (Figure 4 (d)). 2-*endo*-Hydroxy-1,8-cineole **3** reduced root growth above  $0.005 \text{ mol}$   
275  $\text{L}^{-1}$ , with complete inhibition of root growth above  $0.1 \text{ mol L}^{-1}$  (Figure 5 (a)). 2-  
276 *endo*-Hydroxy-1,8-cineole **3** suppressed shoot growth above  $0.01 \text{ mol L}^{-1}$  (Figure 5  
277 (a)). 3-*exo*-Hydroxy-1,8-cineole **4a** suppressed root growth at and above  $0.025 \text{ mol}$   
278  $\text{L}^{-1}$  leading to complete inhibition above  $0.1 \text{ mol L}^{-1}$  (Figure 5 (b)). This hydroxy

279 compound suppressed ryegrass shoot growth above  $0.01 \text{ mol L}^{-1}$  (Figure 5 (b)). 3-  
280 *exo*-Hexoxy-1,8-cineole **4c** completely inhibited ryegrass root growth at and above  
281  $0.0316 \text{ mol L}^{-1}$  with suppression first observed at  $0.01 \text{ mol L}^{-1}$  ( $P = 6.47 \times 10^{-3}$ ) but  
282 promoted root growth at the two lowest concentrations tested of  $0.001$  and  $0.00316$   
283  $\text{mol L}^{-1}$  ( $P = 2.44 \times 10^{-5}$ , and  $P = 2.93 \times 10^{-6}$ , respectively) (Figure 5 (c)). This  
284 compound suppressed ryegrass shoot growth at all concentrations with complete  
285 inhibition at the highest concentration ( $0.1 \text{ mol L}^{-1}$ ) (Figure 5 (c)). 3-*exo-t*-  
286 Butylacetoxy-1,8-cineole **4d** reduced root and shoot growth at and above  $0.01 \text{ mol L}^{-1}$   
287 with roots showing 5% and shoots 27% growth relative to control means at  $0.316 \text{ mol}$   
288  $\text{L}^{-1}$  (Figure 5 (e)).

289 **1,4-Cineole Derivatives on Ryegrass.** 2-*exo*-Hydroxy-1,4-cineole **5a** and 2-*exo*-  
290 acetoxy-1,4-cineole first suppressed ryegrass root and shoot growth at  $0.01 \text{ mol L}^{-1}$   
291 with complete root growth inhibition for both at  $0.1 \text{ mol L}^{-1}$  (Figure 5 (d) and (f)).

292 The ryegrass roots turned brown and became dehydrated when treated with  
293 1,8-cineole **1**, eucalyptus oil, 2-*endo*-hydroxy-1,8-cineole **3**, 3-*exo*-hexoxy-1,8-cineole  
294 **4c**, 2-*exo*-hydroxy-1,4-cineole **5a** and 2-*exo*-acetoxy-1,4-cineole **5b** at their highest  
295 tested concentrations. The browning of the roots was apparent within approximately  
296 5 minutes for the 1,8-cineole and eucalyptus oil.

297 The dose response data for both species closely fitted the sigmoidal curves  
298 generated from log-logistic regression analysis ( $R^2$  values all above 0.9) but the  $I_{50}$   
299 values are approximate with errors in some cases larger than the estimated  $I_{50}$  due to  
300 emphasis being on the wide range of compounds tested rather than on repetition to  
301 achieve high precision for fewer compounds.

302 **Discussion**

303 As for results of germination bioassays (Barton et al. 2010), the post-emergence  
304 results do not support the postulate that cineole esters would be more active than their  
305 respective carboxylic acid and the hydroxy cineole due to metabolic cleavage on  
306 uptake by plants. In general the post-emergence activity of the cineole esters did not  
307 show improvement relative to their respective hydroxylated cineole and carboxylic  
308 acid precursors. Eucalyptus oil was compared to the 1,8-cineole **1** effects to give an  
309 indication of any effects other components of the oil may have on growth of  
310 seedlings. The results suggest limited effect on growth of other components of the  
311 oil.

312 The post-emergent results indicate that for radish, roots were generally more  
313 sensitive to the tested substances than were shoots, as also shown by the pre-  
314 emergence observations. For the ryegrass, shoots were slightly more sensitive post-  
315 emergent but there was no clear trend for sensitivity of roots as compared to shoots  
316 for pre-emergence bioassays.

317 Post-emergence, the carboxylic acids were the most active of the tested  
318 substances against radish with 2-*endo*-hydroxy-1,8-cineole **3** as the overall most  
319 active of the cineole compounds. Although both 3-*exo*-benzoxy-1,8-cineole **4b** and 2-  
320 *exo*-hexoxy-1,4-cineole **5c** initially suppress shoot growth at a lower concentration  
321 than 2-*endo*-hydroxy-1,8-cineole **3**, they do not completely inhibit shoot growth even  
322 at 1 mol L<sup>-1</sup> whilst 2-*endo*-hydroxy-1,8-cineole completely inhibits shoots at 0.2 mol  
323 L<sup>-1</sup>. Several cineole compounds initially suppress root growth at a concentration  
324 lower than that of 2-*endo*-hydroxy-1,8-cineole but some of these compounds do not  
325 give complete root inhibition whilst 2-*endo*-hydroxy-1,8-cineole does.

326 Of all the cineole compounds 3-*exo*-hexoxy-1,8-cineole **4c** had the highest  
327 post-emergence activity against ryegrass with a shoot growth suppression initially

328 occurring at the lowest concentration of the cineole compounds and with complete  
329 suppression at  $0.1 \text{ mol L}^{-1}$ . Whilst other cineole compounds suppressed post-  
330 emergent ryegrass root growth at lower concentrations, this hexanoate ester  
331 completely inhibited root growth at  $0.0316$  and  $0.1 \text{ mol L}^{-1}$ . *2-endo*-Hydroxy-1,8-  
332 cineole **3** was the most active hydroxy-cineole against ryegrass roots but all the  
333 hydroxy-cineoles had similar shoot activity.

334         The lighter green of radish shoots treated with *3-exo*-hydroxy-1,8-cineole  
335 compared to that of shoots of control seedlings indicates this compound may interfere  
336 with chlorophyll production or enhance its breakdown, or increase production of  
337 masking carotenoids. Sing et al. (2002) observed that 1,8-cineole reduced chlorophyll  
338 content in billy goat weed as well as reducing cellular respiration. The content of  
339 chlorophylls a and b in *Amaranthus viridis* were observed to decrease on treatment  
340 with 1,8-cineole, as was the amount of carotenoids (Kaur et al. 2011) suggesting that  
341 the lighter green of radish shoots is more likely a result of lowered chlorophyll  
342 production rather than presence of masking carotenoids. Kaur et al. (2011) also  
343 observed lowered cell respiration in *A. viridis*. The cineole-containing oil of *Ajania*  
344 *tenuifolia* caused decreased activity of nitrate reductase and chlorophyll content of  
345 *Elymus nutans* (Bai and Zhang 1994). There are many other reports of eucalyptus oils  
346 and 1,8-cineole reducing chlorophyll content (Batish et al. 2004; Kohli and Singh  
347 1991; Singh et al. 2005). Reduced chlorophyll content will lower photosynthetic  
348 efficiency and so contribute to the herbicidal activity of 1,8-cineole. The browning of  
349 roots by 1,8-cineole and eucalyptus oil was likely as a result of the volatility of these  
350 substances. Reduced root growth may be a result of inhibition of DNA synthesis in  
351 nuclei and other organelles in the root apical meristem. 1,8-Cineole has been shown  
352 to decrease the DNA synthesis activity in the root tips of *Brassica campestris*



353 (Koitabashi et al. 1997; Nishida et al. 2005) and to inhibit all stages of mitosis in  
354 onion roots (Romagni et al. 2000). The cineole derivatives prepared in this work have  
355 higher melting points and lower volatility at ambient temperatures than 1,8-cineole, so  
356 overcoming limitations in field use of 1,8-cineole as a herbicide due to its volatility  
357 and subsequent low uptake by plants.

### 358 **Conclusion**

359 In conclusion, there is a dose-dependent post-emergence herbicidal activity by 1,8-  
360 cineole and the hydroxy and ester derivatives of 1,8-cineole and 1,4-cineole against  
361 radish and annual ryegrass root and shoot growth. *2-endo*-Hydroxy-1,8-cineole **3** is  
362 the most active of the cineole derivatives against radish and *3-exo*-hexoxy-1,8-cineole  
363 **4c** the most active derivative against ryegrass. Many derivatives have improved  
364 phytotoxicity relative to 1,8-cineole, particularly at the lower concentrations. Results  
365 do not indicate any strong improvement in activity of 1,8-cineole derivatives over 1,4-  
366 cineole derivatives. As in the case of pre-emergence bioassays, in this study the  
367 carboxylic acids were more active and observed phytotoxicity of ester derivatives may  
368 be due to metabolic cleavage of the esters to the hydroxy cineole and carboxylic acid  
369 within the plant. Based on these preliminary results, the hydroxyl and ester  
370 derivatives of 1,8-cineole and 1,4-cineole have potential as herbicides. However,  
371 before further investigation into their potential as herbicides is undertaken it may be  
372 most appropriate to assess their mechanism of phototoxicity. A novel mechanism of  
373 action may provide stimulus to the development of these potentially safer compounds  
374 but research to assess their efficacy in field trials, toxicity against the crop plants that  
375 they might be used for and safety would be needed. Structure-activity studies to  
376 compare *3-endo*-hydroxy-1,8-cineole and *3-exo*-hydroxy-1,8-cineole may also

377 provide clearer understanding of the position and stereochemical role of  
378 hydroxylation of the 1,8-cineole cyclohexane ring.

379 **Acknowledgement**

380 This work was supported by the Rural Industries Research and Development  
381 Corporation of the Australian Government.

382

383 **References**

- 384 Amri I, Gargouri S, Hamrouni L, Hanana M, Fezzani T, Jamoussi B (2012) Chemical  
385 composition, phytotoxic and antifungal activities of *Pinus pinea* essential oil. J  
386 Pest Sci 85:199-207
- 387 Apsland RO (1968) Monoterpenes: relationship between structure and inhibition of  
388 germination. Phytochem 7:1995-1997
- 389 Bai XF, Zhang BC (1994) Study of staple composition of *Ajania tenuifolia* oil on  
390 inhibition mechanism of seedling initial growth of *Elymus nutans*. Acta Ecol  
391 Sin 14:223-224
- 392 Bärberi P (2002) Weed management in organic agriculture: are we addressing the  
393 right issues? Weed Res 42:177-193
- 394 Batish DR, Setia N, Singh HP, Kohli RK (2004) Phytotoxicity of lemon-scented  
395 eucalyptus oil and its potential use as a bioherbicide. Crop Prot 23:1209-1214
- 396 Barton AFM, Dell B, Knight AR (2010) Herbicidal Activity of Cineole Derivatives. J  
397 Food Agric Chem 58:10147-10155. DOI: 10.1021/jf101827v
- 398 Bastiaans L, Drenth H (1999) Late-emerging weeds; phenotypic plasticity and  
399 contribution to weed growth., in: D. Gut (Ed.), 11th Symposium, Eur Weed  
400 Res Soc, Biological Control of Weeds, Wageningen, Netherlands : EWRS,  
401 Basel, Switzerland. pp. 3
- 402 Ben Ghnaya A, Hanana M, Amri I, Hazar B, Gargouri S, Jamoussi B, Hamrouni L  
403 (2013) Chemical composition of *Eucalyptus erythrocorys* essential oils and  
404 evaluation of their herbicidal and antifungal activities J Pest Sci 86:571-577
- 405 Calderone NW, Spivak M (1995) Plant Extracts for the Control of the Parasitic Mite  
406 *Varroa jacobsonii* (Acari: Varroidae) in Colonies of the Western Honey Bee  
407 (Hymenoptera: Apidea). J of Econ Entomol 88:1211-1215

- 408 Chade AR, Kasten M, Tanner CM (2006) Nongenetic causes of Parkinson's disease. J  
409 Neural Transm, Supplement 70:147-151
- 410 Chaimovitsh D, Rogovoy O, Altshuler O, Belausov E, Abu-Abied M, Rubin B, Sadot  
411 E, Dudai N (2011) The relative effect of citral on mitotic microtubules in  
412 wheat roots and BY2 cells. Plant Biol 14:354-364
- 413 Duke SO, Dayan F, Rimando AM, Schrader KK, Aliotta G, Oliva A, Romagni JG  
414 (2002) Chemicals from nature for weed management. Weed Sci 50:138-151
- 415 Edwards PB, Wanjura WJ, Brown WV (1993) Selective herbivory by Christmas  
416 beetles in response to intraspecific variation in *Eucalyptus* terpenoids  
417 Oecologia 95:551
- 418 Fantke P, Friedrich R, Jolliet O (2012) Health impact and damage cost assessment of  
419 pesticides in Europe. Environ Int 49:9-17
- 420 Friesen L, Morrison IN, Marshall G, Wesley R (1990) Effects of volunteer wheat and  
421 barley on the growth and yield of flax. Canadian J Plant Sci 70:1115-1122
- 422 Gianessi L (2009) The Potential for Organic Agriculture to Feed the World is Being  
423 Oversold. Outlooks on Pest Manag 20:4-5
- 424 Heap I (2007) The International Survey of Herbicide Resistant Weeds. Online.  
425 Internet. Available at [www.weedscience.com](http://www.weedscience.com)
- 426 Ibanez S, Dotterl S, Anstett MC, Baudino S, Caissair JC, Gallet C, Despres L (2010)  
427 The role of volatile organic compounds, morphology and pigments of  
428 globeflowers in the attraction of their specific pollinating flies. New Phytol  
429 188:451-463
- 430 Kaur S, Singh HP, Batish DR, Kohli RK (2011) Chemical characterization and  
431 allelopathic potential of *Eucalyptus tereticornis* against *Amaranthus viridis*. J  
432 Plant Interact 6:297-302

- 433 Kohli RK, Singh D (1991) Allelopathic impact of volatile components from  
434 *Eucalyptus* on crop plants. Biol Plant 33:475-483
- 435 Kohli RK, Batish DR, Singh HP (2006) Weeds and Their Management: Rationale  
436 and Approaches, In: Singh H P, et al. (Eds.), Handbook of Sustainable Weed  
437 Management, Haworth Press, Inc., New York. pp. 1-19
- 438 Koitabashi R, Suzuki T, Kawazu T, Sakai A, Kuroiwa H, Kuroiwa T (1997) 1,8-  
439 Cineole Inhibits Root Growth and DNA Synthesis in the Root Apical  
440 Meristem of *Brassica campestris* L. J Plant Res 110: 1-6
- 441 Koul O, Singh R, Kaur B, Kanda D (2013) Comparative study on the behavioural  
442 response and acute toxicity of some essential oil compounds and their binary  
443 mixtures to larvae of *Helicoverpa armigera*, *Spodoptera litura* and *Chilo*  
444 *partellus*. Ind Crops Prod 49:428-436
- 445 Kumar P, Mishra S, Malik A, Satya S (2013) Preparation and characterization of  
446 PEG-*Mentha* oil nanoparticles for housefly control. Colloids and Surf B:  
447 Biointerfaces In Press, Corrected Proof, Available online 15 November 2013
- 448 Laborda R, Manzano I, Gamón M, Gavidia I, Pérez-Bermúdez P, Boluda R (2013)  
449 Effects of *Rosmarinus officinalis* and *Salvia officinalis* essential oils on  
450 *Tetranychus urticae* Koch (Acari: Tetranychidae). Ind Crops Prod 48:106-110
- 451 Lampman R, Eckenbach U, Seigler D, Novak R (2000) Laboratory evaluations of  
452 methylated soy oil and monoterpenes as mosquito larvicides. J Am Mosq  
453 Control Assoc 16:153-157
- 454 Lewis S, Brodie J, Bainbridge Z, Rohde K, Davis A, Masters B, Maughan M, Devlin  
455 M, Mueller J, Schaffelke B (2009) Herbicides: A new threat to the Great  
456 Barrier Reef. Environ Pollut 157:2470-2484

- 457 Li XF, Wang J, Huang D, Wang LX, Wang K (2011) Allelopathic potential of  
458 *Artemisia frigida* and successional changes of plant communities in the  
459 northern China steppe. *Plant Soil* 341:383-398
- 460 Liebman D (2000) Integration of soil, crop and weed management in low-external-  
461 input farming systems. *Weed Res* 40:27-47
- 462 Lucia A, Licastro S, Zerba E, Audino PG, Masuh H (2009) Sensitivity of *Aedes*  
463 *aegypti* adults (Diptera: Culicidae) to the vapours of *Eucalyptus* essential oils.  
464 *Bioresour Technol* 100:6083-6087
- 465 Matsuki M, Foley WJ, Floyd RB (2011) Role of Volatile and Non-Volatile Plant  
466 Secondary Metabolites in Host Tree Selection by Christmas Beetles. *J Chem*  
467 *Ecol* 37:286-300
- 468 McCloskey M, Firbank LG, Watkinson AR, Webb DJ (1996) The dynamics of  
469 experimental arable weed communities under different management practices.  
470 *J Veg Sci* 7:799-808
- 471 Melander B, Rasmussen IA, Bàrberi P (2005) Integrating Physical and Cultural  
472 Methods of Weed Control: Examples from European Research. *Weed Sci*  
473 53:369-381
- 474 Miresmailli S, Bradbury R, Isman IB (2006) Comparative toxicity of *Rosmarinus*  
475 *officinalis* L. essential oil and blends of its major constituents against  
476 *Tetranychus urticae* Koch (Acari: Tetranychidae) on two different host plants.  
477 *Pestic Manag Sci* 62:366-371
- 478 Muller WH, Muller CH (1964) Volatile growth inhibitors produced by *Salvia* species.  
479 *Bull Torrey Bot Club* 91:327-330
- 480 Nishida N, Tamotsu S, Nagata N, Saito C, Sakai, A (2005) Allelopathic effects of  
481 volatile monoterpenoids produced by *Salvia leucophylla*: Inhibition of cell

- 482 proliferation and DNA synthesis in the root apical meristem of *Brassica*  
483 *campestris* seedlings. J Chem Ecol 31:1187-1203
- 484 Obeng-Ofori D, Reichmuth C, Bekele J, Hassanali A (1997) Biological activity of  
485 1,8-cineole, a major component of essential oil *Ocimum kenyense*  
486 (Ayobangira) against stored product beetles. J Appl Entomol 121:237-243
- 487 O'Donovan JT (1992) Seed yields of canola and volunteer barley as influenced by  
488 their relative times of emergence. Canadian J Plant Sci 72:263-267
- 489 Polatoğlu K, Karakoç OC, Gören N (2013) Phytotoxic, DPPH scavenging,  
490 insecticidal activities and essential oil composition of *Achillea vermicularis*, *A.*  
491 *teretifolia* and proposed chemotypes of *A. biebersteinii* (Asteraceae). Ind  
492 Crops Prod 51:35-45
- 493 Roh HS, Lee BH, Park CG (2013) Acaricidal and repellent effects of myrtacean  
494 essential oils and their major constituents against *Tetranychus urticae*  
495 (Tetranychidae). J Asia-Pac Entomol 16:245-249
- 496 Romagni JG, Allen SN, Dayan FE (2000) Allelopathic effects of volatile cineoles on  
497 two weedy plant species. J Chem Ecol 26:303-313
- 498 Rouimi P, Zucchini-Pascal N, Dupont G, Razpotnik A, Fouche E, De Sousa G,  
499 Rahmani R (2012) Impacts of low doses of pesticide mixtures on liver cell  
500 defence systems. Toxicol In Vitro 26:718-726
- 501 Scarfato P, Avallone E, Iannelli P, De Feo V, Acierno D (2007) Synthesis and  
502 characterization of polyurea microcapsules containing essential oils with  
503 antigerminative activity. J Appl Polym Sci 105:3568-3577
- 504 Seefeldt SS, Jensen JE, Fuerst EP (1995) Log-Logistic Analysis of Herbicide Dose-  
505 Response Relationships. Weed Technol 9:218-227

- 506 Singh HP, Batish DR, Kohli RK (2002) Allelopathic effects of two volatile  
507 monoterpenes against billy goat weed (*Ageratum conyzoides* L.). *Crop Prot*  
508 21:347-350
- 509 Singh HP, Batish DR, Setia N, Kohli RK (2005) Herbicidal activity of volatile oils  
510 from *Eucalyptus citriodora* against *Parthenim hysterophorus*. *Ann Appl Biol*  
511 146:89-94
- 512 Vaughn S, Spencer G (1993) Volatile Monoterpenes as Potential Parent Structures for  
513 New Herbicides. *Weed Sci* 41:114-119
- 514 Vaughn S, Spencer G (1996) Synthesis and Herbicidal Activity of Modified  
515 Monoterpenes Structurally Similar to Cinmethylin. *Weed Sci* 44:7-11
- 516 Wang Y, Chen W, Lin L, Yen J (2010) Dissipation of herbicides chlorosulfuron and  
517 imazosulfuron in the soil and the effects on the soil bacterial community. *J*  
518 *Environ Sci Health Part B* 45:449-455
- 519 Willemsen RE, Hailey A (2001) Effect of spraying the herbicides 2,4-D and 2,4,5-T  
520 on a population of the tortoise *Testudo hermanni* in southern Greece. *Environ*  
521 *Pollut* 113:71-78
- 522 Wilson GC, Soltani N, Tardif FJ, Swanton CJ, Sikkema PH (2010) Control of  
523 volunteer cereals with post-emergence herbicides in maize (*Zea mays* L.).  
524 *Crop Prot* 29:1389-1395
- 525 Yang F, Li X, Zhu F, Lei C (2009) Structural characterization of nanoparticles loaded  
526 with garlic essential oil and their insecticidal activity against *Tribolium*  
527 *castaneum* (Herbst) (Coleoptera: Tenebrionidae). *J Agric Food Chem*  
528 57:10156-10162



529 **Figure captions**

530 **Fig. 1** Structures of the substances used in the post-emergence herbicidal assessments

531

532 **Fig. 2** Effect of (a) acetic acid, (b) 1,8-cineole, (c) benzoic acid, (d) 3-oxo-1,8-  
533 cineole, (e) hexanoic acid and (f) eucalyptus oil on post-emergence growth of roots  
534 (—◆—) and shoots (—■—) of radish 48 hours after exposure. Bars = means ± SE; \*means  
535 at and above this concentration were significantly less than (solvent) control means

536

537 **Fig. 3** Effect of (a) 2-*endo*-hydroxy-1,8-cineole, (b) 2-*exo*-hydroxy-1,4-cineole, (c) 3-  
538 *exo*-hydroxy-1,8-cineole, (d) 2-*exo*-acetoxy-1,4-cineole, (e) 3-*exo*-benzoxy-1,8-  
539 cineole and (f) 2-*exo*-hexoxy-1,4-cineole on post-emergence growth of roots (—◆—)  
540 and shoots (—■—) of radish 48 hours after exposure. Bars = means ± SE; \*means at  
541 and above this concentration were significantly less than (solvent) control means

542

543 **Fig. 4** Effect of (a) acetic acid, (b) 1,8-cineole, (c) *t*-butylacetic acid, (d) 3-oxo-1,8-  
544 cineole, (e) hexanoic acid and (f) eucalyptus oil on post-emergence growth of roots  
545 (—◆—) and shoots (—■—) of ryegrass 48 hours after exposure. Bars = means ± SE;  
546 \*means at and above this concentration were significantly less than (solvent) control


547 means

548


549 **Fig. 5** Effect of (a) 2-*endo*-hydroxy-1,8-cineole, (b) 3-*exo*-hydroxy-1,8-cineole, (c) 3-  
550 *exo*-hexoxy-1,8-cineole, (d) 2-*exo*-hydroxy-1,4-cineole, (e) 3-*exo-t*-butylacetoxy-1,8-  
551 cineole and (f) 2-*exo*-acetoxy-1,4-cineole on post-emergence growth of roots (—◆—)  
552 and shoots (—■—) of ryegrass 48 hours after exposure. Bars = means ± SE; \*means at

553 and above this concentration were significantly less than (solvent) control means;


554 means significantly higher than (solvent) control means




**Figure 1**




**Figure 2**



**Figure 3**



**Figure 4**



**Figure 5**

**Table 1 Solvents used for bioassays of test compounds**

<b>Compound</b>	<b>Solvent</b>
acetic acid	water
benzoic acid	trichloromethane (chloroform)
hexanoic acid	hexane
<i>t</i> -butylacetic acid	hexane: chloroform; 99:1
1,8-cineole <b>1</b>	Tween <sup>®</sup> 80 in water (0.34 g L <sup>-1</sup> )
eucalyptus oil	Tween <sup>®</sup> 80 in water (0.34 g L <sup>-1</sup> )
3-oxo-1,8-cineole <b>2</b>	hexane
2- <i>endo</i> -hydroxy-1,8-cineole <b>3</b>	hexane: chloroform; 99:1
3- <i>exo</i> -hydroxy-1,8-cineole <b>4a</b>	hexane
3- <i>exo</i> -benzoxy-1,8-cineole <b>4b</b>	chloroform
3- <i>exo</i> -hexoxy-1,8-cineole <b>4c</b>	hexane
3- <i>exo-t</i> -butylacetoxy-1,8-cineole <b>4d</b>	chloroform
2- <i>exo</i> -hydroxy-1,4-cineole <b>5a</b>	hexane: chloroform; 9:1
2- <i>exo</i> -acetoxy-1,4-cineole <b>5b</b>	hexane
2- <i>exo</i> -hexoxy-1,4-cineole <b>5c</b>	hexane



**Table 2 Concentration of test compounds used in post-emergent herbicidal testing**

Compound	Concentrations (mol L <sup>-1</sup> )	
	Radish	Rye Grass
acetic acid	0.0025, 0.01, 0.025, 0.05	
benzoic acid	0.000316, 0.001, 0.00316, 0.01, 0.0316	
hexanoic acid	0.001, 0.0025, 0.007, 0.025, 0.05	0.001, 0.0025, 0.007, 0.02, 0.05
<i>t</i> -butylacetic acid		0.000316, 0.001, 0.00316, 0.01, 0.0316
1,8-cineole <b>1</b>	0.001, 0.00316, 0.01, 0.0316, 0.1	0.001, 0.01, 0.0316, 0.1, 0.316
eucalyptus oil	<sup>a</sup> 0.001, 0.01, 0.0316, 0.1, 0.316	<sup>a</sup> 0.001, 0.00316, 0.01, 0.0316, 0.1
3-oxo-1,8-cineole <b>2</b>	0.005, 0.01, 0.025, 0.05, 0.075	
2- <i>endo</i> -hydroxy-1,8-cineole <b>3</b>	0.005, 0.01, 0.05, 0.1, 0.2	
3- <i>exo</i> -hydroxy-1,8-cineole <b>4a</b>	0.01, 0.025, 0.05, 0.1, 0.2	
3- <i>exo</i> -benzoxy-1,8-cineole <b>4b</b>	0.01, 0.0316, 0.1, 0.316, 1.0	
3- <i>exo</i> -hexoxy-1,8-cineole <b>4c</b>		0.001, 0.00316, 0.01, 0.0316, 0.1
3- <i>exo-t</i> -butylacetoxy-1,8-cineole <b>4d</b>		0.001, 0.01, 0.0316, 0.1, 0.316
2- <i>exo</i> -hydroxy-1,4-cineole <b>5a</b>	0.01, 0.02, 0.04, 0.08, 0.1	
2- <i>exo</i> -acetoxy-1,4-cineole <b>5b</b>	0.001, 0.00316, 0.01, 0.0316, 0.1	
2- <i>exo</i> -hexoxy-1,4-cineole <b>5c</b>	0.01, 0.0316, 0.1, 0.316, 1	

<sup>a</sup> Concentration units for eucalyptus oil solution is g mL<sup>-1</sup>