



Local restoration of dystrophin expression with the morpholino oligomer AVI-4658 in Duchenne muscular dystrophy: a single-blind, placebo-controlled, dose-escalation, proof-of-concept study

Maria Kinali*, Virginia Arechavala-Gomez*, Lucy Feng, Sebahattin Cirak, David Hunt, Carl Adkin, Michela Guglieri, Emma Ashton, Stephen Abbs, Petros Nihoyannopoulos, Maria Elena Garralda, Mary Rutherford, Caroline McCulley, Linda Popplewell, Ian R Graham, George Dickson, Matthew JA Wood, Dominic J Wells, Steve D Wilton, Ryszard Kole, Volker Straub, Kate Bushby, Caroline Sewry, Jennifer E Morgan, Francesco Muntoni

Summary

Background Mutations that disrupt the open reading frame and prevent full translation of *DMD*, the gene that encodes dystrophin, underlie the fatal X-linked disease Duchenne muscular dystrophy. Oligonucleotides targeted to splicing elements (splice switching oligonucleotides) in *DMD* pre-mRNA can lead to exon skipping, restoration of the open reading frame, and the production of functional dystrophin in vitro and in vivo, which could benefit patients with this disorder.

Methods We did a single-blind, placebo-controlled, dose-escalation study in patients with DMD recruited nationally, to assess the safety and biochemical efficacy of an intramuscular morpholino splice-switching oligonucleotide (AVI-4658) that skips exon 51 in dystrophin mRNA. Seven patients with Duchenne muscular dystrophy with deletions in the open reading frame of *DMD* that are responsive to exon 51 skipping were selected on the basis of the preservation of their extensor digitorum brevis (EDB) muscle seen on MRI and the response of cultured fibroblasts from a skin biopsy to AVI-4658. AVI-4658 was injected into the EDB muscle; the contralateral muscle received saline. Muscles were biopsied between 3 and 4 weeks after injection. The primary endpoint was the safety of AVI-4658 and the secondary endpoint was its biochemical efficacy. This trial is registered, number NCT00159250.

Findings Two patients received 0·09 mg AVI-4658 in 900 µL (0·9%) saline and five patients received 0·9 mg AVI-4658 in 900 µL saline. No adverse events related to AVI-4658 administration were reported. Intramuscular injection of the higher-dose of AVI-4658 resulted in increased dystrophin expression in all treated EDB muscles, although the results of the immunostaining of EDB-treated muscle for dystrophin were not uniform. In the areas of the immunostained sections that were adjacent to the needle track through which AVI-4658 was given, 44–79% of myofibres had increased expression of dystrophin. In randomly chosen sections of treated EDB muscles, the mean intensity of dystrophin staining ranged from 22% to 32% of the mean intensity of dystrophin in healthy control muscles (mean 26·4%), and the mean intensity was 17% (range 11–21%) greater than the intensity in the contralateral saline-treated muscle (one-sample paired *t* test *p*=0·002). In the dystrophin-positive fibres, the intensity of dystrophin staining was up to 42% of that in healthy muscle. We showed expression of dystrophin at the expected molecular weight in the AVI-4658-treated muscle by immunoblot.

Interpretation Intramuscular AVI-4658 was safe and induced the expression of dystrophin locally within treated muscles. This proof-of-concept study has led to an ongoing systemic clinical trial of AVI-4658 in patients with DMD.

Funding UK Department of Health.

Introduction

Duchenne muscular dystrophy (DMD) affects 1 in 3500 newborn boys, causing eventually progressive muscle weakness, cardiomyopathy, and respiratory failure. Patients are diagnosed when they are toddlers, become wheelchair-dependent in their early teens, and die in their 20s. With improvements in standards of care, including non-invasive ventilation and glucocorticoid and cardioprotective treatment, many individuals with DMD survive beyond their mid-20s^{1,2} despite having severe and disabling weaknesses.

DMD is caused by the absence of the protein dystrophin. Dystrophin associates with other sarcolemmal proteins of the dystrophin glycoprotein complex and connects the cytoskeleton to the extracellular matrix. The absence of dystrophin reduces the stability of the sarcolemma and increases intracellular calcium influx, which is followed by degeneration of the muscle fibres. Dystrophin is encoded by *DMD*. Deletions (in about 65% of patients), duplications (in about 10% of patients), point mutations (in about 10% of patients), or other smaller rearrangements can disrupt the open reading frame of *DMD*, leading to

Lancet Neurol 2009; 8: 918–28

Published Online

August 26, 2009

DOI:10.1016/S1474-

4422(09)70211-X

See [Reflection and Reaction](#)

page 873

*Contributed equally

The Dubowitz Neuromuscular Centre, University College London Institute of Child Health London, UK

(M Kinali MD,

V Arechavala-Gomez PhD,

L Feng PhD, S Cirak MD,

Carl Adkin PhD, C Sewry PhD,

J E Morgan PhD, F Muntoni MD);

Department of Pediatrics,

Hammersmith Hospital,

London, UK (M Kinali,

F Muntoni); Department of

Orthopaedics, St Mary's

Hospital, London, UK

(D Hunt FRCS); Institute of

Human Genetics, Newcastle

University, Newcastle, UK

(M Guglieri MD, V Straub MD,

K Bushby MD); DNA Laboratory

Genetics Centre, Guy's

Hospital, London, UK

(E Ashton PhD, S Abbs PhD);

Department of Cardiology

(P Nihoyannopoulos MD),

Division of Neuroscience and

Mental Health (C McCulley PhD,

D J Wells PhD), and Robert

Steiner MRI Unit

(M Rutherford MD),

Hammersmith Hospital

Campus, Imperial College

London, UK; Academic Unit of

Child and Adolescent

Psychiatry, Imperial College

St Mary's Campus, London, UK

(M E Garralda MD);

Hammersmith Hospital

Campus, Imperial College

London, UK; Royal Holloway

University of London, UK

(L Popplewell PhD,

I R Graham PhD, G Dickson PhD);

Department of Physiology,

premature termination of its translation,^{3,4} whereas deletions or duplications that maintain the open reading frame can lead to truncated but functional dystrophin, which underlies the milder disorder Becker muscular dystrophy (BMD).⁵ The spectrum of severity for BMD varies, ranging from difficulties in walking in the late teens to preserved walking ability into late adulthood and a normal lifespan.⁶

Up to 50% of patients with DMD have sporadic dystrophin-positive revertant fibres.⁷ This dystrophin expression arises from alternative processing of *DMD* pre-mRNA that skips some exons, leading to restoration of the open reading frame.⁸ Revertant dystrophin is correctly localised to the sarcolemma and mediates the assembly of other proteins of the dystrophin glycoprotein complex, suggesting that it is physiologically functional. The occurrence of revertant fibres and the mild symptoms of some individuals with BMD with in-frame deletions suggest that it might be feasible to modify the splicing of the *DMD* transcript and, by skipping the mutated exons, produce functional dystrophin (figure 1).

Some exon deletions are more common than others.¹¹ Deletions of exons 50, 52, 52–63, 45–50, 47–50, and 49–50 cumulatively account for 13% of all the deletions in *DMD*.¹² Skipping of exon 51 in patients with these deletions should restore the open reading frame of *DMD* and lead to the expression of functional dystrophin.

Antisense oligonucleotides have been used for experimental gene silencing and recently as splice-switching oligonucleotides to modify splicing and induce exon skipping,¹³ particularly in myoblasts from patients with DMD *in vitro*,^{14,15} and in mouse and dog models of DMD.^{16–18} One patient with DMD who had a deletion of exon 20 received an intravenous infusion of a splice-switching oligonucleotide with a phosphorothioate backbone, which induced skipping of exon 19 and restored the *DMD* open reading frame in lymphocytes but had no effect in skeletal muscle.¹⁹ A recent phase 1 clinical study reported encouraging results in four boys with DMD who received a single intramuscular injection of a 2'-O-methyl-ribooligonucleoside-phosphorothioate splice-switching oligonucleotide that was targeted to skip exon 51. This treatment led to appreciable expression of dystrophin and was well tolerated.²⁰

Other chemically modified oligonucleotides have been used in preclinical models and clinical trials. Phosphorodiamidate morpholino oligomers (PMOs; figure 1) are non-toxic, and in the *mdx* mouse model of DMD they were the most effective oligomer chemistry for inducing exon skipping and restoring long-lasting (weeks) dystrophin expression after intravenous or intramuscular injection.^{21–24} PMOs, unlike other antisense oligonucleotides, are uncharged, not metabolised, and in preclinical or clinical studies were not associated with activation of the immune system, anaphylaxis, hypotension, or anti-arrhythmias.²⁵ On the basis of these data, we have

studied the safety and biochemical efficacy of AVI-4658, a PMO designed to target exon 51 that is delivered by intramuscular injection. Here, we report the results of a single-blind, placebo-controlled, dose-escalation safety and efficacy study of PMOs in patients with DMD.

Methods

Patients

This single-site, non-randomised, single-blind (investigator) study was done at Imperial College NHS Trust, London, UK, in patients with DMD who were recruited nationally. Participants were boys with a classic clinical diagnosis of DMD²⁶ who were aged between 10 and 17 years inclusive when the study drug was given. All participants had a deletion that can be rescued by the skipping of exon 51 (eg, deletion of exons 45–50, 47–50, 48–50, 49–50, 50; 52, or 52–63); had fewer than 5% revertant fibres seen in a muscle biopsy; had the extensor digitorum brevis (EDB) muscle sufficiently preserved (grade 1 to 3: grade 1 is near normal; grade 2 is 30–60% of the muscle is normal; and grade 3 is muscle is almost all abnormal but some normal muscle still present at the periphery), as determined by MRI of the feet;^{27,28} had a forced vital capacity of 25% or more and a normal overnight sleep study before 3 months from the day of injection; were able to comply with all study assessments and return for all study visits; and had adequate

Anatomy and Genetics, University of Oxford, UK (M J A Wood MD); Centre for Neuromuscular and Neurological Disorders, University of Western Australia, Perth, WA, Australia (SD Wilton PhD); AVI Biopharma, Corvallis, Oregon, USA (R Kole PhD); Centre for Inherited Neuromuscular Disorders, Robert Jones and Agnes Hunt NHS Trust, Oswestry, UK (C Sewry)

Correspondence to: Francesco Muntoni, Dubowitz Neuromuscular Centre, University College London Institute of Child Health, 30 Guilford Street, London, WC1N 1EH, UK
f.muntoni@ich.ucl.ac.uk




Figure 1: Deletions and predicted results of exon skipping in the patients who were studied

(A) Pre-mRNA transcripts and dystrophin protein products from full length *DMD*, in patients with Duchenne muscular dystrophy, and predicted protein sequences after exon skipping. (I) The normal dystrophin gene produces the full length dystrophin product. (II) Patients 1 and 2 had a deletion in exon 50 that disrupts the open reading frame, leading to a truncated and unstable dystrophin. (III) Skipping of exon 51 restores the reading frame, producing a truncated but functional dystrophin that lacks exons 50 and 51. (IV) Patient 7 is missing exons 49 and 50. (V) Patients 3 and 4 are missing exons 48–50. (VI) Patients 5 and 6 are missing exons 45–50. All the truncated dystrophins produced after skipping of exon 51 are missing the hinge 3 region and some of the rod domain but have been associated with the milder BMD phenotype.^{3,10} (B) Structure of the phosphorodiamidate morpholino modification of the antisense oligomer.

Age at enrolment (years)	DMD deletion	Mobility	Steroids	Age at first biopsy (years)	Dystrophin-positive fibres in original biopsy	MRI grading of EDB muscle	EDB fibrosis		Time between injection and EDB biopsy (weeks)	PCR primers to exons	Amplicon sizes (bp)	
							Saline injected	Treated				
Low dose												
1	16	14 bp deletion in intron 49 that included the exon 50 acceptor splice site	Wheelchair for 11 years	N	8	A few revertant fibres, (~1–2%); traces on a few fibres	2a	++	++	3	48 and 52	519–286
2	13	Exon 50	Wheelchair for 10 years; rides static bike for 10 min daily	N	7	No revertant fibres; no traces	∞ 2b/3	+++	+++	4	48 and 52	519–286
High dose												
3	11	Exons 48–50	Wheelchair for 10 years	Y	7	No revertant fibres; no traces	∞ 2b/3	+++	++	4	46 and 52	570–337
4	15	Exons 48–50	Walks indoors	Y	3	One revertant fibre; traces on many fibres	∞ 2a/2b	+	+	4	46 and 52	570–337
5	11	Exons 45–50	Walks unaided	Y	7	No revertant fibres; traces on a few fibres	1	++	++	4	43 and 52	486–253
6	12	Exons 45–50	Walks unaided	Y	3	No revertant fibres; traces on many fibres	2a	++	++	4	43 and 52	486–253
7	10	Exons 49–50	Walks unaided	Y	4	No revertant fibres; traces on many fibres	1	+	+	3	47 and 52	539–306

Numbers are patient number. EDB=extensor digitorum brevis. bp=base pair. Y=yes. N=no. ∞=Asymmetrical EDB involvement on muscle MRI grading. +=Moderate increase of perimysial and endomysial connective tissue; some areas had a severe increase in perimysial and endomysial connective tissue. ++=Most fibres were surrounded by large amounts of connective tissue, but some areas had less and were compact. +++=All fibres surrounded by connective tissue; severe fibrosis throughout sample. EDB=extensor digitorum brevis. bp=base pairs. PCR=polymerase chain reaction.

Table 1: Baseline characteristics, exons targeted by PCR primers, and predicted amplicon sizes

psychiatric adjustments, supportive psychosocial circumstances, and full understanding of the study aims, process, and likely outcomes.

Exclusion criteria were: absence of EDB muscles or advanced pathology of EDB muscles (grade 4) on muscle MRI; left ventricular shortening fraction of 25% or less, an ejection fraction of less than 35% seen by echocardiography within 3 months of visit one, or both; respiratory insufficiency defined by the need for invasive or non-invasive ventilation; severe cognitive dysfunction that meant the patient was unable to understand and collaborate with the study protocol; immune deficiency or autoimmune disease; bleeding disorders or chronic anticoagulant treatment within 3 months before study entry; medication with anabolic steroids, creatine protein supplementation, albuterol, or other beta agonists, and intranasal, inhaled, or topical steroids for a disorder other than muscular dystrophy within 1 week before study entry; surgery within 3 months before study entry or planned for anytime during the study; inability to undergo MRI (eg, owing to metal implants); known allergies to products likely to be used in the study (eg, antiseptics or anaesthetics); and participation in another experimental study within 4 weeks of study entry.

Standard-of-care treatment, including glucocorticoids and cardioprotective drugs, was continued in all patients. All study participants were informed before enrolment of the procedures, risks, and possibility of no benefit. All participants provided written assent, and their parents

gave written informed consent before enrolment in this study. This trial was designed and done in compliance with UK good clinical practice, International Conference on Harmonisation (ICH) E6, and all applicable regulatory requirements were met (UK Medicines and Healthcare Products Regulatory Agency, UK Gene Therapy Advisory Committee, and local research ethics committees).

Trial activities and adverse events were monitored by a safety monitoring committee. The safety monitoring committee met on the following occasions: before recruitment of the first patient; to authorise the recruitment of the second patient after the first patient was biopsied; and after the second patient was studied but before recruitment of the first patient in the high-dose cohort without use of an intermediate dose (0.27 mg). The safety monitoring committee also met to discuss and authorise a proposed change to the protocol, which enabled us to increase the dose directly to the higher dose, to authorise the recruitment of the last two patients in the high-dose cohort, and to discuss a severe adverse event (bilateral surgical wound infection after the muscle biopsies) in one of the patients in the second cohort.

The protocol was also amended in May, 2008, so that we did not need to recruit the third and last patient into the low-dose group (0.09 mg) or recruit the three patients into the intermediate dose group (0.27 mg), and permitted us to recruit patients in the high-dose group (0.9 mg). Because we had identified considerable comorbidity that precluded recruitment of some of the

older patients, we also requested and obtained permission to lower the age at inclusion to 10 years and to be able to recruit ambulant patients.

Procedures

To confirm that each patient had less than 5% dystrophin-positive revertant fibres, each of the original muscle biopsies used to diagnose the patients was re-evaluated (table 1).²⁹ Presence of a deletion suitable for exon 51 skipping and no additional mutations were reconfirmed by sequencing of all the intact *DMD* coding exons and their intron–exon boundaries in all patients. The extensor digitorum brevis (EDB) muscle at the back of the foot was selected as the target muscle. This muscle is well preserved in non-ambulant boys with DMD (Kinali and Muntoni, unpublished), is, for the most part, functionally redundant (an important consideration in a study that is not expected to lead to functional benefit), and can even be absent in some individuals.³⁰ MRI confirmed the presence and preservation of the EDB muscles in all patients^{31,32} and that involvement of the EDB muscle was not more than grade 3.^{27,28} Healthy muscle biopsies were obtained from the Dubowitz Neuromuscular Centre biobank.

Psychiatric assessments were done to ascertain the expectations and risk of reactive depression for each patient and their family by documenting previous and current psychiatric adjustment and current psychosocial stresses and supports. Parents and children were interviewed separately. The parental questionnaire included the Strengths and Difficulties Questionnaire, the Parental Stress and Support Questionnaire, the General Health Questionnaire, and the Family Assessment Device, which are all validated assessment tools.^{33–36} Patient interviews with the psychiatrist focused on their understanding of the trial, their general adjustment at home and school, emotional and depressive symptoms, and the Strengths and Difficulties Questionnaire and the Hamilton Anxiety and Depressive Scale.^{34,37}

Cultured fibroblasts from a skin biopsy were analysed to verify oligonucleotide-induced splice switching of exon 51 in all patients. Fibroblasts were forced into myogenic differentiation by transduction with an adenovirus expressing the myogenic regulatory factor protein MyoD,³⁸ and cultures were transfected with AVI-4658 congener on a 2′O-methyl backbone (300 nM) with Lipofectin (Invitrogen, UK).³⁹ RNA was isolated 48 h after transfection and analysed after reverse transcriptase–polymerase chain reaction (RT-PCR) amplification.³⁹ 7 days after transfection, cells were harvested for western blot analysis,³⁹ and lysates were probed with Dys1 (Vector Laboratories, UK), an anti-dystrophin monoclonal antibody. Dysferlin (Vector Laboratories, UK) was used as a loading control. Baseline safety blood analyses, including tests for anti-dystrophin antibodies⁴⁰ and T-cell subsets (CD4: CD8), were also done.

AVI-4658 is an exon 51-targeted PMO (sequence CTCC AACATCAAGGAAGATGGCATTCTAG).³⁹ AVI-4658 was synthesised and purified by AVI BioPharma (Portland, OR, USA) and was supplied as a low endotoxin and low bioburden powder, which was reconstituted in normal saline in the operating theatre.

This dose escalation intramuscular trial was done in seven patients, who received either of two doses: two patients received 0·09 mg and five patients received 0·9 mg of AVI-4658; both doses were diluted in 900 µL normal saline (0·9%) and were injected in one EDB muscle; the contralateral EDB muscle was injected with 900 µL normal saline. The dose was divided into nine 100 µL injections in the first five patients and four 225 µL injections in the last two patients; this regimen reduced leakage into the skin, which was seen to some extent in patients 1–4. To ensure delivery in the muscle, the drug was injected with a 22 gauge EMG delivery needle (Pajunk, Multistim Sensor, Germany) inside a 1 cm² grid drawn in non-permanent ink on the skin over the site of the EDB muscles. The site and depth of the injections were recorded on videotape. After each injection, the needle was manipulated to confirm its correct placement within the muscle. This took about 1 min. Each infusion was completed in about 30 s. After infusion, the needle was left in place for about 30 s to avoid leakage. The choice of which muscle to inject with the PMO or saline was made in the operating theatre on the day of the injection and the person who made the decision (FM) was masked to whether the patient was right-handed or left-handed. Patients and investigators (except MK, SC, and FM) were masked to which site received the active compound. The procedure was done under general anaesthetic in six patients; one patient opted to have the treatment under local anaesthetic. Both types of anaesthesia were available, and the choice was left to the families.

For all patients, an open biopsy of both EDB muscles was done between 3 and 4 weeks after injection. The rationale for this time frame was taken from previous work in mice and in humans.^{17,20} In the *mdx* mouse, the same level of dystrophin expression was detected at 4 weeks as was detected at 2 weeks after an intramuscular injection of a 2′O-methyl antisense oligonucleotide designed to skip dystrophin exon 23.¹⁷ Also, in a study of intramuscular injection of a 2′O-methyl antisense oligonucleotide that targeted exon 51, given to patients with DMD, the skipped products and dystrophin were still detected in muscles that were analysed 4 weeks after intramuscular injection.²⁰ The area immediately below the needle track was exposed and an open biopsy was taken and rapidly frozen in liquid nitrogen-cooled isopentane, according to standard techniques.²⁹ To ensure that all the injection site had been obtained, most of the EDB muscle was removed.

Safety was determined by physical examination and haematological and urinary parameters, which were assessed periodically. The injection sites were monitored for local reaction and reactive pain. Patients were

followed-up at timed intervals for 120 days after treatment. Any immune response against the newly synthesised dystrophin was assessed by the production of anti-dystrophin antibodies: serum samples were used to probe western blots loaded with lysates of muscle from the AD17 transgenic mouse, which overexpresses full-length human dystrophin; goat anti-human-IgG was used as the secondary antibody (Bio-Rad, UK).⁴⁰ The presence of T cells and B cells in each biopsy was ascertained by immunohistochemistry with antibodies raised against human CD3, CD4, CD8, or CD20 (Dako, UK).

RNA extraction and RT-PCR analysis were done on ten serial 7 μ m sections of the frozen muscle sample.³⁹ Direct DNA sequencing of the excised bands was done by University College London Scientific Support Services. For western blotting, proteins from 20 serial 10 μ m sections of muscle were isolated directly in 50 μ L of

loading buffer and analysed as previously described.³⁹

For immunohistochemical detection,²⁹ unfixed, frozen serial sections (7 μ m) were incubated for 1 h with monoclonal antibodies against dystrophin (Dys 2 [exon 77–79]; Vector Laboratories, UK), MANDYS106⁴¹ (exon 43; a gift from G Morris, Oswestry, and the MDA Monoclonal Antibody Resource), and β -spectrin (Vector Laboratories, UK)²⁹ and were then assessed by two investigators (LF and CS) who were masked to the identity of the patient and which side received the active compound. Images were captured with a Leica DMR microscope linked to MetaMorph, version 7.5 (Molecular Devices, CA, USA). Quantitative studies were done as follows: the numbers of dystrophin-positive and dystrophin-negative fibres in the muscle fascicles adjacent to a presumed injection site were counted on the MANDYS106-stained sections of AVI-4658-treated muscles and areas of control muscles chosen at random by two independent investigators (JM and CA). Dystrophin expression was evaluated in 40 muscle fibres selected at random on one representative transverse MANDYS 106-stained region per biopsy. Expression was normalised against the expression of β -spectrin on serial sections and was compared with sections of normal control muscle and the contralateral EDB saline-injected biopsy that were processed in the same way and simultaneously labelled with the same antibodies. Four fields of the immunostained transverse cryosection of each muscle were selected at random (out of focus) and these areas (in focus) were photographed. Ten regions per image, each including an area of membrane and fibre cytoplasm, were selected by moving the cursor across the image and were analysed with MetaMorph. We measured the relative intensity of dystrophin in 100 dystrophin-positive and 100 dystrophin-negative fibres in the same regions of a section of treated muscle from each patient. This trial is registered, number NCT00159250.

Role of the funding source

The study was funded by the UK Department of Health and sponsored by Imperial College London. Neither had a role in the study design, data collection, data analysis, data interpretation, or writing of the report. AVI Biopharma manufactured and supplied AVI-4658 for the study and provided preclinical testing, packaging, labelling and the investigator brochure for the drug. The company supported the toxicity studies and participated in the design of the protocol, the execution and monitoring of the study, and discussions with the regulatory authorities. All authors have seen and approved the submitted version of the manuscript.

Results

MRI confirmed that the EDB muscles in all patients had changes that were less than grade 4 (figure 2). The diagnostic muscle biopsies were re-analysed with Dys1, Dys2, and Dys 3 antibodies to confirm there was no or little dystrophin and less than 5% of the fibres were




Figure 2: Procedure for prescreening of patients before injection of AVI-4658.

Patient 3 is shown as an example; similar results were obtained for all patients. (A) Transverse MRI of the lower leg and coronal MRI of the extensor digitorum brevis muscle (arrow) confirmed the suitability of the muscle. (B) Skin fibroblasts from all patients were forced into myogenic differentiation and treated with an AVI-4658 congener to confirm exon skipping and dystrophin production. RT-PCR analysis shows two bands: the high molecular weight band corresponds to the unskipped transcript (including exons 46, 47, 51, and 52) and the low molecular weight band corresponds to the transcript fragment with size specific skipping of exon 51. (C) Exon 51 skipping was confirmed by sequencing.

revertant (table 1). Treatment of MyoD transfected fibroblasts with the 2XO-methyl congener of AVI-4658 showed exon skipping in the RT-PCR products (confirmed by sequencing) and dystrophin expression on western blot in all patients (figure 2, webappendix).

Four patients who had mild cardiac involvement were on treatment with angiotensin-converting enzyme (ACE) inhibitors before recruitment (table 2). Patient psychiatric adjustment and family psychosocial circumstances were deemed to be adequate for all patients included in the study. One patient who was from a family with unrealistic expectations and psychiatric problems was excluded.

All safety assessments showed no adverse events that were related to AVI-4568. All patients showed some short-lived, bilateral, localised reactions in both EDB muscles after the injections and muscle biopsies (table 2). In one patient, the echocardiogram at 3 months after the screening visit showed deterioration in fractional shortening, despite ACE inhibitors. This was attributed to the natural history of DMD, and the patient's condition stabilised on beta-blockers. One patient developed bilateral cellulitis in the feet after the muscle biopsies and needed intravenous antibiotics. A short period of refusal to bear weight resolved without consequences.

Light microscopy and immunocytochemistry done masked to treatment showed no differences in inflammatory infiltrates between the treated and control

EDB muscles in all patients (table 2). There was no induction of anti-dystrophin antibodies after treatment with the phosphorodiamidate morpholino oligomer, although two patients (patient one and patient six) had low levels of cross reactivity with dystrophin in their pre-treatment and post-treatment serum samples (not shown).

Both patients who had the low dose AVI-4658 showed little expression of dystrophin, despite dystrophin being robustly restored in the cultured fibroblasts, ahead of the injection of the antisense AVI-4658. This suggests that there is a lower threshold effect, and the low dose did not seem to be sufficient to induce exon skipping in patients with Duchenne muscular dystrophy; therefore, we proceeded straight to the high dose.

Biopsies of AVI-4658-injected EDB muscles and the contralateral saline-injected EDB muscles were analysed by assessors who were masked to which muscle was the treated one. This involved quantification of immunostained, dystrophin-positive fibres, the detection of exon 51 skipped RNA (table 1), and immunoblot analysis. After immunostaining of muscle sections with anti-dystrophin antibodies (Dys2 and MANDYS106), all patients treated with high-dose AVI-4658 showed a strong dystrophin signal that prevented masking of which side was the treated one in all patients (figure 3). The results also showed variable low-level immunostaining of the

See Online for webappendix

	Relative intensity of dystrophin		Adverse events	AVI-4658 injection	Inflammatory infiltrate in muscle	Anti-dystrophin antibodies
	Untreated	Treated				
Low dose						
1	9%	6%	Mild bilateral oedema of the forefoot that resolved on day 3	Bilateral mild discomfort, erythema (<25 mm), slight induration, and ecchymosis (>30 mm but ≤50 mm) at the injection sites that resolved on day 3	No difference between the two sides	No difference after injection
2	8%	6%	No local side-effects; decline in cardiac function (FS=22%) but was on ACE inhibitors before the EDB muscle biopsy	Bilateral erythema (50–85 mm) and induration (<25 mm) at the injection sites that resolved on day 3	No difference between the two sides	No difference after injection
High dose						
3	11%	22%	No local side-effects; mild biochemical evidence of myoglobinuria, which was self-limiting and resolved after the third micturition after the muscle biopsies	Bilateral mild discomfort, erythema (<25 mm), and slight induration that resolved on day 2	No difference between the two sides	No difference after injection
4	14%	32%	Bilateral ecchymosis that resolved on day 7	Bilateral ecchymosis that resolved on day 3	No difference between the two sides	No difference after injection
5	10%	31%	Mild biochemical evidence of myoglobinuria that was self-limiting and resolved after the third micturition after the muscle biopsies; cellulitis (local pain and redness) in both feet at the sites of the biopsies was treated with a short course of intravenous and then oral antibiotics; refusal to bear weight for 10 days owing to moderate discomfort	Ecchymosis (<20 mm) and slight induration at the AVI-4658 injection site that resolved on day 2	No difference between the two sides	No difference after injection
6	8%	25%	Large ecchymosis (>50 mm) that resolved on day 7	Mild biochemical evidence of myoglobinuria that was self limiting and resolved after three micturitions after the general anaesthetic; mild ecchymosis (<20 mm) on the control foot that resolved after 2 days	No difference between the two sides	No difference after injection
7	4%	22%	No local side-effects or problems	No local side-effects and no problems	No difference between the two sides	No difference after injection

Numbers are patient number. EDB=extensor digitorum brevis. ACEI=angiotensin converting enzyme. FS=shortening fraction.

Table 2: Safety studies and adverse reactions documented during the trial




Figure 3: Dystrophin expression in patients treated with high-dose AVI-4658
 Transverse sections of treated and contralateral EDB muscles that were immunostained for dystrophin with MANDYS106. (A) Low-power micrograph of a whole section taken with $\times 10$ objective lens shows widespread expression of dystrophin in fibres from the treated muscle in patient 4. (B) Higher magnification ($\times 20$ objective lens) of dystrophin immunolabelling in treated and untreated sections in patients 3-7. Scale bars=100 μm .

	Untreated		Treated	
	Total	Positive	Total	Positive
3	443	21 (5%)	377	182 (49%)
4	662	2 (<1%)	792	623 (79%)
5	475	2 (<1%)	263	116 (44%)
6	554	5 (1%)	404	264 (65%)
7	405	3 (<1%)	262	164 (63%)

Table 3: Dystrophin expression in muscle myofibres in patients 3-7, who were treated with high-dose AVI-4658

saline-treated muscles in all patients, which underscores the importance of this control (webappendix). Sarcolemmal colocalisation of dystrophin with other proteins of the dystrophin glycoprotein complex (webappendix) suggested that dystrophin interacted with other members of this protein complex and was therefore presumed to be functional. However, in patients one and two, who received low-dose AVI-4658, there was no clear difference in protein expression between the treated and control EDB muscle biopsies (webappendix).

The results from the high-dose group were quantified further by image analysis of fluorescent sections stained with the MANDYS106 antibody, which has been used previously for this purpose.²⁰ Masked measurements of the fluorescent intensity of dystrophin staining were done in 40 fibres chosen at random per drug-treated and saline-treated muscle sample. In the high-dose group, the mean difference in measurements of the fluorescent intensity of dystrophin expression in all treated muscles

was about 17% (range 11–21%) more than that in the contralateral saline-injected muscles (one-sample paired *t* test $p=0.002$; figure 4). Because the random measurements took into account areas that contained both dystrophin-positive and dystrophin-negative fibres for the intensity measurements, we targeted the dystrophin-positive fibres within the same area. The intensity in these fibres in patient 4 was 42% of that in healthy muscle (figure 4).

Counting of the positive fibres stained with MANDYS106 was done after confirmation of which muscle was the treated muscle and which was the control, owing to low-level staining in several of the patients (figures 3 and 4). If the low-level immunostaining was not factored in, the number of positive fibres in the side that received AVI-4658 reached 100% in several patients. We therefore adjusted the detection threshold so that only the rare revertant fibres were seen in the saline-injected muscle. This threshold was then subtracted from the contralateral AVI-4658 injected muscle. A mean of 419 fibres (range 262–792 fibres) were seen in the four areas counted from the treated muscles in patients from the high-dose group. In the five muscles treated with high-dose AVI-4658 there was a mean of 269.8 (SD 204.5) dystrophin-positive fibres by contrast with 6.6 (8.1) dystrophin-positive fibres in the saline-treated side (one-sample paired *t* test $p=0.02$). When the proportion of positive fibres in the fascicles that were assumed to relate to the needle track were counted, the number varied between 44% and 79% (mean 59.8% [SD 13.9]; table 3). At least 262 fibres were counted

in each sample. There was no correlation between the proportion of dystrophin-positive fibres and the severity of fibrosis (table 1).

The results of the dystrophin immunostaining were corroborated by those from the RT-PCR and western blots, which were done masked to treatment. Exon 51 skipping and distinct bands of dystrophin protein were seen in the drug-treated muscles but not in the saline-treated muscles of patients in the high-dose group (figure 5). Exon 51 skipping was also seen in the two patients in the low-dose group, but this was less abundant and only detected when high-sensitivity conditions were used (additional cycles);²⁰ however, these two patients did not have detectable dystrophin on western blot (webappendix). Sequencing of the RT-PCR products confirmed accurate skipping of exon 51 in the treated muscles of all patients (figure 5). Immunoblot analysis showed bands of the expected molecular weight in the AVI-4658-treated muscles.

Discussion

This single-blind study assessed the local safety and biochemical efficacy of intramuscular injection of AVI-4658 in patients with DMD who have deletions that are responsive to skipping of *DMD* exon 51. We showed, *in vivo*, that the PMO AVI-4658 induced specific skipping of exon 51 and the production of dystrophin that was correctly localised at the sarcolemma. The treatment was not associated with any systemic or local adverse events or with any immune response against dystrophin.

Specific dose-dependent exon skipping was seen in the treated EDB muscles compared with the contralateral saline-injected muscles. Patients in the low-dose group showed RT-PCR evidence of exon skipping, confirmed by sequencing, but no dystrophin protein was detected.

Strong expression of dystrophin was seen in the treated muscle in all patients in the high-dose cohort after subtraction of the signal intensity for the low-level of antibody labelling in the sarcolemma of the saline-injected contralateral muscle (figure 3). Localisation of dystrophin to the sarcolemma suggests appropriate interaction with other proteins of the dystrophin glycoprotein complex. Western blot analysis detected increased expression of dystrophin in the AVI-4658-treated muscle of all patients who received the high dose, and the immunoblot detected expression of dystrophin of the expected molecular weight in all patients. However, the quantification of dystrophin from a western blot after an intramuscular injection is difficult (eg, ensuring that equivalent amounts of sarcolemmal proteins are loaded in each track, the transfer of large proteins is efficient, and development of the signal in the linear range). In view of the highly localised delivery of AVI-4658, we chose an immunohistochemistry-based method to quantify dystrophin in specific myofibres. This method has several advantages over other techniques, such as western blot and real-time RT-PCR, because it enables the *in situ* visualisation of the correct localisation




Figure 4: Intensity of dystrophin expression in patients treated with high-dose AVI-4658 relative to control

(A) Mean random intensity measurements. (B) Measurement of mean dystrophin intensity in positive fibres: intensity measurements exclusively targeted to 100 dystrophin-positive and 100 dystrophin-negative fibres within the same area in patient 4. Bars are SEM.

of the expressed protein. Additionally, western blot and RT-PCR are not relevant for proteins that are only expressed in a subset of fibres, which is the case here because of the local injection.

The quantitative immunohistochemistry method we used goes a step further than the one used in a recent exon-skipping study,²⁰ because fibres are randomly selected to calculate the mean expression in the treated muscle and compared with that in the control muscle (figure 4). We extended our analysis by measuring the intensity of selected dystrophin-positive and dystrophin-negative fibres within the same areas of the treated muscles. We show that the fluorescent intensity of the dystrophin-positive fibres was 42% of that of fibres in healthy, non-dystrophic control muscles (figure 4).

In a recent clinical trial in which exon 51 was targeted, the investigators showed dystrophin expression in the tibialis anterior muscle after one injection of PRO051, a 2'-O-methyl antisense oligonucleotide.²⁰ Although PRO051 and AVI-4658 target the same region of exon 51, we used a phosphorodiamidate morpholino oligomer chemistry, used a longer splice-skipping oligonucleotide, and treated a small intrinsic and relatively non-functional foot muscle, which enabled us to obtain bilateral muscle biopsies that were not available in the PRO051 study. Because staining with the MANDYS106 antibody, which was used in both studies, results in low-level labelling in some patients with DMD (figure 4), negative controls are




Figure 5: Exon 51 skipping in amplified RNA from treated muscles

(A) RT-PCR analysis of RNA extracted from treated (X), untreated (O), and control (C) muscle sections detects shorter transcript fragments in the treated muscles, with sizes that correspond to the specific skipping of exon 51. (B) Exon 51 skipping was confirmed by sequencing. (C) Western blot analysis of homogenates of treated and untreated muscle (20×10 μm sections) and control muscle (2×10 μm sections [to avoid overexposure]) shows dystrophin expression in extracts from the control muscles (C) and treated (X) extensor digitorum brevis but not in the contralateral muscles (O). Loading was monitored with protogold. Low dose=0.09 mg AV-4658. High dose=0.9 mg AV-4658.

crucial for accurate quantification of the protein. This low-level labelling of dystrophin is absent only in patients with DMD in whom the MANDYS106 epitope (coded in exon 43) is deleted (data not shown), implying that a genuine product of *DMD* is detected by this high-affinity antibody. Our controlled study of exon skipping for DMD clearly shows an increase in the expression of dystrophin in drug-treated muscles compared with the saline-injected contralateral muscle. Because a negative saline-injected control was not included in the previous study,²⁰ and the background concentration of endogenous dystrophin expression was not taken into account, the relative efficacy of the two compounds for inducing dystrophin-positive fibres cannot be compared directly. If we estimate that we obtained an average dystrophin intensity value of 26.4% (range 22.0–32.0%) in a much larger biopsy (about 0.5×2×2 cm; webappendix) compared with the 27% (range 17–35%) in the previous study, which reported biopsy sizes between 120–726 fibres,²⁰ this might suggest that the phosphorodiamidate morpholino oligomer AVI-4658 compares favourably with the 2XO-methyl chemistry.²⁰ This seems to be particularly relevant because the method used to measure intensity values in the PRO051 study averaged the intensities of the whole image and did not subtract the low-level background expression. However the methods for quantifying immunocytochemistry in the two studies are not identical, different muscles were studied, and the volume of the

injected drug was different; therefore, direct comparison between the two studies cannot be made with precision. This implies that, in terms of the different chemistries used for the antisense oligonucleotides, the efficacies of the results of these two studies cannot be directly compared. Although PMOs that were designed to skip exon 23 were more efficient than were 2XO-methyl oligomers after intramuscular injection in *mdx* mice,^{22,24} any differences in length among the antisense oligonucleotides might contribute to their efficacy.²⁴ Nevertheless, both studies reported unequivocal expression of dystrophin at similar concentrations.

Whether this expression of dystrophin resulted in improved muscle function was not studied. However, in-frame deletions of the exons 48–51, 50–51, and 45–51 in *DMD*, which will lead to dystrophin that is similar to that induced in this study, have been described in multi-generation families in which the affected members were asymptomatic.^{9,10} We therefore anticipate that the dystrophin produced by the patients in our study would be functional and speculate that the missing domain is not essential for protein function or structure. The concentration of dystrophin needed to improve or preclude clinical symptoms will probably depend on the quantity and quality (molecular structure) of the protein. We recently reported that dystrophin expression of 27% of the concentration in healthy muscle was sufficient to avoid skeletal muscle symptoms,⁴² but the concentration of truncated dystrophin without exon 51

that is sufficient to provide a clinical benefit is not known. Nevertheless, if the increases in dystrophin concentration that we observed along the needle track were achieved after systemic delivery, then this might lead to a clinically significant response.

Preclinical studies in *mdx* mice have shown that seven weekly doses of PMOs resulted in a high number of dystrophin-positive fibres, varying between 10% and 70% of the muscle fibres in the different muscles analysed.¹⁷ Translation of dose from intramuscular to systemic studies or studies with the same delivery route but in different species is difficult. However, in a recent study¹⁸ in which dogs with canine X-linked muscular dystrophy were given an intravenous cocktail of morpholinos designed to skip exons 6 and 8 of dystrophin, expression of the protein was successfully restored and a clinical benefit was noticed without adverse reactions to the high-doses of morpholinos used. This bodes well for systemic studies in humans. On the basis of these observations, we have initiated a dose-ranging study in ambulant patients with DMD to assess the safety and efficacy of repeated doses of systemic intravenous AVI-4658 (ClinicalTrials.gov, number NCT00844597).

Contributors

MK participated in the design of the clinical trial protocol, drafted the ethics applications and regulatory submissions, identified and coordinated the selection of patients and clinical and laboratory teams, consented and followed-up the study participants, participated in the acquisition of data and had full access to all data and records keeping, including trial codes, managed budgets and finances regarding the MRI and cardiac studies, and supervised the nursing care. VA-G participated in the design of the clinical trial protocol, managed the budget regarding the laboratory consumables, coordinated the laboratory teams, participated in the acquisition, analysis, and interpretation of the data, particularly the RT-PCR, western blotting analysis, image capture, and intensity analysis, and drafted the manuscript. LF participated in the design of the clinical trial protocol, acquisition, analysis, and interpretation of data, particularly in muscle morphology, dystrophin expression, and T-cell studies, and drafted the manuscript. SC selected and screened patients, assisted in the consent process, was in charge during the trial period, followed-up the study participants, coordinated the work between the clinical and laboratory teams, participated in the acquisition, had full access to all data and records keeping, including trial codes, and participated in drafting the manuscript. DH participated in the acquisition of data and drafting of the manuscript. CA participated in the acquisition, analysis, and interpretation of data. RK participated in the drafting of the manuscript. MG participated in the acquisition of data and drafting of the manuscript. EA and SA participated in the acquisition of the genetic data. PN participated in the acquisition of the cardiac data. MEG participated in the psychiatric screening of patients and families. MR participated in the design, analysis, and interpretation of data, and the drafting of the manuscript. CM did the anti-dystrophin antibody analysis and participated in the drafting of the manuscript. LP, IG, GD, VS, KB, JEM, MJW, DJW, and SW participated in the design, analysis, and interpretation of data and the drafting of the manuscript. CS participated in the immunohistochemical assessment of the diagnostic and blinded trial samples, the evaluation of the data, and the drafting of the manuscript. FM participated in the design of the clinical trial protocol, drafted the ethics applications and regulatory submissions, identified patients and coordinated their selection and that of the clinical and laboratory teams, obtained consent from and followed-up the study participants, participated in the acquisition of data and had full access to all data and patient records, including trial codes, managed the budget regarding the entire study, and drafted the manuscript.

Scientific advisory board

UK—Ian Eperon (Leicester); Kay Davies (chair), David Hilton-Jones (Oxford); Christopher Mathew (London [until January, 2009]); Stephen Meech (Duchenne Family Support Group); Marita Polschmidt (Muscular Dystrophy Campaign); Nick Catlin (UK Parent Project). France—Serge Braun (Paris).

Safety monitoring committee

John Harris (chair), Kate Bushby (Newcastle, UK); Andrew George, Duncan Geddes, Francesco Muntoni, Maria Kinali, K Ganeshaguru, Wojtek Rakowicz (London, UK); John Newsom Davis (deceased Aug 24, 2007).

Conflicts of interest

SW is a consultant for AVI Biopharma in exchange for licensing several oligo sequences to them. RK is a full-time employee of AVI. The other authors have no conflicts of interest.

Acknowledgments

The authors wish to thank the participating patients and the charities Muscular Dystrophy Campaign, Action Duchenne, and Duchenne Family Support Group for participating in the UK MDEX consortium (www.mdex.org.uk), which did this study. We particularly want to thank the members of the scientific advisory board for their constructive criticism. The authors wish to thank: Jenny Versnel and Kanagasabai Ganeshaguru for the study coordination in London, and Geoff Bell for patient co-ordination in Newcastle upon Tyne; Laurelle Hughes-Carre for nursing support, D Chambers and J Kim for their help in processing the muscle samples, and A Glover (Radiology Department, Hammersmith Hospital) and KG Hollingsworth (Newcastle University) for support in doing the muscle MRIs. We thank Glenn Morris, Oswestry, and the MDA Monoclonal Antibody Resource for MANDSY106. The contributions of T Partridge and Q Lu (Carolinas Medical Center, Charlotte, NC, USA) are acknowledged for their preclinical work and as the members of the UK MDEX consortium, which initiated the MDEX Consortium grant application and the clinical trial. S Shrewsbury (AVI Biopharma) gave helpful comments on the manuscript. We are grateful for the support of the National Institute of Health Research (NIHR) Biomedical Research Centre Funding Scheme to the Paediatric Clinical Trial Unit at St Mary's Hospital, Imperial College, London, UK, the Muscular Dystrophy Campaign to the Dubowitz Neuromuscular Centre (Centre Grant), and to the North-Star Clinical network, which contributed patients through A Manzur, S Robb, H Jungbluth, J Gosalakal, H Roper, R Quinlivan, and AM Childs. The University College London UK Medical Research Council Neuromuscular Centre Biobank is also acknowledged. MEG is supported by the NIHR Biomedical Research Centre Funding Scheme, Imperial College Healthcare NHS Trust, London, UK. JM is supported by a Wellcome Trust University Award.

References

- 1 Simonds AK, Muntoni F, Heather S, Fielding S. Impact of nasal ventilation on survival in hypercapnic Duchenne muscular dystrophy. *Thorax* 1998; **53**: 949–52.
- 2 Eagle M, Baudouin SV, Chandler C, Giddings DR, Bullock R, Bushby K. Survival in Duchenne muscular dystrophy: improvements in life expectancy since 1967 and the impact of home nocturnal ventilation. *Neuromuscul Disord* 2002; **12**: 926–29.
- 3 Hoffman EP, Brown RH Jr, Kunkel LM. Dystrophin: the protein product of the Duchenne muscular dystrophy locus. *Cell* 1987; **51**: 919–28.
- 4 Morris GE, Ellis JM, Nguyen TM. A quantitative ELISA for dystrophin. *J Immunol Methods* 1993; **161**: 23–28.
- 5 Koenig M, Beggs AH, Moyer M, et al. The molecular basis for Duchenne versus Becker muscular dystrophy: correlation of severity with type of deletion. *Am J Hum Genet* 1989; **45**: 498–506.
- 6 Bushby KM, Gardner-Medwin D, Nicholson LV, et al. The clinical, genetic and dystrophin characteristics of Becker muscular dystrophy. II Correlation of phenotype with genetic and protein abnormalities. *J Neurol* 1993; **240**: 105–12.
- 7 Klein CJ, Covert DD, Bulman DE, Ray PN, Mendell JR, Burghes AH. Somatic reversion/suppression in Duchenne muscular dystrophy (DMD): evidence supporting a frame-restoring mechanism in rare dystrophin-positive fibers. *Am J Hum Genet* 1992; **50**: 950–59.

- 8 Lu QL, Morris GE, Wilton SD, et al. Massive idiosyncratic exon skipping corrects the nonsense mutation in dystrophic mouse muscle and produces functional revertant fibers by clonal expansion. *J Cell Biol* 2000; **148**: 985–96.
- 9 Melis MA, Cau M, Muntoni F, et al. Elevation of serum creatine kinase as the only manifestation of an intragenic deletion of the dystrophin gene in three unrelated families. *Eur J Paediatr Neurol* 1998; **2**: 255–61.
- 10 Lesca G, Testard H, Streichenberger N, et al. Family study allows more optimistic prognosis and genetic counselling in a child with a deletion of exons 50–51 of the dystrophin gene. *Arch Pediatr* 2007; **14**: 262–65.
- 11 Muntoni F, Torelli S, Ferlini A. Dystrophin and mutations: one gene, several proteins, multiple phenotypes. *Lancet Neurol* 2003; **2**: 731–40.
- 12 Aartsma-Rus A, Fokkema I, Verschuuren J, et al. Theoretic applicability of antisense-mediated exon skipping for Duchenne muscular dystrophy mutations. *Hum Mutat* 2009; **30**: 293–99.
- 13 Sazani P, Graziewicz MA, Kole R. Splice switching oligonucleotides as potential therapeutics. In: ST Crooke, ed. *Antisense drug technology: principles, strategies, and applications*. London: Taylor and Francis, 2008: 89–114.
- 14 Aartsma-Rus A, Janson AA, Kaman WE, et al. Therapeutic antisense-induced exon skipping in cultured muscle cells from six different DMD patients. *Hum Mol Genet* 2003; **12**: 907–14.
- 15 Aartsma-Rus A, van Ommen GJ. Antisense-mediated exon skipping: a versatile tool with therapeutic and research applications. *RNA* 2007; **13**: 1609–24.
- 16 McClorey G, Moulton HM, Iversen PL, Fletcher S, Wilton SD. Antisense oligonucleotide-induced exon skipping restores dystrophin expression in vitro in a canine model of DMD. *Gene Ther* 2006; **13**: 1373–81.
- 17 Lu QL, Mann CJ, Lou F, et al. Functional amounts of dystrophin produced by skipping the mutated exon in the *mdx* dystrophic mouse. *Nat Med* 2003; **9**: 1009–14.
- 18 Yokota T, Lu QL, Partridge T, et al. Efficacy of systemic morpholino exon-skipping in duchenne dystrophy dogs. *Ann Neurol* 2009; **65**: 667–76.
- 19 Takeshima Y, Yagi M, Wada H, et al. Intravenous infusion of an antisense oligonucleotide results in exon skipping in muscle dystrophin mRNA of Duchenne muscular dystrophy. *Pediatr Res* 2006; **59**: 690–94.
- 20 van Deutekom JC, Janson AA, Ginjaar IB, et al. Local dystrophin restoration with antisense oligonucleotide PRO051. *N Engl J Med* 2007; **357**: 2677–86.
- 21 Sazani P, Gemignani F, Kang SH, et al. Systemically delivered antisense oligomers upregulate gene expression in mouse tissues. *Nat Biotechnol* 2002; **20**: 1228–33.
- 22 Fletcher S, Honeyman K, Fall AM, Harding PL, Johnsen RD, Wilton SD. Dystrophin expression in the *mdx* mouse after localised and systemic administration of a morpholino antisense oligonucleotide. *J Gene Med* 2006; **8**: 207–16.
- 23 Alter J, Lou F, Rabinowitz A, et al. Systemic delivery of morpholino oligonucleotide restores dystrophin expression bodywide and improves dystrophic pathology. *Nat Med* 2006; **12**: 175–77.
- 24 Heemskerk HA, de Winter CL, de Kimpe SJ, et al. In vivo comparison of 2'-O-methyl phosphorothioate and morpholino antisense oligonucleotides for Duchenne muscular dystrophy exon skipping. *J Gene Med* 2009; **11**: 257–66.
- 25 Muntoni F, Bushby KD, van Ommen G. 149th ENMC International Workshop and 1st TREAT-NMD Workshop on: planning phase I/II clinical trials using systemically delivered antisense oligonucleotides in Duchenne muscular dystrophy. *Neuromuscul Disord* 2008; **18**: 268–75.
- 26 Emery AE, Muntoni F. *Duchenne muscular dystrophy*, 3rd edn. Oxford: Oxford University Press; 2003.
- 27 Mercuri E, C Cini, Counsell S, et al. Muscle MRI findings in a three-generation family affected by Bethlem myopathy. *Eur J Paediatr Neurol* 2002; **6**: 309–14.
- 28 Hawley RJ Jr, Schellinger D, O'Doherty DS. Computed tomographic patterns of muscles in neuromuscular diseases. *Arch Neurol* 1984; **41**: 383–87.
- 29 Dubowitz V, Sewry C. *Muscle biopsy: a practical approach*, 3rd edn. Netherlands: Saunders Elsevier; 2007.
- 30 Macalister A. Additional observation on muscular anomalies in human anatomy (third series), with a catalogue of the principal muscular variations hitherto published. *Trans Roy Irish Acad Sci* 1875; **25**: 1–134.
- 31 Mercuri E, Pichiecchio A, Allsop J, Messina S, Pane M, Muntoni F. Muscle MRI in inherited neuromuscular disorders: past, present, and future. *J Magn Reson Imaging* 2007; **25**: 433–40.
- 32 Mercuri E, Pichiecchio A, Counsell S, et al. A short protocol for muscle MRI in children with muscular dystrophies. *Eur J Paediatr Neurol* 2002; **6**: 305–07.
- 33 Bailey D, Garralda M E. The use of social stress and support interview in families with deviant children: methodological issues. *Social Psychiatry* 1987; **22**: 209–15.
- 34 Goodman R, Ford T, Simmons H, Gatward R, Meltzer H. Using the Strengths and Difficulties questionnaire to screen for child psychiatric disorders. *Br J Psychiatry* 2000; **177**: 534–39.
- 35 Goldberg D. *Manual of the General Health Questionnaire*: NFER Publishing Company, Wiltshire: UK, 1978.
- 36 Miller IW, Epstein NB, Bishop DS, Keitner GI. The McMaster Family Assessment Device: Reliability and Validity. *J Marital Fam Ther* 1985; **11**: 345–56.
- 37 White D, Leach C, Atkinson M, Cottrell D. Validation of the Hospital Anxiety and Depression Scale for use with adolescents. *Br J Psychiatry* 1999; **175**: 452–54.
- 38 Roest PA, van der Tuijn AC, Ginjaar HB, et al. Application of in vitro Myo-differentiation of non-muscle cells to enhance gene expression and facilitate analysis of muscle proteins. *Neuromuscul Disord* 1996; **6**: 195–202.
- 39 Arechavala-Gomez V, Graham IR, Popplewell LJ, et al. Comparative analysis of antisense oligonucleotide sequences for targeted skipping of exon 51 during dystrophin pre-mRNA splicing in human muscle. *Hum Gene Ther* 2007; **18**: 798–810.
- 40 Ferrer A, Foster H, Wells KE, Dickson G, Wells DJ. Long-term expression of full-length human dystrophin in transgenic *mdx* mice expressing internally deleted human dystrophins. *Gene Ther* 2004; **11**: 884–93.
- 41 Nguyen TM, Morris GE. Use of epitope libraries to identify exon-specific monoclonal antibodies for characterization of altered dystrophins in muscular dystrophy. *Am J Hum Genet* 1993; **52**: 1057–66.
- 42 Neri M, Torelli S, Brown S, et al. Dystrophin levels as low as 30% are sufficient to avoid muscular dystrophy in the human. *Neuromuscul Disord* 2007; **17**: 913–18.