



Murdoch
UNIVERSITY

MURDOCH RESEARCH REPOSITORY

This is the author's final version of the work, as accepted for publication following peer review but without the publisher's layout or pagination.

The definitive version is available at

<http://dx.doi.org/10.1098/rspb.2013.3245>

Kopps, A.M., Ackermann, C.Y., Sherwin, W.B., Allen, S.J., Bejder, L. and Krützen, M. (2014) Cultural transmission of tool use combined with habitat specializations leads to fine-scale genetic structure in bottlenose dolphins. *Proceedings of the Royal Society B: Biological Sciences*, 281 (1782).

<http://researchrepository.murdoch.edu.au/21717/>

Copyright: © 2014 The Author(s)

It is posted here for your personal use. No further distribution is permitted.

1 **Cultural transmission of tool use combined with habitat specialisations leads to fine-scale**
2 **genetic structure in bottlenose dolphins**

3

4 Kopps, Anna M.^{a,b} (anna.kopps@gmx.com), Ackermann, Corinne Y.^b, Sherwin, William B.^{a,c},
5 Allen, Simon J.^c, Bejder, Lars^c & Krützen, Michael^{b,a}

6 ^aEvolution & Ecology Research Centre, School of Biological, Earth and Environmental
7 Sciences, University of New South Wales, Sydney, NSW 2052, Australia

8 ^bEvolutionary Genetics Group, Anthropological Institute and Museum, University of Zurich,
9 Winterthurerstrasse 190, 8057 Zurich, Switzerland

10 ^cMurdoch University Cetacean Research Unit, Centre for Fish, Fisheries and Aquatic
11 Ecosystems Research, School of Veterinary and Life Sciences, Murdoch University,
12 South Street, Murdoch, WA 6150, Australia

13

14 **Abstract**

15 Socially learned behaviours leading to genetic population structure have rarely been
16 described outside humans. Here, we provide evidence of fine-scale genetic structure that
17 has likely arisen based on socially transmitted behaviours in bottlenose dolphins (*Tursiops*
18 sp.) in western Shark Bay, Western Australia. We argue that vertical social transmission in
19 different habitats has led to significant geographic genetic structure of mitochondrial DNA
20 (mtDNA) haplotypes. Dolphins with mtDNA haplotypes E or F are found predominantly in
21 deep (> 10m) channel habitat, while dolphins with a third haplotype (H) are found
22 predominantly in shallow habitat (< 10m), indicating a strong haplotype-habitat correlation.
23 Some dolphins in the deep habitat engage in a foraging strategy using tools. These
24 “sponging” dolphins are members of one matriline, carrying haplotype E. This pattern is
25 consistent with what had been demonstrated previously at another research site in Shark
26 Bay, where social vertical transmission of sponging had been shown using multiple lines of
27 evidence. Using an individual-based model, we found support that in western Shark Bay,
28 socially transmitted specialisations may have led to the observed genetic structure. The
29 reported genetic structure appears to present an example of cultural hitchhiking of mtDNA
30 haplotypes on socially transmitted foraging strategies, suggesting that, as in humans, genetic
31 structure can be shaped through cultural transmission.

32

33 Key index words: bottlenose dolphin, cultural hitchhiking, genetic structure, *Tursiops* sp.,
34 social learning

35 **Introduction**

36 Darwinian selection acts on phenotypes, which are manifested through both genetically and
37 non-genetically inherited traits. Both inheritance mechanisms may be adaptive and, thus, of
38 evolutionary consequence [1]. Vertical social transmission (i.e., learning from a biological
39 parent) closely follows genetic inheritance patterns. For example, the diets and/or foraging
40 strategies of offspring have been shown to resemble that of their mother in a wide range of
41 mammalian taxa (e.g. orangutans *Pongo pygmaeus wurmbii*, sea otters *Enhydra lutris nereis*
42 and bottlenose dolphins *Tursiops* sp.) [2-4]. All three species share prolonged maternal
43 dependence and there is overlap between nursing and offspring-foraging during
44 development, providing opportunities for social learning [2, 3, 5].

45 Social transmission can affect the evolutionary outcomes of genetic transmission and vice
46 versa. Social transmission may change selection pressure on genes [gene-culture
47 coevolutionary theory, 6], as for example documented for the spread of lactose tolerance in
48 adult humans [7] and proposed for hundreds of human genes [8]. Further, individual
49 learning capacity or the exposure to a socially learned trait can correlate with specific
50 genetic marker systems [6]. For instance, in vertical social transmission, patterns of
51 transmission of the socially learned trait and uniparentally inherited genetic units, such as
52 mitochondrial DNA (mtDNA), and the Y chromosome, may be closely correlated. If the
53 socially learned trait increases its bearer's fitness, population frequencies of these correlated
54 genetic units will increase, even if there is no other active selection on genetic units. This
55 phenomenon is called "cultural hitchhiking" [9] and was posed as a possible explanation for
56 the low genetic diversity in human mtDNA and Y chromosome sequences [10] and the low
57 mtDNA diversity in matrilineal whales [9]. However, stochastic modelling has shown that a

58 reduction of genetic diversity is not a necessary consequence of parallel transmission of
59 genes and socially-learned phenotypes [10, 11].

60 Cultural hitchhiking might occur with little or no fitness differences if there is fine-scale
61 geographic population structure that relates to the cultural trait, as we discuss below. There
62 are two general types of geographic population structure: continuous clines or sharp
63 boundaries [12]. Genetic and geographic distances often correlate in natural populations
64 [i.e. isolation-by-distance, e.g., 13]. On the other hand, geographic features resulting in
65 habitat fragmentation or behavioural patterns limiting movement and genetic exchange,
66 may lead to discontinuous genetic structuring of populations. There is ample evidence for
67 structuring by geographical boundaries [14]. However, there is limited evidence for
68 geographic structuring of genetic variation due to behaviour and even less evidence due to
69 cultural behaviour. Possible mechanisms include assortative mating and microhabitat
70 specialisation. For example, two species of cichlid fish (*Amphilophus xiloaensis* and *A.*
71 *sagittae*) preferentially mate with a partner of the same colour morph, which has led to
72 genetic differentiation between morphs [15]. Moreover, in killer whales (*Orcinus orca*),
73 socially transmitted foraging specialisations within an ecosystem were proposed to have led
74 to sympatric ecotypes in the absence of physical barriers [16, 17].

75 To date, studies describing the influence of social transmission or culture on genetic
76 structure or selection have focused on genetic variation between populations [8]. Here, we
77 provide an example of how social transmission may drive genetic structure within a single
78 dolphin population.

79 Thirteen foraging strategies have been described for bottlenose dolphins in the eastern gulf
80 of Shark Bay (ESB), Western Australia [3]. Individual females have been observed to engage

81 in one to seven of these strategies [3], some of which are only observed in specific habitats.
82 The most prominent foraging strategies observed only in shallow water include beach
83 hunting, bottom grubbing and kerplunking [18, 19]. Foraging strategies observed in deep
84 water include 'sponging' [20-23]. The question of whether sponging is a socially transmitted
85 behaviour has been studied extensively in Shark Bay in the past [3, 20, 21, 24-28]. These
86 studies used different approaches (i.e. genetics, individual follows, survey data, network-
87 based approaches, individual based modelling or combinations thereof) to infer social
88 transmission of tool use in this population.

89 Sponging dolphins ('spongers' hereafter) carry conical marine sponges on their rostrums,
90 most likely to protect them while foraging for prey hiding in the substrate [21, 27]. The
91 behaviour is almost exclusively vertically transmitted from mothers to their offspring
92 through social learning [3, 20, 24]. With one exception, spongers in ESB share a maternally
93 inherited mtDNA haplotype [24]. Spongers in ESB are also biparentally more closely related
94 than expected by chance [24]. Haplotype sharing between sponging individuals is expected
95 in cases of strong vertical social transmission [24].even though a genetic basis for sponging
96 has been ruled out [24, 25, 28], Sponging has also been documented in the western gulf of
97 Shark Bay (WSB) [23, 25, 29]. The ESB and WSB study sites are approximately 120 km apart
98 by sea divided by a prominent peninsula and there is no direct evidence of dispersal
99 between the sites [30]. In both gulfs, sponging is almost exclusively limited to deep channels
100 where sponges occur [22, 23]. However, not all individuals inhabiting deep channels use
101 sponges to forage. Therefore, a purely ecological explanation for sponging would not
102 account for the heterogeneity of foraging strategies observed in these deep channels [22,
103 24].

104 This study focuses on the fine-scale population genetic structure of bottlenose dolphins in
105 WSB. We investigated the relationship between maternal relatedness, habitat and
106 behaviour. Our study significantly extends previous studies in that it explores the possibility
107 of social transmission shaping genetic structure in an animal population. This is a very
108 important advance because, to our knowledge, our study provides the first example of the
109 potential impact of social learning on within-population structure in non-human animals (as
110 opposed to its influence on between-population structure [15-17]). Briefly, we observed
111 striking spatial homogeneity in mtDNA, in contrast to other parts of the bay where
112 haplotype distribution was much more heterogeneous [30]. We aimed to test the hypothesis
113 that vertical social transmission of a habitat-dependent trait can lead to fine-scale genetic
114 structure. To do this we modelled the possible process using an individual-based model
115 based on empirical data. We used this model to investigate whether the observed
116 homogenous mtDNA haplotype structure could be a result of a random process, or
117 alternatively, would require that transmission patterns of a socially learned trait and
118 uniparentally inherited genetic units correlate, as is already known in this case [3, 20, 21, 24-
119 28]. Briefly, the model showed, that the observed geographic structuring of mtDNA
120 haplotypes is only possible if genetic and social transmission are correlated.

121

122 **Material and methods**

123 *(a) Study site and data collection*

124 Shark Bay is located on the west coast of Australia, 850 km north of Perth. It consists of an
125 eastern and western gulf, divided by Peron Peninsula. Our main study area is off the
126 township of Useless Loop in the western gulf and consists of approximately 260 km², about
127 five times the size of the average home range of an adult ESB bottlenose dolphin [31]. We
128 will refer to our main study area as WSB. WSB is characterized by channels (> 10 m deep)
129 dividing shallow water areas that are closer to land (figure 1) [23].

130 We carried out systematic photo-identification and behavioural surveys on bottlenose
131 dolphins from a 5.4 m boat with an outboard motor. Dolphin group composition [10 m chain
132 rule, 32], water depth, GPS and predominant dolphin activity (forage, rest, travel, social or
133 unknown) were recorded. For our analyses we did not discriminate between these activities.
134 Dolphins were individually identified by natural markings and the shape of their dorsal fin
135 [33]. Biopsy sampling [34] was conducted under a License to use and/or supply Animals for
136 Scientific Purposes from the Western Australian Department of Environment and
137 Conservation, and ethics approval was obtained from the University of New South Wales
138 (08/33B) and the University of Zurich.

139 A sponger was defined as a dolphin which was seen sponging at least twice [20]. We were
140 able to use water depth as a proxy for habitat because Tyne *et al.* [23] reported that in WSB,
141 sponges only occur in water deeper than 10 m, while seagrass occurs in shallow water (< 10
142 m).

143 *(b) Analyses*

144 The genetic sampling and laboratory methods are detailed in the supplement. We ran a
145 nested ANOVA in PASW Statistics 18 to test for significance of association between water
146 depth distribution of dolphins with different mtDNA haplotypes. Because multiple sightings
147 of the same individuals were included, individual dolphin identification (ID) was used as a
148 random factor. To exclude the possibility that the observed mtDNA haplotype distribution
149 might simply be an artefact of significant autosomal population structure, we tested
150 whether the observed segregation of mtDNA haplotypes and depth is reflected in
151 biparentally inherited microsatellite markers. Such autosomal structure would have to be
152 very strong, causing a severe restriction of gene flow, if it was to secondarily produce the
153 very clear mitochondrial geographic structuring. To investigate this, we ran a STRUCTURE
154 analysis (burn-in length of 10^5 and 10^6 Markov chain Monte Carlo steps) using mtDNA
155 haplotype as Locprior. [35-37]. The use of a Locprior model significantly increases the
156 chance to detect population structure even in weakly structured populations [37]

157 We analysed whether spongers shared an mtDNA haplotype more often than would be
158 expected by chance and whether they were biparentally more related than would be
159 expected by chance. Individuals (spongers and non-spongers) were included in these
160 analyses when they were sampled in an area specified by a 95% kernel utilization
161 distribution of locations where sponging had been observed by any animal [24] (figure S1).
162 The kernel was calculated in ArcMap 9.2 (ESRI) using the extension Home Range Tools [38].
163 Pairwise comparisons between the observed mtDNA haplotype distribution within spongers
164 were compared to pairwise comparisons of 10,000 times the number of sampled spongers
165 randomly drawn. The haplotype frequencies used as a basis for randomisation reflected the
166 frequencies calculated for the WSB dolphins. The randomizations were carried out in a
167 “macro” written in Microsoft Excel.

168 To test whether spongers are biparentally more related than expected by chance, we
169 calculated the average pairwise relatedness among spongers and compared it to the average
170 pairwise relatedness of the population. We calculated the Queller and Goodnight [39]
171 estimator of pairwise relatedness (R) in SPAGEDi [40]. In order to assess statistical
172 significance, we programmed a randomization test in MATLAB R2010a. Biparental genotypes
173 of all sampled individuals of the population were randomized and the average pairwise
174 relatedness among 22 or 15 (for WSB and ESB respectively, representing the number of
175 biopsy sampled spongers) randomly drawn genotypes were calculated. In the
176 randomization, the observed sex ratio of spongers was retained.

177 *(c) Individual-based model*

178 We used simulations to test the hypothesis that vertical social transmission of a habitat-
179 dependent trait can lead to fine-scale genetic structure using an individual-based model
180 based on empirical data. The model is described in detail in the supplement. We included
181 simulations of a null model with no fitness benefits for specialist, as well as one with fitness
182 benefits. Most simulations were run with 5% fitness benefits for specialists even though
183 empirical data suggest that fitness advantages of spongers may be larger than that, albeit
184 being non-significant [20].

185

186

187 **Results**

188 *(a) mtDNA haplotypes and habitat*

189 We found a strong and significant association between dolphin mtDNA haplotypes and
190 water depth in WSB. Dolphins with mtDNA haplotypes E or F were predominantly found in
191 deep water (> 10 m deep) and dolphins with haplotype H in shallow water (< 10 m, figures 1
192 and 2). The geographic segregation of mtDNA haplotypes in WSB was not reflected in
193 biparentally inherited genetic markers: STRUCTURE plots showed no indication of genetic
194 structure (figure S2). The depth distribution for both adult males and females with one of
195 the three common haplotypes E, F, and H differed significantly (tables 1 and 2), as revealed
196 by a nested ANOVA and a subsequent Tukey's HSD *post hoc* test, based on 90 dolphins (59
197 females and 31 males) and 756 sightings (mean number of sightings \pm SE/individual: $8.4 \pm$
198 0.68 , range = 1-33). Three other haplotypes (B, D, I) were rare (frequency < 0.03).

199 *(b) Genetic relatedness of spongers*

200 Forty spongers have been identified in WSB which represents about a quarter (25.9%) of
201 dolphins identified in deep water [29]. All 22 biopsied spongers in WSB share the same
202 mtDNA haplotype E, which is significantly different ($P < 0.001$) from what would be expected
203 by chance, based on the observed mtDNA haplotype frequencies in the population. The
204 sponger haplotype E in WSB is different from the haplotype found among ESB spongers
205 [haplotype H, 24]. All 22 sampled spongers were biparentally more closely related than the
206 population average ($R_{\text{spongers}} = 0.0259$, $R_{\text{population}} = -0.0104$, $N_{\text{population}} = 108$), although this
207 difference was not statistically significant ($P = 0.092$). We also re-ran the pairwise
208 relatedness analyses for ESB [24], including additionally genotyped microsatellite loci and
209 individuals. Our result confirms the previous publication [24]: spongers in ESB were found to

210 be biparentally more closely related than the population average and than expected by
211 chance ($R_{\text{spongers}} = 0.0712$, $R_{\text{population}} = -0.0044$, $N_{\text{spongers}} = 15$, $N_{\text{population}} = 238$, $P = 0.006$).

212 (c) *Individual-based model*

213 Simulations showed that fine-scale genetic structure based on mtDNA haplotypes can be
214 driven by vertically, socially transmitted, habitat-dependent traits (figures 3, S3). The
215 geographic segregation of haplotypes occurred even if the learning fidelity was not 100%
216 (figure S3), but in the absence of vertically, socially transmitted specialisations, no
217 geographic segregation of mtDNA haplotypes was observed (figure 3, fourth column).

218

219 **Discussion**

220 Through a combination of empirical data and individual-based modelling, we were able to
221 infer that the vertical cultural transmission of a foraging behaviour involving tools has led to
222 the clear separation of mtDNA haplotypes within our study area. The observed geographic
223 distribution of dolphins with different mtDNA haplotypes might represent a case of cultural
224 hitchhiking. To our knowledge, this is one of the first studies to provide evidence for fine-
225 scale geographic genetic structure driven by socially transmitted behaviour within a single
226 wild animal population.

227 In what circumstances would we predict correlations between habitats and mtDNA
228 haplotypes? We propose that four prerequisites must be met. First, a population must
229 exhibit vertically socially transmitted, habitat-dependent skills, such as foraging or predator
230 avoidance strategies. Second, philopatry must keep haplotypes localised, although if one sex
231 disperses, the pattern could still be found in the philopatric sex. Third, habitat differentiation
232 needs to be on a scale that is larger than an individual's home range. Fourth, in order for
233 habitat specialisations to drive genetic differentiation over small distances, these
234 specialisations must be stable over an individual's life time [41] and also present in following
235 generations. The clear separation of matriline and habitats we present here suggests that
236 habitat specialisations have been in place for many more generations than, for instance, the
237 four generations of spongers which have been observed in ESB so far.

238 The segregation of mtDNA haplotypes by habitat appears to be driven by vertically
239 transmitted foraging specialisations (and/or other behaviours leading to habitat
240 specialisation). The absence of fine-scale genetic structure based on biparentally inherited
241 markers shows that the correlation of maternally inherited mtDNA haplotype and water

242 depth cannot be explained solely by geographic separation. Of course, it was possible that
243 there is undetected autosomal structure, but it is most likely that this would be strong
244 enough to secondarily cause the extremely sharp mtDNA patterning that we observed. That
245 any autosomal structuring within our study area must be only weak is supported by two
246 points. Firstly, our use of Locprior ensures detection of relatively weak structure [37].
247 Secondly, the relatively weak autosomal genetic differentiation between ESB and WSB was
248 documented using a much smaller number of markers [30] than used in the current study.

249 Different depth preferences exhibited by dolphins with different mtDNA haplotypes coincide
250 with different habitats. Seagrass meadows are characteristic of shallow water, and conical
251 sponges only grow in water deeper than 10 m in WSB [23]. Shallow and deep habitats
252 intergrade and are not divided by a barrier or distance that could prevent dolphins from
253 moving between deep and shallow water. The absence of genetic structure based on
254 biparental genetic markers indicates that there is no mating barrier between deep and
255 shallow habitat and may be explained by the dispersal patterns of Shark Bay dolphins. Both
256 sexes are philopatric, with males expanding their natal range [30, 31]. In the WSB study site,
257 we observed the dolphins foraging, travelling, resting and socialising. Therefore the
258 geographic segregation of dolphins with different mtDNA haplotypes indicates that dolphins
259 stay in their natal habitat for all those activities (i.e. we do not know about mating). Male
260 bottlenose dolphins in Shark Bay cooperate with one another in pairs and trios to consort
261 females [42]. It appears that allied males can direct a female outside her regular home
262 range. Some sightings of females with haplotype H in deep water or haplotype E in shallow
263 water might be explained by the coercion of males. It is possible that these groupings
264 involved consorting of females by males, however, we did not confirm the consortships by
265 the standards of ESB [42]. In ESB, males have been observed consorting females never seen

266 in the study site before and well-studied females altered their depth-use during consortships
267 [43].

268 Although our study site is not likely to encompass the entire home ranges of some
269 individuals [44] included in this study, it is about five times the size of an average adult ESB
270 dolphin home range [31]. Accordingly, we expect our identification of each individual's
271 habitat usage to be suitably representative. In addition, the number of resightings per
272 individual (up to 33) suggests that the study site covers a great portion of the home ranges
273 of at least some animals.

274 Our individual-based simulations emphasise that social transmission of habitat-dependent
275 specialisations can lead to fine-scale genetic structure (figure 3, first and second column);
276 whereas this fine-scale structure is unlikely to be seen if there is no social transmission
277 (figure 3, fourth column). Social transmission of foraging strategies in WSB appears likely,
278 given the mtDNA haplotype sharing among WSB spongers, as occurs in ESB [24]. Our finding
279 that spongers in WSB are not more closely related than expected by chance provides further
280 evidence that sponging is transmitted socially, rather than genetically, and in a vertical
281 fashion. Furthermore, our simulations indicate that two specialisations are necessary, one
282 for the deep and one for the shallow habitat, in order to obtain the observed segregation of
283 two mtDNA haplotypes by habitat. It is conceivable that traits other than sponging are
284 vertically socially transmitted, such that matriline H and F also exhibit habitat specialisation.
285 Without any habitat specialisation, no mtDNA haplotype segregation is observed (figure 3,
286 fourth column).

287 Although all spongers sampled in WSB belong to one matriline, it is a different matriline
288 from that of spongers in ESB [24]. The presence of different sponging matriline in each gulf

289 can be explained most parsimoniously by the occurrence of at least two independent
290 innovations, one in each gulf. Alternatively, there could have been horizontal transmission
291 between matriline. However, this would have to be extremely rare to fit the patterns that
292 we observe.

293 What we have inferred for sponging may be just one example of a more widespread
294 phenomenon of interactions between culture and geographic structure of genetic variation.
295 Genetic differentiation on relatively small scales has been reported for many bottlenose
296 dolphin populations (both *T. aduncus* or *T. truncatus*) in all oceans. This structuring was
297 attributed largely to habitat specialisation and was found based on maternal and
298 biparentally inherited markers [45-49]. However, most of these studies (except [45]) gave no
299 indication of what the specialisations might be, or how they could have driven the genetic
300 structure. Furthermore, the geographic scales at which genetic differentiation have been
301 documented (tens to hundreds of km) in previous studies are much greater than those
302 reported here: deep and shallow habitats in WSB are separated by just tens of meters. The
303 largest delphinid, the killer whale, may present a similar case. Sympatric ecotypes were
304 shown to differ genetically and in their foraging behaviour [16, 50]. The genetic
305 differentiation between ecotypes based on microsatellites suggests that this separation may
306 be more stringent and/or has been in place for longer than in WSB.

307 A correlation between habitat and genetic structure has also been found in tool-using New
308 Caledonian crows (*Corvus moneduloides*) [51]. Rutz *et al.* [51] found biparental and maternal
309 (e.g. mtDNA haplotypes) genetic structure among three habitats (dry forest, farmland and
310 beachside habitat, < 10 km apart), in which different tool use has been observed. Restricted
311 dispersal may lead to localised occurrence of particular tool use [51]. In female mountain

312 gorillas (*Gorilla beringei beringei*) and western gorillas (*G. gorilla diehli*), Guschanski *et al.*
313 [52] reported a correlation between microsatellite-based genetic and geographic structure
314 (i.e. altitude). This correlation did not hold for males. In gorillas, both sexes disperse, but
315 females usually join a neighbouring group. Hence, female dispersal distance is limited and
316 often occurs within habitat types. The authors concluded that dispersal usually results in
317 females staying within the habitat into which they were born, probably due to food
318 preferences [52]. Unfortunately, data on maternally inherited markers was not presented.
319 The correlation between habitat and mitochondrial genetic structure in bottlenose dolphins
320 reported here holds for both adult females and males.

321 If vertical social transmission influences the fine-scale genetic structure of a population (e.g.
322 within WSB), it may have consequences for the genetic structure at larger scales (e.g.
323 between the two gulfs of Shark Bay). Since WSB matrilineal groups seem to stay in their natal
324 habitats, it is likely that they have behaviourally adapted to those habitats. If male and
325 female dolphins in Shark Bay are philopatric because of learnt foraging strategies, genetic
326 differentiation would occur – as observed - over small geographic distances. Limited
327 dispersal for both sexes could potentially cause inbreeding. Indeed, elevated levels of
328 inbreeding were measured in ESB, but the costs may not be high enough to outweigh
329 benefits of philopatry [53]. Philopatry bears benefits of habitat familiarity and the potential
330 for kin cooperation [54]. This study highlights that cultural forces can shape genetic
331 population structure outside humans and on a small geographic scale.

332

333

334 **Acknowledgements**

335 We are extremely grateful to Shark Bay Resources Inc. and the Useless Loop community for
336 their ongoing and extensive logistical support. We thank our field crew, including Kathrin
337 Bacher, Deirdre McElligott, Sina Kreicker, Dominik Noll and Richard Mardens. Thanks to
338 Celine Frere for helpful comments on the manuscript. This work was supported by the Sea
339 World Research and Rescue Foundation, National Geographic Society, Claraz-Schenkung, A.-
340 H. Schultz Stiftung and Julius-Klaus Stiftung.

341

342 **References**

- 343 [1] Bonduriansky, R. & Day, T. 2009 Nongenetic Inheritance and Its Evolutionary Implications.
344 *Annual Review of Ecology, Evolution, and Systematics* **40**, 103-125.
345 (doi:doi:10.1146/annurev.ecolsys.39.110707.173441).
- 346 [2] Jaeggi, A.V., Dunkel, L.P., Van Noordwijk, M.A., Wich, S.A., Sura, A.A.L. & Van Schaik, C.P.
347 2010 Social learning of diet and foraging skills by wild immature Bornean orangutans:
348 implications for culture. *American Journal of Primatology* **72**, 62-71.
349 (doi:10.1002/ajp.20752).
- 350 [3] Mann, J. & Sargeant, B.L. 2003 Like mother, like calf: The ontogeny of foraging traditions
351 in wild Indian Ocean bottlenose dolphins (*Tursiops* sp.). In *The biology of traditions: models*
352 *and evidence* (eds. D. Fragaszy & S. Perry), pp. 236-266. Cambridge, Cambridge University
353 Press.
- 354 [4] Estes, J., Riedman, M., Staedler, M., Tinker, M. & Lyon, B. 2003 Individual variation in
355 prey selection by sea otters: patterns, causes and implications. *Journal of Animal Ecology* **72**,
356 144-155. (doi:10.1046/j.1365-2656.2003.00690.x).

- 357 [5] Staedler, M.M., Tinker, M.T. & Estes, J. 2009 Individual variation in maternal care and
358 provisioning in the southern sea otter (*Enhydra lutris nereis*): Causes and consequences of
359 diet specialization in a top predator. In *18th Biennial Conference on the Biology of Marine*
360 *Mammals* (Quebec City, Canada).
- 361 [6] Feldman, M.W. & Laland, K.N. 1996 Gene-culture coevolutionary theory. *Trends in*
362 *Ecology & Evolution* **11**, 453-457. (doi:10.1016/0169-5347(96)10052-5).
- 363 [7] McCracken, R.D. 1971 Lactase Deficiency: An Example of Dietary Evolution. *Current*
364 *Anthropology* **12**, 479-517. (doi:10.1086/201234).
- 365 [8] Laland, K.N., Odling-Smee, J. & Myles, S. 2010 How culture shaped the human genome:
366 bringing genetics and the human sciences together. *Nat Rev Genet* **11**, 137-148.
367 (doi:10.1038/nrg2734).
- 368 [9] Whitehead, H. 1998 Cultural selection and genetic diversity in matrilineal whales. *Science*
369 **282**, 1708-1711. (doi:10.1126/science.282.5394.1708).
- 370 [10] Whitehead, H., Richerson, P.J. & Boyd, R. 2002 Cultural Selection and Genetic Diversity
371 in Humans. *Selection* **3**, 115-125. (doi:10.1556/Select.3.2002.1.9).
- 372 [11] Whitehead, H. 2005 Genetic diversity in the matrilineal whales: Models of cultural
373 hitchhiking and group-specific non-heritable demographic variation. *Marine Mammal*
374 *Science* **21**, 58-79. (doi:10.1111/j.1748-7692.2005.tb01208.x).
- 375 [12] Manel, S., Schwartz, M.K., Luikart, G. & Taberlet, P. 2003 Landscape genetics: combining
376 landscape ecology and population genetics. *Trends in ecology & evolution (Personal edition)*
377 **18**, 189-197. (doi:10.1016/S0169-5347(03)00008-9).
- 378 [13] Pogson, G.H., Taggart, C.T., Mesa, K.A. & Boutilier, R.G. 2001 Isolation by distance in the
379 atlantic cod, *Gadus morhua*, at large and small geographic scales. *Evolution* **55**, 131-146.
380 (doi:10.1111/j.0014-3820.2001.tb01279.x).

- 381 [14] Storfer, A., Murphy, M.A., Spear, S.F., Holderegger, R. & Waits, L.P. 2010 Landscape
382 genetics: where are we now? *Molecular Ecology* **19**, 3496-3514. (doi:10.1111/j.1365-
383 294X.2010.04691.x).
- 384 [15] Elmer, K.R., Lehtonen, T.K. & Meyer, A. 2009 Color assortative mating contributes to
385 sympatric divergence of neotropical cichlid fish. *Evolution* **63**, 2750-2757.
386 (doi:10.1111/j.1558-5646.2009.00736.x).
- 387 [16] Morin, P.A., Archer, F.I., Foote, A.D., Vilstrup, J., Allen, E.E., Wade, P., Durban, J.,
388 Parsons, K., Pitman, R., Li, L., et al. 2010 Complete mitochondrial genome phylogeographic
389 analysis of killer whales (*Orcinus orca*) indicates multiple species. *Genome Research* **20**, 908-
390 916. (doi:10.1101/gr.102954.109).
- 391 [17] Hoelzel, A.R. & Dover, G.A. 1991 Genetic differentiation between sympatric Killer whale
392 populations. *Heredity* **66**, 191-195. (doi:10.1038/hdy.1991.24).
- 393 [18] Connor, R.C., Heithaus, M.R., Berggren, P. & Miksis, J.L. 2000 "Kerplunking": Surface
394 fluke-splashes during shallow-water bottom foraging by bottlenose dolphins. *Marine*
395 *Mammal Science* **16**, 646-653. (doi:10.1111/j.1748-7692.2000.tb00959.x).
- 396 [19] Sargeant, B.L., Mann, J., Berggren, P. & Krützen, M. 2005 Specialization and
397 development of beach hunting, a rare foraging behavior, by wild bottlenose dolphins
398 (*Tursiops* sp.). *Canadian Journal of Zoology-Revue Canadienne De Zoologie* **83**, 1400-1410.
399 (doi:10.1139/z05-136).
- 400 [20] Mann, J., Sargeant, B.L., Watson-Capps, J., Gibson, Q.A., Heithaus, M.R., Connor, R.C. &
401 Patterson, E. 2008 Why do dolphins carry sponges? *PLoS ONE* **3**.
402 (doi:10.1371/journal.pone.0003868).

- 403 [21] Smolker, R., Richards, A., Connor, R., Mann, J. & Berggren, P. 1997 Sponge carrying by
404 dolphins (Delphinidae, *Tursiops* sp.): A foraging specialization involving tool use? *Ethology*
405 **103**, 454-465. (doi:10.1111/j.1439-0310.1997.tb00160.x).
- 406 [22] Sargeant, B.L., Wirsing, A.J., Heithaus, M.R. & Mann, J. 2007 Can environmental
407 heterogeneity explain individual foraging variation in wild bottlenose dolphins (*Tursiops* sp.)?
408 *Behavioral Ecology and Sociobiology* **61**, 679-688. (doi:10.1007/s00265-006-0296-8).
- 409 [23] Tyne, J., Loneragan, N., Kopps, A.M., Allen, S.J., Krützen, M. & Bejder, L. 2012 Ecological
410 characteristics contribute to sponge distribution and tool use in bottlenose dolphins
411 (*Tursiops* sp.). *Marine Ecology Progress Series* **444**, 143-153. (doi:10.3354/meps09410).
- 412 [24] Krützen, M., Mann, J., Heithaus, M.R., Connor, R.C., Bejder, L. & Sherwin, W.B. 2005
413 Cultural transmission of tool use in bottlenose dolphins. *Proceedings of the National*
414 *Academy of Sciences of the United States of America* **102**, 8939-8943.
415 (doi:10.1073/pnas.0500232102).
- 416 [25] Bacher, K., Allen, S., Lindholm, A., Bejder, L. & Krützen, M. 2010 Genes or Culture: Are
417 Mitochondrial Genes Associated with Tool Use in Bottlenose Dolphins (*Tursiops* sp.)?
418 *Behavior Genetics* **40**, 706-714. (doi:10.1007/s10519-010-9375-8).
- 419 [26] Mann, J., Stanton, M.A., Patterson, E.M., Bienenstock, E.J. & Singh, L.O. 2012 Social
420 networks reveal cultural behaviour in tool-using dolphins. *Nat Commun* **3**, 980.
421 (doi:10.1038/ncomms1983).
- 422 [27] Patterson, E.M. & Mann, J. 2011 The Ecological Conditions That Favor Tool Use and
423 Innovation in Wild Bottlenose Dolphins (*Tursiops* sp.). *PLoS ONE* **6**, e22243.
424 (doi:10.1371/journal.pone.0022243).

- 425 [28] Kopps, A.M. & Sherwin, W.B. 2012 Modelling the emergence and stability of a vertically
426 transmitted cultural trait in bottlenose dolphins. *Animal Behaviour* **84**, 1347-1362.
427 (doi:10.1016/j.anbehav.2012.08.029).
- 428 [29] Kopps, A., Krützen, M., Allen, S., Bacher, K. & Sherwin, W. in press Characterizing the
429 socially transmitted foraging tactic 'sponging' by bottlenose dolphins (*Tursiops* sp.) in the
430 western gulf of Shark Bay, Western Australia. *Marine Mammal Science*.
431 (doi:10.1111/mms.12089).
- 432 [30] Krützen, M., Sherwin, W.B., Berggren, P. & Gales, N. 2004 Population structure in an
433 inshore cetacean revealed by microsatellite and mtDNA analysis: Bottlenose dolphins
434 (*Tursiops* sp.) in Shark Bay, Western Australia. *Marine Mammal Science* **20**, 28-47.
435 (doi:10.1111/j.1748-7692.2004.tb01139.x).
- 436 [31] Tsai, Y.-J.J. & Mann, J. 2012 Dispersal, philopatry, and the role of fission-fusion dynamics
437 in bottlenose dolphins. *Marine Mammal Science*, no-no. (doi:10.1111/j.1748-
438 7692.2011.00559.x).
- 439 [32] Smolker, R.A., Richards, A.F., Connor, R.C. & Pepper, J.W. 1992 Sex-Differences in
440 Patterns of Association among Indian-Ocean Bottle-Nosed Dolphins. *Behaviour* **103**, 38-69.
- 441 [33] Würsig, B. & Würsig, M. 1977 The Photographic Determination of Group Size,
442 Composition, and Stability of Coastal Porpoises (*Tursiops truncatus*) *Science* **198**, 755-756.
443 (doi:10.1126/science.198.4318.755).
- 444 [34] Krützen, M., Barre, L.M., Möller, L.M., Heithaus, M.R., Simms, C. & Sherwin, W.B. 2002 A
445 biopsy system for small cetaceans: Darting success and wound healing in *Tursiops* spp.
446 *Marine Mammal Science* **18**, 863-878. (doi:10.1111/j.1748-7692.2002.tb01078.x).
- 447 [35] Pritchard, J.K., Stephens, M. & Donnelly, P. 2000 Inference of Population Structure Using
448 Multilocus Genotype Data. *Genetics* **155**, 945-959.

- 449 [36] Falush, D., Stephens, M. & Pritchard, J.K. 2003 Inference of Population Structure Using
450 Multilocus Genotype Data: Linked Loci and Correlated Allele Frequencies. *Genetics* **164**,
451 1567-1587.
- 452 [37] Hubisz, M.J., Falush, D., Stephens, M. & Pritchard, J.K. 2009 Inferring weak population
453 structure with the assistance of sample group information. *Molecular Ecology Resources* **9**,
454 1322-1332. (doi:10.1111/j.1755-0998.2009.02591.x).
- 455 [38] Rodgers, A.R. & Carr, A.P. 2005 HRT: Home Range Tools for ArcGIS. (ed. C.o.N.F.E.R.
456 Ontario Ministry of Natural Resources). Thunder Bay, Ontario, Canada,
457 <http://www.blueskytelemetry.com/downloads.asp>.
- 458 [39] Queller, D.C. & Goodnight, K.F. 1989 Estimating Relatedness Using Genetic-Markers.
459 *Evolution* **43**, 258-275.
- 460 [40] Hardy, O.J. & Vekemans, X. 2002 SPAGEDi: a versatile computer program to analyse
461 spatial genetic structure at the individual or population levels. *Molecular Ecology Notes* **2**,
462 618-620. (doi:10.1046/j.1471-8286.2002.00305.x).
- 463 [41] Knudsen, R., Primicerio, R., Amundsen, P.-A. & Klemetsen, A. 2010 Temporal stability of
464 individual feeding specialization may promote speciation. *Journal of Animal Ecology* **79**, 161-
465 168. (doi:10.1111/j.1365-2656.2009.01625.x).
- 466 [42] Connor, R.C., Smolker, R.A. & Richards, A.F. 1992 Two Levels of Alliance Formation
467 among Male Bottle-Nosed Dolphins (*Tursiops* Sp). *Proceedings of the National Academy of*
468 *Sciences of the United States of America* **89**, 987-990. (doi: 10.1073/pnas.89.3.987).
- 469 [43] Watson, J. 2005 Female mating behavior in the context of sexual coercion and female
470 ranging behavior of bottlenose dolphins (*Tursiops* sp.) in Shark Bay, Western Australia
471 Washington, DC, Georgetown University.

- 472 [44] Nicholson, K., Bejder, L., Allen, S.J., Krützen, M. & Pollock, K.H. 2012 Abundance, survival
473 and temporary emigration of bottlenose dolphins (*Tursiops* sp.) off Useless Loop in the
474 western gulf of Shark Bay, Western Australia. *Marine and Freshwater Research* **63**, 1059-
475 1068. (doi:http://dx.doi.org/10.1071/MF12210).
- 476 [45] Hoelzel, A.R., Potter, C.W. & Best, P.B. 1998 Genetic differentiation between parapatric
477 'nearshore' and 'offshore' populations of the bottlenose dolphin. *Proceedings of the Royal*
478 *Society of London. Series B: Biological Sciences* **265**, 1177-1183.
479 (doi:10.1098/rspb.1998.0416).
- 480 [46] Natoli, A., Birkun, A., Aguilar, A., Lopez, A. & Hoelzel, A.R. 2005 Habitat structure and
481 the dispersal of male and female bottlenose dolphins (*Tursiops truncatus*). *Proceedings of*
482 *the Royal Society B-Biological Sciences* **272**, 1217-1226. (doi:10.1098/rspb.2005.3076).
- 483 [47] Sellas, A.B., Wells, R.S. & Rosel, P.E. 2005 Mitochondrial and nuclear DNA analyses
484 reveal fine scale geographic structure in bottlenose dolphins (*Tursiops truncatus*) in the Gulf
485 of Mexico. *Conservation Genetics* **6**, 715-728. (doi:10.1007/s10592-005-9031-7).
- 486 [48] Parsons, K.M., Durban, J.W., Claridge, D.E., Herzing, D.L., Balcomb, K.C. & Noble, L.R.
487 2006 Population genetic structure of costal bottlenose dolphins (*Tursiops truncatus*) in the
488 northern Bahamas. *Marine Mammal Science* **22**, 276-298. (doi:10.1111/j.1748-
489 7692.2006.00019.x).
- 490 [49] Möller, L.M., Wiszniewski, J., Allen, S.J. & Beheregaray, L.B. 2007 Habitat type promotes
491 rapid and extremely localised genetic differentiation in dolphins. *Marine and Freshwater*
492 *Research* **58**, 640-648. (doi:10.1071/MF06218).
- 493 [50] Hoelzel, A.R., Hey, J., Dahlheim, M.E., Nicholson, C., Burkanov, V. & Black, N. 2007
494 Evolution of Population Structure in a Highly Social Top Predator, the Killer Whale. *Molecular*
495 *Biology and Evolution* **24**, 1407-1415. (doi:10.1093/molbev/msm063).

- 496 [51] Rutz, C., Ryder, T. & Fleischer, R. 2012 Restricted gene flow and fine-scale population
497 structuring in tool using New Caledonian crows. *Naturwissenschaften* **99**, 313-320.
498 (doi:10.1007/s00114-012-0904-6).
- 499 [52] Guschanski, K., Caillaud, D., Robbins, M.M. & Vigilant, L. 2008 Females Shape the
500 Genetic Structure of a Gorilla Population. *Current Biology* **18**, 1809-1814. (doi:DOI:
501 10.1016/j.cub.2008.10.031).
- 502 [53] Frère, C.H., Krützen, M., Kopps, A.M., Ward, P., Mann, J. & Sherwin, W.B. 2010
503 Inbreeding tolerance and fitness costs in wild bottlenose dolphins. *Proceedings of the Royal*
504 *Society B: Biological Sciences* **277**, 2667-2673. (doi:10.1098/rspb.2010.0039).
- 505 [54] Lawson Handley, L.J. & Perrin, N. 2007 Advances in our understanding of mammalian
506 sex-biased dispersal. *Molecular Ecology* **16**, 1559-1578. (doi:10.1111/j.1365-
507 294X.2006.03152.x).
- 508 [55] Koerper, H.C. & Stickel, E.G. 1980 Cultural Drift: A Primary Process of Culture Change.
509 *Journal of Anthropological Research* **36**, 463-469.

510

511 **Table 1:** Depth preference of WSB dolphins. Nested ANOVA comparing depth spectra of
 512 dolphins with different mtDNA haplotypes. Dolphin ID was used as a random factor nested
 513 within haplotypes.

		type III	df	mean	F	sig.
		sum of		square		
		squares				
Intercept	hypothesis	19563.33	1	19563.33	561.81	< 0.001
	error	3744.74	107.54	34.822		
Haplotype	hypothesis	1320.41	2	660.21	18.31	< 0.001
	error	3824.89	106.09	36.05		
ID(haplotype)	hypothesis	6815.46	87	78.34	13.34	< 0.001
	error	3912.11	666	5.87		

514

515

516 **Table 2:** Differences in depth preference for dolphins with different haplotypes (WSB). Tukey

517 HSD *Post Hoc* Test for nested ANOVA.

hap 1	hap 2	depth diff. [m]	std. error	sig.	95 % CI depth [m]	
					lower	upper
E	H	5.7*	0.19	< 0.001	5.2	6.1
E	F	1.5*	0.28	< 0.001	0.9	2.2
F	H	4.1*	0.28	< 0.001	3.5	4.8

518 hap: Haplotype, depth diff.: Mean depth difference, * indicates significance

519

520 **Figure legends**

521 **Figure 1:** Segregation of dolphin haplotypes by habitats in WSB. Survey locations of dolphins
522 with known mtDNA haplotypes are indicated. Survey colours represent haplotypes of
523 dolphins. Each sighting of a sampled dolphin was plotted, thus individuals can appear
524 multiple times. The sightings include all types of behaviour, including foraging, travelling,
525 resting, socialising and unknown. White areas represent shallow (< 10 m) and grey areas
526 represent deep (> 10 m) water. Insert illustrates the study area within Shark Bay; ESB
527 indicates the location of the eastern gulf of Shark Bay.

528


529 **Figure 2:** Depth preference of bottlenose dolphins with the three common haplotypes: H, F
530 and E (sponger haplotype). Boxes contain 50% of data points. Medians are indicated by black
531 horizontal lines within boxes. Whiskers delimit the lower and upper quartiles respectively.
532 Circles and asterisks represent outliers that are more than 1.5 and 3 times the box length
533 away from either end of the box, respectively.

534


535 **Figure 3:** MtDNA haplotype segregation by habitat in an individual-based model. Three
536 mtDNA haplotypes (Hap1, 2 and 3) and two habitat specialisations (sponging and strategy2)
537 were present. All spongers had Hap1 and all strategy2 individuals had Hap2. Top row:
538 number of different females per strategy (No IDs/strategy). Bottom row: proportion of
539 individuals with a particular mtDNA haplotype in deep water relative to all individuals with
540 this particular mtDNA haplotype. Proportions were calculated for every haplotype
541 separately. Error bars represent one standard error. Dashed lines indicate the observed

542 haplotype proportion in deep water for the three mtDNA haplotypes E, F and H in WSB.
543 Fitness benefits for specialists are shown below graphs. Because random cultural drift [55] is
544 a strong force counteracting the establishment of new innovation we indicated the
545 likelihood (%) of at least one specialist/strategy to persist for 100 time periods; this is shown
546 below the fitness benefits. Learning fidelities equalled 1 for daughters born to specialists for
547 the simulations shown here. Simulation results with various other learning fidelities and
548 fitness benefits are shown in figure S3. In the first column, individuals of both strategies had
549 the same fitness benefits. In the second column, only one strategy (sponging) was present,
550 in the third column, strategy2 was innovated 50 time periods after sponging and had a
551 higher fitness, and in the fourth column, there were no specialisations present.


552



553



554



555