

# New species of free-living aquatic nematodes from south-western Australia (Nematoda: Axonolaimidae and Desmodoridae)

M. Hourston<sup>1</sup> and R.M. Warwick<sup>2</sup>

<sup>1</sup> Centre for Fish and Fisheries Research, School of Biological Sciences and Biotechnology, Division of Science and Engineering, Murdoch University, South St, Perth, Western Australia 6150, Australia.

<sup>2</sup> Plymouth Marine Laboratory, Prospect Place, The Hoe, Plymouth PL1 3DH, United Kingdom.

**ABSTRACT** – Descriptions of 10 new species of free-living aquatic nematodes are presented. Four of the new species belong to the Axonolaimidae, i.e. *Ascolaimus australis*, *Parascolaimus brevisetus*, *Odontophora serrata* and *Parodontophora aurata*. The remaining six species belong to the Desmodoridae, i.e. *Bolbonema spiralis*, *Onyx cephalispiculus*, *O. potteri*, *Eubostrichus otti*, *Catanema australis* and *Leptonemella peronensis*. All 10 species were found in sediments taken from near-shore waters of either the Swan River Estuary or marine waters of the Perth region (Western Australia). *Ascolaimus australis* differs from its only congener *A. elongatus* in that its spicules are non-barbed and have rounded proximal cephalations. *Parascolaimus brevisetus* is different from its four other congeners by its short cephalic setae, which are less than one quarter the length of any other *Parascolaimus*, and the spicule cephalations are rounded rather than simple or spade-shaped. *Parodontophora aurata* is distinguished from six morphologically similar species by the unique arrangement ((4D-2V)2) and greater length (6 µm) of the opisthocephalic setation. *Odontophora serrata* is distinct from other *Odontophora* species in that each odontium bears seven dentate projections, which is more projections than similar species, *O. bermudensis* (5), *O. villoti* (3) and *O. paravilloti* (3). *Bolbonema spiralis* has spiral amphids with two turns, in contrast to those of *B. longisetosum*, which are only a single loop. *Onyx cephalispiculus* is most similar to *O. sagittarius* but has more robust spicules with large, bilobed cephalations. *Onyx potteri* is the only *Onyx* species to exhibit spicules with a distinct constriction and one of only two species with ten tubular precloacal supplements. The other, *O. dimorphus*, has sexually dimorphic amphids whereas *O. potteri* does not. *Eubostrichus otti* is most similar to *E. topiarus* but is only half the size. Cephalic setation is also divergent, with *E. otti* having two cephalic crowns, eight setae in the anterior, four in the posterior, whereas this arrangement is reversed for *E. topiarus*. *Catanema australis* is most similar to *C. exilis* but is distinguished from that species by the lack of prominent, tubular, postcloacal supplements and having only four pairs of normal setae on the tail. *Leptonemella peronensis* is most similar to *L. granulosa* but differs in its non-granular cephalic capsule, shorter tail and different circumcloacal setation (two precloacal, four postcloacal and one terminal double pair v. *L. granulosa*'s seven postcloacal pairs).

**KEYWORDS:** morphology, taxonomy.

## INTRODUCTION

During extensive studies of the free-living marine and estuarine nematode assemblages of south-western Australia, many previously undescribed species were found, ten of which are now formally described. These comprise two separate families, the Axonolaimidae Filipjev, 1918 (four species) and the Desmodoridae Filipjev, 1922 (six species) and encompass all of the new species which belong to those families that were encountered during these studies. The four representatives of the Axonolaimidae belong to four separate genera, i.e. *Ascolaimus*, *Parascolaimus*, *Odontophora* and *Parodontophora*. Of the six new species within Desmodoridae, one species belongs to the

Desmodorinae (*Bolbonema*), two species to the Spirininae (two *Onyx* species), and the final three species to the Stilbonematinae (one each of *Eubostrichus*, *Catanema* and *Leptonemella*).

Greenslade (1989) provides a comprehensive checklist of the nematode species recorded in Australian marine and estuarine environments prior to 1989 and also details the relevant taxonomic articles. Since that publication, a considerable number of taxonomic papers on Australian free-living nematodes have been produced (e.g. Greenslade and Nicholas 1991; Nicholas 1993, 1996, 2002; Blome and Riemann 1994; Gourbault and Vincx 1994; Stewart and Nicholas 1994, 1995; Nicholas and Trueman 2002). However, these latter publications consider only

species from the east and south coast of Australia, primarily as a reflection of the centers of Australian nematological research. The most recent descriptions of local Western Australian free-living nematode species are those of Inglis (1970, 1971), which describe several species of Cyatholaimidae and Enoplidae, respectively.

The specimens used to describe the 10 species below were collected during the studies detailed by Hourston et al. (2005, 2009). From those publications it is apparent that more than half of the species encountered could not be assigned to previously described species and in some cases, genera. In many instances this is because they represent new species, although it is acknowledged that some species have only been identified to a putative level and that the completion of the taxonomy is an ongoing task given the paucity of local relevant taxonomic research.

Taxonomic research is ongoing for the remaining unidentified species encountered during recent ecological studies in Western Australia (Hourston et al. 2005, 2009). The present paper represents the first of several taxonomic works required to complete this body of work.

## METHODS

The methodology for the collection, preservation, mounting and identification for individuals is detailed in Hourston et al. (2005). Reference slides for type material, were prepared by transferring specimens from mixed assemblage slides to individual species reference slides in glycerol, and sealed with slide sealant. All drawings were made using an Olympus CH40 compound light microscope with incorporated drawing tube. All type material for each of the 10 species has been deposited at the Western Australian Museum, Perth (WAM), the specimens indicated as 'other material examined' are currently held by Murdoch University. Further specimens may be lodged as they become available. The registration numbers for each type series are included in the descriptions below.

### Family Axonolaimidae Filipjev, 1918

### Subfamily Axonolaiminae Filipjev, 1918

### Genus *Ascolaimus* Ditlevsen, 1919

*Ascolaimus* Ditlevsen, 1919: 229.

#### TYPE SPECIES

*Bathylaimus filiformis* Ditlevsen, 1918 (junior synonym of *Monhystera elongata* Bütschli, 1874), by monotypy.

### *Ascolaimus australis* sp. nov.

Figures 1A–C; Table 1

#### MATERIAL EXAMINED

##### *Holotype*

**Australia: Western Australia:** ♂, Leighton Beach

(32°01'43"S, 115°44'51"E), 19 February 2001, M. Hourston (WAM V7450).

##### *Paratypes*

**Australia: Western Australia:** 1 ♀, 1 juvenile, Leighton Beach (32°01'43"S, 115°44'51"E), 19 February 2001, M. Hourston (WAM V7450).

#### DIAGNOSIS

*Ascolaimus australis* differs from its only other valid congener, *A. elongatus* Bütschli, 1874, in that the male spicules have simple not barbed distal tips, and rounded proximal cephalations as opposed to spade-shaped cephalations.

#### DESCRIPTION

##### *Male (holotype)*

*Body:* slender, medium length, predominantly transparent, distinct but fine annulations from posterior of amphid to caudal tip. *Tail:* short, conical. *Setation:* labial; 6, minute; cephalic; 4. Somatic; 4 files, over oesophagus, alternating insertion point between dorso-lateral and ventro-lateral files (Figure 1a.). Terminal; absent. *Buccal cavity:* double conical, weakly sclerotised, unarmed. *Oesophagus:* no bulbs/swellings, cardia indistinct. *Amphid:* large, circular loop, over posterior of buccal cavity. *Spicules:* short, evenly arcuate (90°), distally simple point, proximally cephalate with dorso-lateral twist to the cephalation. *Gubernaculum:* encases distal spicule tips, substantial dorso-caudal apophysis. *Testes:* paired, outstretched, right of intestine. *Supplementary organs:* absent.

##### *Female (paratype)*

Same diagnostic structures for body, setation, buccal cavity, amphids. Ovaries: paired, opposed, outstretched, right of intestine.

#### HABITAT

Type specimens were collected from sub- and supratidal sediments at a moderately high-energy beach in south-western Australia (Leighton Beach). Sediments at this location are predominantly biogenic and calcareous, medium grained and have very low organic content.

#### REMARKS

*Ascolaimus* contains only one other valid species, i.e. *A. elongatus*, which has been reported from many locations throughout Europe and North America in many separate articles, many of which provide detailed illustrations. While some details of *A. elongatus* differ among depictions, this species' spicules invariably have distinctive barbed distal ends and spade-shaped proximal cephalation (e.g. De Coninck and Schuurmans-Stekhoven 1932; Gerlach 1953; Warwick et al. 1998). Both distinctive characteristics of the above species are lacking in the presently described species. The spicules of *A. australis* have simple straight distal tips and rounded cephalations.




FIGURE 1 *Ascolaimus australis* sp. nov.: A, head and oesophageal region of holotype; B, caudal region of holotype; C, total body of paratype male.

**TABLE 1** Morphological measurements for the holotype of *Ascolaimus australis*, as well as those for the male and female paratype specimens. All measurements are in  $\mu\text{m}$  unless otherwise stated in tables 1–4 and 6–11. a = body length / maximum body diameter, b = body length/oesophagus length, c = body length/tail length, cbd = corresponding body diameter, from ant = distance from anterior, V (%) = percentage of vulva distance from anterior to total body length, ABD = anal body diameter. All abbreviations used are consistent across tables 1–4 and 6–11. Where sufficient paratype material was available, the ranges of values for the morphological measurements are given.

		Holotype	Paratype ♀	Paratype ♂
Body length (mm)		1.186	1.376	1.355
a		49.4	49.1	56.4
b		8.8	9.3	8.7
c		15.2	16.9	16.1
head diameter		10	10	13
Setae length	Cephalic	12	15	20
	Somatic	6	6	5
Buccal cavity length/diameter		13/4	14/5	13/5
Amphid:	from ant	5	8	9
	length/width	7/7	7/7	7/8
	cbd	12	14	18
Nerve ring:	from ant	72	80	88
	cbd	22	23	22
Oesophagus length		134	147	155
Max diameter		24	28	24
Vulva:	from ant	-	780	-
	V (%)	-	57	-
	cbd	-	28	-
Spicule length (arc)		34	-	31
ABD		25	20	22
Tail length		78	81	84

#### ETYMOLOGY

Since the only other *Ascolaimus* species (*A. elongatus*) has, to date, only been found in northern hemisphere localities, *A. australis* has been named in contrast, to reflect its southern hemisphere type locality.

#### Genus *Parascolaimus* Wieser, 1959

*Parascolaimus* Wieser, 1959: 66.

#### TYPE SPECIES

*Parascolaimus tau* Wieser, 1959, by original designation.

#### *Parascolaimus brevisetus* sp. nov.

Figures 2A–D, Table 2

#### MATERIAL EXAMINED

##### Holotype

**Australia: Western Australia:** ♂, Fish Market

Reserve (31°53'43"S, 115°57'33"E), 17 August 2005, M. Hourston (WAM V7451).

##### Paratypes

**Australia: Western Australia:** 2 ♂, 4 ♀, Fish Market Reserve (31°53'43"S, 115°57'33"E), 17 August 2005, M. Hourston (WAM V7451).

#### OTHER MATERIAL EXAMINED

**Australia: Western Australia:** 3 ♂, 3 ♀, Fish Market Reserve (31°53'43"S, 115°57'33"E), 20 August 2005, M. Hourston.

#### DIAGNOSIS

*Parascolaimus brevisetus* can be discriminated from its four described congeners in that its cephalic setae are only 5  $\mu\text{m}$  long, less than 1/4 the length of any other *Parascolaimus* species. Also the proximal ends of the spicules have distinct and rounded cephalations rather than simple, spade-shaped or weakly rounded.




FIGURE 2 *Parascolaimus brevisetus* sp. nov.: A, head and oesophageal region of holotype; B, caudal region of holotype; C, total body of holotype; D, total body of paratype female.

**TABLE 2** Morphological measurements for the holotype of *Parascolaimus brevisetus*, as well as the range for female and male paratype specimens.

	Holotype	Paratype ♀	Paratype ♂
Body length (mm)	1.035	0.935–0.045	0.801–1.035
a	41.4	30.4–34.8	36.3–41.4
b	6.0	7.3–8.3	5.8–6.7
c	11.0	9.3–9.7	9.2–11.0
Setae length			
Cephalic	5	4–5	4–5
Somatic	3	3–4	3–4
Buccal cavity Length/Diameter	15/4	14/4–16/4	12/4–15/5
Head diameter	9	6–8	8–9
Amphid:			
from ant	7	5–8	4–7
length/width	6/6	5/5–6/6	5/5–6/6
cbd	11	10–12	10–11
Oesophagus length	182	113–142	119–182
Nerve ring:			
from ant	88	71–86	73–88
cbd	20	18–20	17–20
Max diameter	25	27–31	20–25
Vulva:			
from ant	–	475–585	–
V (%)	–	51–56	–
cbd	–	27–31	–
Spicule length (arc)	26	–	25–26
ABD	22	17–19	17–22
Tail length	94	100–107	87–94

## DESCRIPTION

### *Male (holotype)*

*Body*: slender, small, colourless, fine and indistinct annulations visible on posterior of body. *Tail*: short, conical, slight terminal swelling. *Setation*: Cephalic; four, 5 µm from the anterior. Cervical; four pairs, dorso-lateral/ventro-lateral position. Somatic; sparse, short. Sub-ventral; 7 µm, 2 files on posterior body and tail. Terminal; absent. *Buccal cavity*: double conical, lightly sclerotised, labial ‘claws’ *sensu* Wieser (1959). *Oesophagus*: slight posterior bulb, distinct cardia. *Amphid*: circular loop, 7 µm from anterior, over posterior cone of buccal cavity. *Spicules*: slim, moderately arcuate, distinctive rounded cephalations. *Gubernaculum*: two parts, Dorsal part; with apophysis. Ventral part; paired, tubular, parallel to distal end of spicules. *Supplementary organs*: none visible, possible minute structures. *Testis*: large, single, outstretched, left of intestine. Spermatozoa; spherical, readily visible.

### *Female (paratypes)*

Same diagnostic structures for body; setation; buccal cavity, amphids. Ovaries: paired, opposed, outstretched, left of intestine.

## HABITAT

Individuals of this species are present in the mid to upper reaches of the Swan River Estuary. Type specimens were found in sediment types ranging from clay/mud to coarse sand, but more commonly occurred in the latter.

## REMARKS

*Parascolaimus brevisetus* is assigned to *Parascolaimus* on the basis of the ‘claw-like’ structures in the buccal cavity as described by Wieser (1959) and a complex, two-part gubernaculum. Within the genus, the present species can be easily discriminated from the other four described species in that the length of the cephalic setae (5 µm) are less than one quarter the length of that described for *P. tau* Wieser, 1959 (36–47 µm), *P. unguatus* Belogurov and Kartavtseva, 1975 (26–40 µm), *P. amphidoporus* Wieser, 1959 (c. 60 µm) and *P. proprius* Belogurov and Kartavtseva, 1975 (c. 40 µm). Furthermore, males of the present species possess distinctly rounded cephalate spicules whereas those of the other four species are either simple, spade-shaped or slightly cephalate. Additionally, the amphids of the present species (50% cbd) are smaller than those of *P. tau* (75% cbd), larger than those of *P. unguatus* (25% cbd) and *P. proprius* (30% cbd), and of a different shape to *P. amphidoporus*.

## ETYMOLOGY

*Parascolaimus brevisetus* has been named for one of the primary features that discriminates it from other *Parascolaimus* species, i.e. its relatively much shorter cephalic setae.

**Genus *Odontophora* Bütschli, 1874**

*Odontophora* Bütschli, 1874: 49.

## TYPE SPECIES

*Odontophora marina* Bütschli, 1874 by subsequent designation of Allgén, 1929.

***Odontophora serrata* sp. nov.**

Figures 3A–D; Table 3

## MATERIAL EXAMINED

*Holotype*

**Australia: Western Australia:** ♂, Chidley Point (32°01'03"S, 115°46'52"E), 6 February 2005, M. Hourston (WAM V7452).

*Paratypes*

**Australia: Western Australia:** 1 ♂, 1 ♀, Chidley Point (32°01'03"S, 115°46'52"E), 6 February 2005, M. Hourston (WAM V7452).

*Other material examined*

**Australia: Western Australia:** 2 ♂, 3 ♀, 5 juveniles, Penguin Island (32°18'25"S, 115°41'30"E), 20 July 2001, M. Hourston; 1 ♂, 2 ♀, 2 juveniles, Becher Point (32°22'8"S, 115°43'05"E), 20 July 2001, M. Hourston.

## DIAGNOSIS

*Odontophora serrata* is distinct from other *Odontophora* species in that each odontium bears 7 'dentate' projections, creating a serrated appearance. *O. bermudensis*, Jensen and Gerlach, 1976, *O. villoti* Luc and De Coninck, 1959 and *O. paravilloti* Blome, 1982 are the only other recorded species to bear dentate odontia, however, they have only 5, 3 and 3 projections per odontium, respectively.

## DESCRIPTION

*Male (holotype)*

*Body:* long, slender. *Cuticle:* mainly smooth, fine

**TABLE 3** Morphological measurements for the holotype of *Odontophora serrata*, as well as those for the female and male paratype specimens.

		Holotype	Paratype ♀	Paratype ♂
Body length (mm)		3.080	3.086	3.061
a		64.1		66.5
b		18.9	19.2	19.8
c		29.3	30.2	24.6
Head diameter		26	30	31
Setae length	Cephalic 1	36	32	37
	Cephalic 2	8	6	8
	Subcephalic	33	31	30
	Subterminal	64	-	65
Amphid:	from ant	6	2	3
	length/width	6/6	7/7	8/7
	cbd	25	30	27
Nerve ring:	from ant	115	114	113
	cbd	36	44	38
Oesophagus length		163	161	154
Max diameter		48	63	46
Vulva:	from ant	-	1700	-
	V (%)	-	55	-
	cbd	-	63	-
Spicule length (arc)		50	-	52
ABD		41	36	40
Tail length		105	102	124




FIGURE 3 *Odontophora serrata* sp. nov.: A, head of paratype female, external only; B, head of holotype, internal and external; C, total body of paratype female; D, caudal region of holotype.



annulations on tail. *Setation*: Labial; six, papillae. Anterior cephalic; six, long, insert level with posterior of amphid. Posterior cephalic; four, shorter, 4 µm behind anterior cephalic. Subcephalic; four, long, 4 µm behind posterior cephalic. Cervical; four files, extending along oesophagus. Somatic; absent. Caudal; thick, four submedian files. Subterminal; conspicuous, thick, 32 and 64 µm long. *Amphid*: small, round loop, extruded corpus gelatum. *Buccal cavity*: typical of *Odontophora*, conical, posterior walls cuticularised, contains odontia. *Odontia*: six, well developed, each has 7 projections, serrated appearance. *Inter-odontial plates*: cuticularised, triangular, anterior edge denticulate. *Oesophagus*: anterior/ posterior bulbs absent, cardia present. *Excretory pore*: level with subcephalic setae. *Spicules*: well developed, strongly arcuate, c. 90° angle, proximal end slightly cephalate and dorsally angled. *Gubernaculum*: cuticularised, conspicuous, long dorsal apophysis. *Supplementary organs*: 17, preanal, minute papillae with fine pores. *Testis*: single, outstretched, left of intestine.

#### Female (paratype)

Similar to male in body shape, anterior setation, amphid, buccal cavity. *Caudal setation*: sparser, conspicuous subterminal setae absent. *Ovaries*: paired, opposed, outstretched, left of intestine, mature eggs large (44 x 160 µm), elongate, conspicuous.

#### HABITAT

*Odontophora serrata* is found in calcareous sediments within both marine and estuarine environments of south-western Australia, but is usually only present in low numbers.

#### REMARKS

Jensen and Gerlach (1976) discussed the difficulties involved in separation of the large number of species in this genus, which differ only in minor features. They describe the 'denticulate projections' on the odontia of *O. bermudensis* Jensen and Gerlach, 1976 as peculiar and suggest that their presence is best seen on specimens on which the odontia are protruded, and that the form of the odontia in descriptions where this is not the case is unknown. However, the three pairs of lateral projections on each odontium of *O. serrata* are very obvious even on specimens where they are not protruded and we do not feel that they could have been overlooked if this species had been described earlier. There are three other species on which similar projections on the odontia have been identified, i.e. *O. villoti* Luc and De Coninck, 1959 and *O. paravilloti* Blome, 1982 which have one pair of projections on each odontium (three-lobed) and *O. bermudensis* which has two pairs (five-lobed). In contrast, *O. serrata* has three pairs of projections on each odontium (seven-lobed). Species in this morphologically similar sub-group also share the presence of a pair of long dorsolateral setae near the tail tip, but none so long as *O. serrata*.

#### ETYMOLOGY

This species had been named for its characteristic odontia, which have a distinctly serrated appearance due to the projections along their edges.

#### Genus *Parodontophora* Timm, 1963

*Parodontophora* Timm, 1963: 35.

#### TYPE SPECIES

*Pseudolella paraganulifera* Timm, 1952, by original designation.

#### *Parodontophora aurata* sp. nov.

Figures 4A–D, Tables 4, 5

#### MATERIAL EXAMINED

##### Holotype

**Australia: Western Australia:** ♂, Fish Market Reserve (31°53'43"S, 115°57'33"E), 8 February 2005, M. Hourston (WAM V7453).

##### Paratypes

**Australia: Western Australia:** 1 ♂, 2 ♀, Fish Market Reserve (31°53'43"S, 115°57'33"E), 8 February 2005, M. Hourston (WAM V7453).

##### Other material examined

**Australia: Western Australia:** 2 ♂, 2 ♀, Coode St Jetty (31°58'23"S, 115°52'08"E) 08 February 2005, M. Hourston.

#### DIAGNOSIS

*Parodontophora aurata* is one of six morphologically similar species. It is distinguished from those other five primarily by the unique arrangement (two dorsal files of three or four setae each and two ventral files of two setae) and greater length (6 µm) of the opisthocephalic setation.

#### DESCRIPTION

##### Holotype

*Body*: medium size, stout, dark amber in colour. *Tail*: conical. *Cuticle*: light annulations over entire body. *Setation*: Labial; 6, papillae. Cephalic; 4. Opisthocephalic; (*sensu* Wu et al. 2000), four files, dorso-lateral files 3–4 setae, ventro-lateral files 2 setae. Cervical; sparse, fine. Caudal; 2 subventral files. *Buccal cavity*: long, narrow, parallel and sclerotised walls. *Odontia*: six, simple, well developed. *Amphid*: 'crook-shaped', entirely over buccal cavity, shorter leg c. 70% of longer leg. *Oesophagus*: narrow, not surrounding buccal cavity, wider posterior to nerve ring. *Oesophageal bulb*: absent. *Renette cell*: large, 54 µm long, 36% oesophagus length. *Spicules*: slender, arcuate, split proximal cephalations. *Gubernaculum*: triangular, narrow dorso-caudal apophysis. *Supplementary organs*: absent. *Testes*: paired, outstretched, right of intestine.




FIGURE 4 *Parodontophora aurata* sp. nov.: A, head of holotype; B, total body of holotype; C, total body of paratype female showing renette cell (\*); D, caudal region of holotype.

**TABLE 4** Morphological measurements for the holotype of *Parodontophora aurata* sp. nov., as well as the range of measurements for female and male paratype specimens.

		Holotype	Paratype ♀	Paratype ♂
Body length (mm)		1.170	1.160–1.322	1.174–1.329
a		39	27.5–35.1	39.0–44.0
b		8.2	7.7–8.1	7.1–8.9
c		9.1	9.6–11.8	8.5–11.0
Head diameter		18	14–14	12–18
Setae length	Cephalic	8	7–8	7–8
	Opisthocephalic	6	5–6	5–6
	Cervical	6	5–5	5–6
Buccal cavity Length/Diameter		33/6	30/7–33/8	30/5–48/7
Amphid:	from ant	5	2–5	2–5
	dorsal arm	20	16–17	18–20
	ventral arm	14	13–14	14–15
	cbd	18	31–33	25–30
Nerve ring:	from ant	94	95–97	84–94
	cbd	29	28–31	25–30
Oesophagus length		142	150–162	142–167
Max diameter		30	33–45	28–30
Vulva:	from ant	–	620–680	–
	V (%)	–	50–54	–
	cbd	–	33–45	–
Spicule length (arc)		40	–	33–40
ABD		26	22–26	26–28
Tail length		129	98–135	120–140

#### *Female (paratypes)*

Similar to male, lower DeMan a ( $\delta=39$  v.  $\delta<35$ ).  
*Ovaries*: paired, opposed, outstretched, right of intestine.

#### HABITAT

Individuals of *Parodontophora aurata* are relatively common in sediments at upper estuarine/riverine locations in the Swan River Estuary. This species does not exhibit any marked preference for sediment type and is found in types ranging from gravels to fine clay/mud.

#### REMARKS

*Parodontophora aurata* can be distinguished from the majority of other congeneric species by the fact that the two arms of the amphid are not extremely different in length, and the longer arm does not extend past the base of the buccal cavity, such as the case for example, *P. wuleidaowanensis* Zhang, 2005. Of the remaining species, a group of morphologically similar species was identified by Wu et al. (2000) in which *P. aurata* should also now be included. The latter authors present a table summarising the distinguishing characteristics for

those five allied *Parodontophora* species. It is evident from Table 5 that although comparable values for most characters can be found among the other five species, the particular combination of characters found for *P. aurata* is unique. Those characters for which there are no exact match in the other five species are the length and arrangement of the opisthocephalic setae (see Diagnosis).

#### ETYMOLOGY

This species has been named for the distinct and consistent colouration of individuals, which ranges from light yellow through to dark amber.

#### Family Desmodoridae Filipjev, 1922

#### Subfamily Desmodorinae Filipjev, 1922

#### Genus *Bolbonema* Cobb, 1920

*Bolbonema* Cobb, 1920: 264.

#### TYPE SPECIES

*Bolbonema brevicolle* Cobb, 1920, by monotypy.

**TABLE 5** An adaptation of Table 2 as presented in Wu *et al.* (2000) with *P. aurata* included for comparison. In those situations where a character is similar for both *P. aurata* and another species, that character has been highlighted.

	<i>Parodontophora aurata</i>	<i>Parodontophora limnophilla</i>	<i>Parodontophora marisjaponici</i>	<i>Parodontophora breviampifida</i>	<i>Parodontophora quadristicha</i>	<i>Parodontophora marina</i>
Cephalic setae length (µm)	8	4	7	7-8	10.5	6-7.5
Opisthocephalic setae length (µm)	6	2	3.6-4.0	?	?	?
Opisthocephalic setae arrangement	(3/4D-2V)2	(3D-1V)2	(3D-1V)2	(1D-2V)2	(4D-3V)2	(3D-1V)2
amphidial dorsal/ventral arm	70%	50%	80%	< 50%	?	> 65%
Position of excretory pore	mid-stoma	anterior stoma	anterior stoma	anterior stoma	mid-stoma	mid-stoma
Renette cell/ oesophagus length	36%	34-47%	35-40%	40%	50-55%	45-57%




FIGURE 5 *Bolbonema spiralis* sp. nov.: A, head and oesophageal region of holotype; B, caudal region of holotype.

***Bolbonema spiralis* sp. nov.**

Figures 5A, B, Table 6

**MATERIAL EXAMINED**

*Holotype*

**Australia: Western Australia:** ♂, Mangles Bay (32°16'08"S, 115°41'34"E), 14 February 2004, R. Caccianiga (WAM V7454).

*Paratypes*

**Australia: Western Australia:** 6 ♂, 6 ♀, Mangles Bay (32°16'08"S, 115°41'34"E), 14 February 2004, R. Caccianiga (WAM V7454).

*Other material examined*

**Australia: Western Australia:** 1 ♂, 1 ♀, Chidley Point (32°01'03"S, 115°46'52"E), 6 February 2005, M. Hourston; 2 ♀, Mosman Waters (32°01'29"S, 115°46'26"E), 10 July 2005, M. Hourston.

**DIAGNOSIS**

*Bolbonema spiralis* differs from its described congeners by having distinctly spiral amphids with two turns, whereas the others have only a single loop. *B. spiralis* also differs from *B. longisetosum* as that former species has no pre- or post-cloacal supplements.

**DESCRIPTION**

*Male (holotype)*

*Body:* medium size, brown. *Tail:* short, conical. *Cuticle:* annulated, coarse/irregular anteriorly, fine/regular posteriorly. *Cephalic capsule:* distinct from body, smooth, abscission at 30% from anterior is semi-complete and interrupted by amphid. *Amphid:* large, spiral, 2 turns. *Setation:* Labial; 6, papillae. Cephalic; 4, level with amphid centre. Subcephalic; 4, at cephalic capsule base. Somatic; eight files regularly spaced radially, setae longitudinally alternate long (c. 22 µm) and short (c. 4 µm), extend from cephalic capsule base to posterior of cloaca. Terminal; absent. *Buccal cavity:* minute, unarmed. *Oesophagus:* short, narrow, non-muscular, oesophageal bulb distinct, cardia absent. *Spicules:* even arc, simple proximal cephalation. *Gubernaculum:* indistinct, small, simple, slender dorsal apophysis. *Supplementary organs:* indistinct or absent. *Testis:* single, outstretched, right of intestine.

*Female (paratypes)*

Similar to male. Comparable length, much greater girth. *Amphid:* 1.75 turns. *Ovaries:* paired opposed, reflexed, right of intestine.

**TABLE 6** Morphological measurements for the holotype of *Bolbonema spiralis* sp. nov., as well as the range of measurements for the female and male paratype specimens.

		Holotype	Paratype ♀	Paratype ♂
Body length (mm)		1.093	0.974–1.145	0.984–1.079
a		32.1	26.6–31.3	34.2–39.3
b		11.5	10.0–13.0	10.1–11.5
c		12.8	10.2–13.7	10.8–11.8
Head diameter		19	17–19	17–18
Setae length	Cephalic	9	10–11	9–12
	Subcephalic	4	4–5	4–5
Amphid:	from ant	5	3–5	4–6
	length/width	11/11	9/9–10/10	10/10–13/13
	cbd	19	17–19	17–18
Nerve ring:	from ant	56	55–64	58–63
	cbd	26	24–27	22–25
Oesophagus length		94	88–97	91–102
Max diameter		34	33–42	25–30
Vulva:	from ant	–	520–587	–
	V (%)	–	50–53	–
	cbd	–	33–38	–
Spicule length (arc)		48	–	43–57
ABD		18	16–18	18–20
Tail length		84	71–104	87–100

TABLE 7 Morphological measurements for the holotype of *Onyx cephalispiculus* sp. nov., as well as the range of measurements for the female and male paratype specimens.

	Holotype	Paratype ♀	Paratype ♂
Body length (mm)	1.204	1.215–1.268	1.232–1.284
a	20.7	21.7–22.0	25.1–27.2
b	5.1	5.7–6	6.3–6.4
c	11.7	11.8–13.0	9.7–14.0
Head diameter	34	33–33	29–32
Setae length			
	Labial	5–5	4–6
	Cephalic	13	12–15
	Subcephalic	7	6–11
Amphid:			
	from ant	1	0–1
	length/width	6/7	5/7–5/7
	cbd	24	12–22
Nerve ring:			
	from ant	115	102–112
	cbd	47	45–49
Oesophagus length	235	202–223	193–201
Max diameter	58	55–58	47–49
Vulva:			
	from ant	–	–
	V (%)	–	–
	cbd	–	–
Spicule length (arc)	69	–	65–75
No. of supplements	24	–	18–23
ABD	47	34–36	45–49
Tail length	112	93–107	88–128

#### HABITAT

All type specimens are from near-shore, sandy marine sediments. Additional individuals were recorded from the marine reaches of the Swan River Estuary, also in similar sediments.

#### REMARKS

Vershelde et al. (1998) ascribe two other species to the genus *Bolbonema*, namely *B. brevicolle* Cobb, 1920, and *B. longisetosum* Jensen, 1985. The latter species has longitudinally oval amphids describing a single loop in the male, with distinct pre- and post-cloacal supplements. *B. brevicolle* was rather inadequately described by Cobb (1920) on the basis of a single female specimen, the amphids of which are in the form of a single circular loop like those depicted by Jensen (1985) for a juvenile *B. longisetosum*. The synonymy of *B. brevicolle* and *B. longisetosum* cannot be dismissed on the basis of the published taxonomic literature alone.

#### ETYMOLOGY

The specific epithet refers to the spiral nature of the amphids, which are a primary diagnostic feature within the genus.

#### Subfamily Spiriniinae Gerlach and Murphy, 1965

##### Genus *Onyx* Cobb, 1891

*Onyx* Cobb, 1891: 146.

#### TYPE SPECIES

*Onyx perfectus* Cobb, 1891, by monotypy.

##### *Onyx cephalispiculus* sp. nov.

Figures 6A–C, Table 7

#### MATERIAL EXAMINED

##### *Holotype*

**Australia: Western Australia:** ♂, Mosman Waters (32°01'29"S, 115°46'26"E), 6 February 2005, M. Hourston (WAM V7455).

##### *Paratypes*

**Australia: Western Australia:** 7 ♂, 5 ♀, Mosman Waters (32°01'29"S, 115°46'26"E), 6 February 2005, M. Hourston (WAM V7455).




FIGURE 6 *Onyx cephalispiculus* sp. nov.: A, total body of holotype; B, head and oesophageal region of holotype; C, caudal region of holotype.



*Other material examined*

**Australia: Western Australia:** 10 ♂, 12 ♀, Mosman Waters (32°01'29"S, 115°46'26"E), 20 February 2005, M. Hourston; 2 ♂, 4 ♀, Chidley Point (32°01'03"S, 115°46'52"E), 6 February 2005, M. Hourston.

## DIAGNOSIS

*Onyx cephalispiculus*, and the previously described *O. sagittarius* Gerlach, 1950, share the largest number of precloacal supplements of any *Onyx* species (24). *O. cephalispiculus* differs from *O. sagittarius* as it has larger, more robust spicules which are more arcuate and have large, bilobed proximal cephalations.

## DESCRIPTION

*Male (holotype)*

*Body:* medium sized, red/brown. *Tail:* short, conical. *Cuticle:* very finely annulated. *Amphid:* simple single loop, extreme anterior placement. *Setation:* Labial; 6, setiform. Cephalic; 4, level with amphid. Subcephalic; 8, posterior to amphid. Cervical; four dense fields, very short and fine. Somatic; short, fine, sparse. Postcloacal; 2 subventral files, conspicuous cluster immediately post cloacal. Terminal; absent. *Buccal cavity:* strongly sclerotised forming a ring anterior to tooth. *Dorsal tooth:* well-developed, hollow, sclerotised. *Oesophagus:* narrow isthmus, anterior bulb well developed, posterior bulb very well developed, elongate, sclerotised inner lining, lacunae present *sensu O. macramphis* Blome and Riemann, 1994. *Nerve ring:* distinct, 50% oesophagus length. *Spicules:* distinct, arcuate, large bilobed cephalations. *Supplements:* 24, S-shaped, equally-sized. *Testis:* single outstretched, left of intestine.

*Female (paratypes)*

Similar to holotype, slightly less hirsute. *Ovaries:* paired, opposed, reflexed, left of intestine.

## HABITAT

All type specimens for *Onyx cephalispiculus* were collected subtidally in the predominantly marine lower Swan River Estuary. Sediment had heterogeneous grain size with low to moderate particulate organic content.

## REMARKS

An annotated list of *Onyx* species is presented in Blome and Riemann (1994) which presents a summary of the diagnostic characters of the eight species previously described for the genus, to which *O. cephalispiculus* (and *O. potteri* sp. nov., see later) should now be added.

*Onyx cephalispiculus* most resembles *Onyx sagittarius* in that the males of both species possess up to 24 strongly cuticularised, S-shaped, precloacal supplements; a feature which distinguishes these two species from all of the other known species of this

genus. *Onyx cephalispiculus* can be discriminated from *O. sagittarius* by the shapes of both the spicules and gubernaculum. The spicules of *O. cephalispiculus* are strongly arcuate and cephalated, with the cephalation being abscised laterally giving it a bilobed appearance while those of *O. sagittarius* are non-cephalated and not as strongly arcuate. Also, the gubernaculum and its dorsal apophysis are larger and more robust for the present species than depicted for *O. sagittarius*. The post cloacal, subventral cuticular elevations noted for *O. sagittarius* are absent on the present species. The fields of short setae in the cervical region of individuals of both sexes are a unique feature of this species.

## ETYMOLOGY

The species epithet 'cephalispiculus' reflects the distinctive cephalations of this species' spicules, which are unique within the genus *Onyx*.

***Onyx potteri* sp. nov.**

Figures 7A–D, Table 8

## MATERIAL EXAMINED

*Holotype*

**Australia: Western Australia:** ♂, Leighton Beach (32°01'43"S, 115°44'51"E), 30 January 2001, M. Hourston (WAM V7456).

*Paratypes*

**Australia: Western Australia:** 1 ♀, 4 juveniles, Leighton Beach (32°01'43"S, 115°44'51"E), 30 January 2001, M. Hourston (WAM V7456).

## DIAGNOSIS

*Onyx potteri* is the only *Onyx* species to exhibit spicules which are distinctly constricted as depicted in Figure 7c. Furthermore, except *O. dimorphus* Gerlach, 1963 it is only 1 of 2 species with 10 tubular precloacal supplements. *Onyx dimorphus* displays marked sexual dimorphism in amphid structure where as *O. potteri* does not.

## DESCRIPTION

*Male (holotype)*

*Body:* moderately long, slim, dark brown. *Cuticle:* fine distinct annulations, posterior of amphid to tail tip. *Amphid:* anterior placement, conspicuous, spiral, 2.75 turns. *Setation:* Cephalic; 2 crowns, 6 short anterior, 4 longer posterior. Sub-cephalic; 8, at posterior amphid edge. Cervical; irregular, long, over oesophagus. Somatic; short, fine, irregular, sparse. Terminal; absent. *Dorsal spines:* dorsal file, caudal region, regular wide spacing. Most very short (3 µm), two long on tail (20 µm). *Cloacal thorn:* present, minute. *Buccal cavity:* cup shaped, lightly sclerotised, distinct rugae. *Dorsal tooth:* well developed, hollow, sclerotised. *Oesophagus:* well




FIGURE 7 *Onyx potteri* sp. nov.: A, head and oesophageal region of holotype; B, head of paratype female; C, caudal region of holotype; D, total body of paratype male.

**TABLE 8** Morphological measurements for the holotype of *Onyx potteri*, as well as those for the female paratype and the range of measurements for the juvenile paratype specimens.

	Holotype	Paratype ♀	Paratype. Juv
Body length (mm)	1.112	0.776	0.660–0.839
a	39.7	26.8	25.3–33.5
b	8.1	5.6	5.8–7.0
c	13.7	12.9	10.4–12.1
Head diameter	20	21	18–23
Setae length:			
Labial	4	3	3–3
Cephalic	14	11	9–14
Subcephalic	9	6	6–10
Amphid:			
from ant	–	2	1–7
length/width	8/8	8/8	7/8–10/10
cbd	21	18	17–21
Nerve ring:			
from ant	63	67	60–63
cbd	23	28	21–25
Oesophagus length	137	138	106–125
Max diameter	28	29	25–26
Vulva:			
from ant	–	426	–
V (%)	–	55	–
cbd	–	29	–
Spicule length (arc)	50	–	–
ABD	27	23	20–21
Tail length	81	60	60–69

developed anterior/posterior bulbs, narrow isthmus, posterior bulb elongate, lacunae absent, non-sclerotised, cardia absent. *Spicules*: moderately sclerotised, slender, distinct angle, simple distal tip, distinct constriction, small proximal cephalations. *Supplements*: 10; posterior eight well defined, closely spaced, thick surrounding cuticle. anterior two inconspicuous, set apart, thin surrounding cuticle. *Gubernaculum*: oval, sclerotised posterior margin, surrounds spicule tips. *Testis*: single, left of intestine.

#### Female (paratype)

Similar to holotype. *Amphid*: similar dimensions, 2.5 turns. *Ovaries*: paired, opposed, reflexed, left of intestine. Uteri each contain a single large elongate egg (80 x 25 µm).

#### HABITAT

The type specimens were collected from nearshore marine waters along the lower west coast of Western Australia during the study by Hourston et al. (2005). The individuals were present in the calcareous sediments at the relatively high-energy site (Leighton Beach).

#### REMARKS

According to the annotated list of *Onyx* species in

Blome and Reimann (1994) only *O. dimorphus* also has ten tubular precloacal supplements. *Onyx dimorphus* is further described as having sexually dimorphic amphid structure, and depicted as relatively hirsute. This is in contrast to *O. potteri* in which the sexes have similar amphids, and are comparatively glabrous. Furthermore, there are no described species of *Onyx* that have spicules with the distinct constriction below the cephalation.

#### ETYMOLOGY

This species is dedicated to Prof. Ian Potter for his major contribution to marine and estuarine research in Western Australia.

#### Subfamily Stilbonematinae Chitwood, 1936

#### Genus *Eubostrichus* Greef, 1869

*Eubostrichus* Greef, 1869: 117.

#### TYPE SPECIES

*Eubostrichus filiformis* Greef, 1869, by subsequent designation of Stiles and Hassal (1905).




FIGURE 8 *Eubostrichus otti* sp. nov.: A, head and oesophageal region of holotype; B, head of paratype female; C, caudal region of holotype; D, total body of paratype female.

TABLE 9 Morphological measurements for the holotype of *Eubostrichus otti* sp. nov., as well as the range of measurements for the female and juvenile paratype specimens.

	Holotype	Paratype ♀	Paratype Juv
Body length (mm)	3.230	2.314–3.147	1.239–2.678
a	134.5	96.4–125.2	72.8–127.5
b	27.3	24.8–33.1	14.7–29.4
c	36.2	24.3–36.6	16.5–28.7
Head diameter	18	15–20	13–15
Setae length:			
Cephalic	15	15–20	8–14
Subcephalic 1	15	10–14	4–12
Subcephalic 2	14	8–12	9–14
Amphid:			
from ant	2	0–2	2–3
length/width	16/12	7/7–8/10	6/7–8/9
cbd	18	15–20	13–15
Nerve ring:			
from ant	64	63–71	54–70
cbd	22	21–24	16–23
Oesophagus length	118	93–109	88–93
Max diameter	24	24–30	17–21
Vulva:			
from ant	–	1142–1746	–
V (%)	–	52–61	–
cbd	–	24–30	–
Spicule length (arc)	43	–	–
ABD	24	16–18	12–14
Tail length	89	86–95	75–83

***Eubostrichus otti* sp. nov.**

Figures 8A–D, Table 9

**MATERIAL EXAMINED**

*Holotype*

**Australia: Western Australia:** ♂, Pelican Point (32°01'43"S, 115°44'51"E), 7 February 2005, M. Hourston (WAM V7457).

*Paratypes*

**Australia: Western Australia:** 1 ♀, Pelican Point (31°59'14"S, 115°49'28"E), 9 April 2005, M. Hourston (WAM V7457); 2 ♀, Applecross (32°00'38"S, 115°49'44"E), 9 April 2005, M. Hourston; 3 juveniles, Chidley Point (32°01'03"S, 115°46'52"E), 8 April 2005, M. Hourston.

**DIAGNOSIS**

*Eubostrichus otti* is most similar to *E. topiarus* Berger et al., 1996 but is approximately half the size. Also the cephalic setation is divergent, with *E. otti* having an anterior cephalic crown of eight setae and a posterior of four (8+4), which is the opposite to *E. topiarus* which has 4+8. The other similar species,

*E. longisetosus* Berger et al., 1996 and *E. africanus* Muthumbi et al., 1995, which both have both 4+4.

**DESCRIPTION**

*Male (holotype)*

*Body:* extremely elongate, slender, colourless. *Cuticle:* fine indistinct annulations, sickle-shaped epibiotic bacteria. *Amphid:* large, oval, single loop. *Setation:* Labial; obscured. Cephalic; eight long, anterior margin of amphid. Sub-cephalic; four, long, mid-amphid, sublateral. Cervical; eight, long. Caudal; all c. 8 µm long, 1 pair subventral-precloacal, five pairs subventral-postcloacal, one pair lateral-postcloacal, one double pair lateral-subterminal. Somatic; absent. *Buccal cavity:* minute, unarmed. *Oesophagus:* very narrow, non-muscular, distinct round posterior bulb, cardia absent. *Tail:* conico-cylindrical. *Spicules:* evenly arcuate, slim, well defined, proximal cephalations notched. *Gubernaculum:* small, narrow, conspicuous sickle-shaped dorsal apophysis. *Testis:* single, outstretched, right of intestine

*Female (paratype)*

Similar to holotype in cuticular ornamentation and epibionts. Differences to holotype. *Body:* substantially

longer. *Tail*: relatively longer. *ABD*: relatively narrower. *Amphid*: much smaller. *Cephalic setation*: posterior cephalic at posterior edge of amphid (Figures 8a v. 8b). *Caudal setation*: absent. *Ovaries*: paired, opposed, reflexed, right of intestine.

**HABITAT**

All of the type material for *Eubostrichus otti* was collected from sediment cores taken from the saline, lower/middle reaches of the Swan River Estuary in Melville Water. Individuals of this species were present in only a few samples and never in large quantities. Specimens were found in sediment types ranging from coarse biogenic sands to fine muds. All specimens found have been lodged as type material.

**REMARKS**

The bacterial coat of the holotype specimen covers only small areas of the body; however, inspection of the paratype material suggests that in life, the coat would have covered the majority of the body.

Notably, one of the juvenile paratype specimens hosted two different types of epibiont. It was covered by the crescent shaped epibionts on the posterior half of the body, while the anterior half was covered with a coccoid species, similar to those displayed by *Catanema australis*.

Of the members of the subfamily Stilbonematinae, *Eubostrichus otti* most resembles *E. topiarius*, sharing features such as amphid, spicule and gubernaculum

structure, cuticle ornamentation, crescent-shaped epibionts and caudal setation arrangement. However these two species are distinct, with the most conspicuous discriminating feature being the size of adults, which is up to 6 mm in *E. topiarius* while the largest individual found for the presently described species was only 3.2 mm. Additionally, the different arrangement of anterior setation is a notable divergence.

**ETYMOLOGY**

This species is named after Prof. Jörg Ott in recognition of his pioneering work with stilbonematids.

**Genus *Catanema* Cobb, 1920**

*Catanema* Cobb, 1920: 271.

**TYPE SPECIES**

*Catanema exilis* Cobb, 1920, by original designation.

***Catanema australis* sp. nov.**

Figures 9A–F, Table 10


**MATERIAL EXAMINED**

*Holotype*

**Australia: Western Australia:** ♂, Penguin Island (32°18'25"S, 115°41'30"E), 10 February 2001, M. Hourston (WAM V7458).

**TABLE 10** Morphological measurements for the holotype of *Catanema australis* sp. nov., as well as the range of measurements for the female and male paratype specimens.

		Holotype	Paratype ♀	Paratype ♂
Body length (mm)		2.166	2.750–3.217	2.110–2.529
a		77	88.7–103.7	78.1–97.2
b		26	23.9–33.5	23.7–24.5
c		25	28.9–36.5	25.7–30.7
Head diameter		19	21–23	19–22
Setae Length	Cephalic 1	15	12–14	11–11
	Cephalic 2	6	4–5	5–5
	Cervical	4	3–4	3–3
Nerve ring:	from ant	52	53–65	51–51
	cbd	24	25–28	22–25
Oesophagus length		82	96–115	89–103
Max diameter		28	30–31	26–27
Vulva:	from ant	–	1810–1490	–
	V (%)	–	54–57	–
	cbd	–	30–31	–
Spicule length (arc)		38	–	33–39
ABD		26	20–21	26–26
Tail length		87	86–95	82–84



**FIGURE 9** *Catanema australis* sp. nov.: A, head and oesophageal region of paratype female; B, head and cervical region of holotype showing GSOs; C, caudal region of holotype; D, total body of paratype female; E, paratype female midbody showing GSOs; F, caudal region of paratype female showing epibionts.

**Paratypes**

**Australia: Western Australia:** 3 ♂, 3 ♀, Penguin Island (32°18'25"S, 115°41'30"E), 10 February 2001, M. Hourston (WAM V7458).

**Other material examined**

**Australia: Western Australia:** 5 ♂, 5 ♀, Penguin Island (32°18'25"S, 115°41'30"E), 10 October 2001, M. Hourston (bulk mounted specimens).

**DIAGNOSIS**

*Catanema australis* is most similar to *C. exilis* Gerlach, 1963 in the location of the amphid and arrangement of the cephalic setae. The species are distinguished by *C. australis* possessing no supplements and only four pairs of normal setae or spines on the tail, whereas *C. exilis* has seven pairs of subventral, tubular, postcloacal supplements.

**DESCRIPTION****Male (holotype)**

**Body:** extremely elongate, slender, dark brown. **Cuticle:** indistinct ornamentation, obscured, coccoid epibiotic bacteria. **Amphid:** extreme anterior, obscured, corpus gelatum conspicuous. **Setation:** Anterior cephalic; 4. Subcephalic; 8. Cervical; four. Somatic; absent. Caudal; 4 unpaired ventral precloacal, 4 pairs subventral post cloacal. **Buccal cavity:** minute, unarmed. **Oesophagus:** muscular anterior bulb surrounding buccal cavity, narrow isthmus, small posterior bulb. **Glandular sensory organs (GSOs):** present, no associated spines, largest and most numerous in ventral cervical region. **Spicules:** evenly arcuate, short, very slender, proximally cephalate. **Gubernaculum:** small, block-shaped caudal apophysis. **Testis:** single, outstretched, right of intestine

**Female (paratypes)**

Similar to holotype in cuticular ornamentation and epibionts. Differences to holotype. **Body:** substantially longer. **Caudal setation:** absent. **GSOs:** large and numerous near vulva as well as cervical region. **Ovaries:** paired, opposed, reflexed, right of intestine

**HABITAT**

*Catanema australis* occurs in marine sediments, type specimens were collected at a low energy beach with high sedimentary organic material. See locality details in Hourston et al. (2005) under the incorrect designation '*Leptonemella* sp.'

**REMARKS**

Confusion exists as to the validity of the genus *Catanema* and whether its species should be reassigned to *Leptonemella*. Until the taxonomic issue is resolved, this species remains within *Catanema* since it meets the generic diagnostic criteria of an unarmed and minute buccal cavity enclosed in a muscular anterior oesophageal bulb, in contrast to *Leptonemella*, which

has no anterior oesophageal bulb.

*Catanema australis* is most similar to *C. exilis* in the extreme anterior position of the amphids (anterior to the four long cephalic setae) and the fact that the subcephalic and subamphidial seate (terminology of Platt and Zhang 1982) comprise a single circle of eight at the same level (collectively termed Subcephalic in the above description). The form of the amphids cannot be ascertained, but their location is indicated by the origin of the corpus gelatum, which is extruded in most specimens. In contrast, *C. exilis* has seven pairs of prominent subventral tubular supplements on the male tail, whereas *C. australis* has only four pairs of apparently normal setae or spines.

GSOs, similar to those described for *Eubostrichus topiarus* by Berger et al. (1996), are present but are not associated with hollow spines nor any visible setae as were those that latter species. GSOs are present over the majority of the body, but tend to be more numerous, larger and more closely spaced on the ventral surface in the cervical region of the male type specimens (Figure 9b v. 9e) than on the rest of the body, and additionally, near the vulva in female paratypes.

**ETYMOLOGY**

This species is named to reflect its southern hemispheric distribution in comparison to the northern hemispheric distribution of its described congeners.

**Genus *Leptonemella* Cobb, 1920**

*Leptonemella* Cobb, 1920: 244.

**TYPE SPECIES**

*Leptonemella cincta* Cobb, 1920, by monotypy.

***Leptonemella peronensis* sp. nov.**

Figures 10A–E, Table 11

**MATERIAL EXAMINED****Holotype**

**Australia: Western Australia:** ♂, Point Peron (32°16'08"S, 115°41'34"E), 14 February 2005, R. Caccianiga (WAM V7459).

**Paratypes**

**Australia: Western Australia:** 1 ♂, 1 ♀, Point Peron (32°16'08"S, 115°41'34"E), 14 February 2005, R. Caccianiga (WAM V7459).

**DIAGNOSIS**

*Leptonemella peronensis* is most similar to *L. granulosa* Boucher, 1975 but differs in its non-granular cephalic capsule and a shorter tail. Furthermore, *L. peronensis* has 2 precloacal, 4 postcloacal and one double pairs of setae near the tail tip, in contrast to *L. granulosa* which has seven post cloacal pairs of setae.



TABLE 11 Morphological measurements for the holotype of *Leptonemella peronensis* sp. nov., as well as those for the female paratype specimen.

		Holotype	Paratype ♀
Body length (mm)		2.040	3.113
a		68.1	103.7
b		20.6	27.1
c		25.2	30.2
Head diameter		24	26
Setae length:	Cephalic 1	23	23
	Cephalic 2	13	18
	Cervical	4	12
Nerve ring:	from ant	47	
	cbd	28	
Oesophagus length		99	115
Max diameter		30	30
Vulva:	from ant	–	2
	V (%)	–	53
	cbd	–	28
Spicule (arc)		40	–
ABD		27	24
Tail length		81	103

## DESCRIPTION

### Male (holotype)

*Body*: very elongate, slender, dark brown. *Cuticle*: coarse regular annulations. *Tail*: short, conical. *Amphid*: extreme anterior, obscured. *Cephalic capsule*: distinct, smooth. *Setation*: Labial; obscured. Anterior cephalic; four, 23 µm. Posterior cephalic, six, 13 µm. Cervical; eight, 4 µm, on cephalic capsule. Caudal; one pair subventral prelocal, four pairs subventral postcloacal, two pairs subdorsal post cloacal, one double pair subterminal. *Buccal cavity*: minute, unarmed. *Oesophagus*: narrow, poorly developed anterior and posterior bulbs. *Spicules*: strongly arcuate, cuticularised, distinct cephalation. *Gubernaculum*: narrow, parallel to posterior of spicules, apophysis absent. *Testis*: single, outstretched, right of intestine.

### Female (paratype)

Similar to holotype in body morphology, cuticular ornamentation, setation pattern. Differences to holotype. *Body*: longer (♀ = 3.1 mm v. ♂ = 2.04 mm). *DeMan a*: greater (♀ = 103 v. ♂ = 68). *Tail*: relatively longer. *Caudal setation*: absent. *Ovaries*: paired, opposed, outstretched, right of intestine.

## HABITAT

Type specimens were collected from a low energy, shallow, near-shore marine environment at Point Peron, south-western Australia. The sediment was relatively

coarse and calcareous, and contained appreciable amounts of seagrass detritus.

## REMARKS

In contrast to many other species of Stilbonematinae, the type specimens of the present species lack any sheath of epibionts. It is not implicit that this sheath does not occur on individuals of this species, only that it is absent for the described individuals in their preserved state.

This species is similar to *L. granulosa* in many respects. The only appreciable differences are the surface texture of the cephalic capsule, the tail length and the circum-cloacal setation. *Leptonemella peronensis* does not have the noticeably granular cephalic capsule which is characteristic of *L. granulosa*. Furthermore, *L. peronensis* has a shorter tail than that described for *L. granulosa*, particularly males where it is only 3 ABD compared to 5.6 ABD. The holotype of *L. granulosa* is depicted as having 7 pairs of short post-cloacal spines on the subventral surfaces and none anterior to the cloaca, whereas *L. peronensis* has two pre-cloacal pairs of setae, only four post-cloacal pairs and a double pair near the tail terminus.

## ETYMOLOGY

This species is named after its type locality, Point Peron, Western Australia.




FIGURE 10 *Leptonemella peronensis* sp. nov.: A, diagram of the arrangement of anterior cephalic (+), posterior cephalic (^), and cervical (\*) setation, not to scale. Setation symbology is consistent through the remainder of Figure 10; B, caudal region of holotype; C, head and oesophageal region of paratype female; D, head and oesophageal region of holotype; E, total body of paratype female.

## ACKNOWLEDGEMENTS

Gratitude is expressed to Michelle Wildsmith, Natasha Coen and Fiona Valesini as well as the undergraduate students of the Marine and Estuarine Biology course (Murdoch University) for help with sampling. Financial support was provided by the Fisheries Research and Development Corporation, Murdoch University and The Royal Society of London. RMW would like to acknowledge his honorary fellowship at the Plymouth Marine Laboratory and his adjunct professorship at Murdoch University.

## REFERENCES

- Allgén, C. (1929). Was ist die von Bütschli 1874 aufgestellte *Odontophora marina*? *Zoologischer Anzeiger* **81**: 305–309.
- Belogurov, O.I. and Kartavtseva, I.V. (1975). Diagnoses of the genera *Parascolaimus*, *Paradontophora*, *Synodontium*, *Synodontoides* of the subfamily Axonolaiminae with the description of some far eastern species. *Zoologicheskii Zhurnal* **54**: 1284–1292.
- Berger, E., Urbancik, W. and Ott, J.A. (1996). *Eubostrichus topiarus* sp. n., A new free-living, marine species of Stilbonematoinae (Nematoda: Desmodoridae) from a shallow subtidal sand bottom. *Nematologica* **42**: 521–536.
- Blome, D. (1982). Systematik der Nematoda eines Sandstrandes der Nordseeinsel Sylt. *Mikrofauna Meeresboden* **86**: 1–25.
- Blome, D. and Riemann, F. (1994). Sandy beach meiofauna of eastern Australia (Southern Queensland and New South Wales). III. Revision of the nematode genus *Onyx* Cobb, 1891, with a description of three new species (Nematoda : Desmodoridae). *Invertebrate Taxonomy* **8**: 1483–1492.
- Boucher, G. (1975). Nématodes des sables fins infralittoraux de al Pierre Noire (Manche occidentale) I. Desmodorida. *Bulletin du Muséum National d' Histoire Naturelle 3 Zoologie* **195**: 101–128.
- Bütschli, O. (1874). Zur Kenntnis der freilebenden Nematoden, insbesondere der des Kieler Hafens. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* **9**: 236–292.
- Chitwood, B.G. (1936). Some marine nematodes from North Carolina. *Proceedings of the Helminthological Society of Washington* **3**: 1–16.
- Cobb, N.A. (1891). *Onyx* and *Dipeltis* new nematode genera, with a note on *Dorylaimus*. *Proceedings of the Linnean Society of New South Wales* **6**: 143–158.
- Cobb, N.A. (1920). One hundred new nemas (type species of 100 new genera). *Contributions to a Science of Nematology* **9**: 217–343.
- De Coninck, L.A. and Schuurmans-Stekhoven, J.H. (1933). The freelifving marine nemas of the Belgian coast II. *Mémoires du Musée Royal D'Histoire Naturelle de Belgique* **58**: 1–163.
- Ditlevsen, H. (1919). Marine free-living nematodes from Danish waters *Videnskabelige Meddelelser Fra Dansk Naturhistorik Forening* **70**: 147–214.
- Filipjev, I. (1918). Free-living marine nematodes of the Sevastopol area (In Russian). *Trudy Osoboi Zoologicheskoi Laboratorii I Sevastopol'skoi Biologicheskoi Stantsii* **2**: 1–350.
- Filipjev, I. (1922). Encore sur les nématodes libres de la mer Noire. *Acta Instituti Agronomici Staurpolitani* **1**: 83–184.
- Gerlach, S.A. (1950). Über einige Nematoden aus der Familie der Desmodoriden. *Zoologischer Anzeiger (Klatt-Festschrift, Neue Ergebnisse und Probleme der Zoologie)* **45**: 178–198.
- Gerlach, S.A. (1953). Die Nematodenbesiedlung des Sandstrandes und des Küstengrundwassers an der italienischen Küste. I. Systematischer Teil. *Archivio Zoologico Italiano* **37**: 517–640.
- Gerlach, S.A. (1963). Freilebende Meeresnematoden von den Malediven II. *Kieler Meeresforschungen* **19**: 67–103.
- Gerlach, S.A. and Murphy, D.G. (1965). Die Familiengruppen-Namen der freilebenden Nematoda-Adenophora (Aphasmidia). *Zoologischer Anzeiger* **175**: 211–216.
- Gourbault, N. and Vincx, M. (1994). New species of *Parapinnanema* (Nematoda: Chromadoridae) are described, with a discussion of the genus. *Australian Journal of Marine and Freshwater Research* **45**: 141–159.
- Greif, R. (1869). Untersuchungen über einige merkwürdige Formen des Arthropoden- und Wurm-Typus. *Archiv für Naturgeschichte* **35**: 71–100.
- Greenslade, P. (1989). Checklist of free-living marine nematodes from Australia, Macquarie Island and Heard Island. *Records of the South Australia Museum* **23**: 7–19.
- Greenslade, P. and Nicholas, W.L. (1991). Some Thoracostomopsidae (Nematoda : Enoplida) from Australia, including descriptions of two new genera and diagnostic keys. *Invertebrate Taxonomy* **4**: 1031–1052.
- Hourston, M., Warwick, R.M., Valesini, F.J. and Potter, I.C. (2005). To what extent are the characteristics of nematode assemblages in nearshore sediments on the west Australian coast related to habitat type, season and zone? *Estuarine, Coastal and Shelf Science* **64**: 601–612.
- Hourston, M., Potter, I.C., Warwick, R.M., Valesini F.J. and Clarke, K.R. (2009). Spatial and seasonal variations in the ecological characteristics of the free-living nematode assemblages in a large microtidal estuary. *Estuarine Coastal and Shelf Science* **82**: 309–322.
- Inglis, W.G. (1970). Cyatholaimidae (Nematoda) from the coast of Western Australia. *Records of the South Australia Museum* **16**: 1–13.
- Inglis, W.G. (1971). Marine Enoplida (Nematoda) from Western Australia. *Transactions of the Royal Society of South Australia* **95**: 65–78.
- Jensen, A. and Gerlach, S.A. (1976). Three new marine nematodes from Bermuda. *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* **16**: 31–44.
- Jensen, P. (1985). The nematode fauna in the sulphide rich brine seep and adjacent bottom of the East Flower Garden, NW Gulf of Mexico. I. Chromadorida. *Zoologica Scripta* **14**: 247–263.
- Luc, M. and De Coninck, L.A. (1959). Travaux de la Station Biologique de Roscoff, LII. Nématodes libres marins de la région de Roscoff. *Archives de Zoologie Experimentale Générale* **98**: 103–165.
- Muthumbi, A.W., Verschelde, D. and Vincx, M. (1995). New Desmodoridae (Nematoda: Desmodoroidea): three new species from *Ceriops* mangrove sediments (Kenya) and one related new species from the North Sea. *Cahiers De Biologie Marine* **36**: 191–195.
- Nicholas, W.L. and Trueman, J.W.H. (2005). Biodiversity of marine nematodes in Australian sandy beaches from tropical and temperate regions. *Biodiversity and Conservation* **14**: 823–839.
- Nicholas, W.L. (1993). Two new species of nematode

- (Nematoda : Enoplida : Thoracostomopsidae) from Lake Alexandrina, South Australia. *Transactions of the Royal Society of South Australia* **117**: 163–170.
- Nicholas, W.L. (1996). *Robustnema fosteri* sp. gen. nov. (Xyalidae, Monhysterida, Nematoda), a common nematode of mangrove mudflats in Australia. *Transactions of the Royal Society of South Australia* **120**: 161–165.
- Nicholas, W.L. (2002). Ultrastructure of the head of *Okranema eileenae* Greenslade and Nicholas, 1991 (Thoracostomopsidae: Nematoda). *Journal of Nematology* **34**: 50–61.
- Platt, H.M. and Zhang, Z. (1982). New species of marine nematodes from Loch Ewe, Scotland. *Bulletin of the British Museum of Natural History* **42**: 227–246.
- Schuurmans-Stekhoven, J.H. and De Coninck, L.A. (1932). Zur Synonymie von *Ascolaimus elongatus* (Bütschli). *Zoologischer Anzeiger* **99**: 149–163.
- Stewart, A.C. and Nicholas, W.L. (1994). New species of Xyalidae (Nematoda : Monhysterida) from Australia. *Invertebrate Taxonomy* **8**: 91–115.
- Stewart, A.C. and Nicholas, W.L. (1995). *Manunema pectenophora* sp. nov. (Peresianidae, Leptolaimina), a nematode possessing unusual male supplementary organs. *Transactions of the Royal Society of South Australia* **119**: 163–169.
- Stiles, C.W. and Hassal, A. (1905). *The determination of generic types and a list of roundworm genera, with their original and type species*. Bureau of Animal Industry, Bulletin 79. U.S. Department of Agriculture, Beltsville, Maryland, 150 pp.
- Timm, R.W. (1952). *A Survey of the marine nematodes of Chesapeake Bay, Maryland*. Chesapeake Biology Laboratory Publication **95**: 1–70.
- Timm, R.W. (1963). Marine nematodes of the suborder Monhysterina from the Arabian Sea at Karachi. *Proceedings of the Helminthological Society of Washington* **30**: 34–49.
- Vershelde, D., Goubault, N. and Vincx, M. (1998). Revision of *Desmodora* with descriptions of new *Desmodorids* (Nematoda) from hydrothermal vents of the Pacific. *Journal of the Marine Biological Association of the United Kingdom* **78**: 75–112.
- Warwick, R.M., Platt, H.M. and Somerfield, P.J. (1998). *Free-Living Marine Nematodes: Part III. Monhysterids*, 53. Field Studies Council, Shrewsbury, U.K.
- Wieser, W. (1959). Freelifing marine nematodes and other small invertebrates of Puget Sound beaches. *University of Washington Publication of Biology* **19**: 1–17.
- Wu, J.H., Somerfield, P.J., Austen, M.C. and Liang, Y.L. (2000). The freelifing nematode genus *Parodontophora* Timm, 1963 (Nematoda: Axonolaimidae) is not exclusively marine: *Parodontophora limnophila* sp. nov. from freshwater in China. *Hydrobiologia* **431**: 205–210.
- Zhang, Z. (2005). Three new species of free-living marine nematodes from the Bohai Sea and Yellow Sea, China. *Journal of Natural History* **39**: 2109–2123.