

Special Focus Review

Locked nucleic acid as a novel class of therapeutic agents

Rakesh N. Veedu and Jesper Wengel*

Nucleic Acid Center; Department of Physics and Chemistry; University of Southern Denmark; Odense, Denmark

Abbreviations: LNA, locked nucleic acid; mRNA, messenger RNA; AONs, antisense oligonucleotides; HIV, human immunodeficiency virus; RNAi, RNA interference; siRNA, small interfering RNA; sisiRNA, small internally segmented interfering RNA; miRNA, microRNA; TAR, trans-activation responsive element; SELEX, systematic evolution of ligands by exponential enrichment

Key words: LNA, oligonucleotides, antisense, siRNA, miRNA, aptamers, RNA targeting

Locked Nucleic Acid (LNA) is a nucleic acid analogue with unprecedented binding affinity and excellent specificity toward complementary RNA and DNA oligonucleotides. The remarkable properties of LNA have led to applications within various gene silencing strategies both *in vitro* and *in vivo*. In the present review, we highlight the uses of LNA for regulation of gene expression with emphasis on RNA targeting.

Introduction

Nucleic acid derivatives have gained a lot of interest for the treatment of many diseases, and the first oligonucleotide to enter the clinic was Vitravene[®],¹ an antisense oligonucleotide for the treatment of cytomegalovirus infection. The use of chemically modified nucleic acids is needed as naturally occurring DNA or RNA have some limitations like poor RNA binding affinity, inefficient cellular uptake and very limited nuclease resistance. Among the numerous modifications known, LNA has shown broad usefulness within chemical biology.²⁻⁹

LNA: Structural Features and Key Properties

LNA (Fig. 1) contains a ribose ring which is locked by a O2'-C4'-methylene linkage, imposing conformational restriction to adopt an *N*-type sugar pucker.^{2-5,10} Structural investigation of LNA oligonucleotides by NMR spectroscopy revealed their similarities with natural nucleic acid duplexes, and confirmed the RNA mimicking structures adopted by LNA.^{11,12}

LNA offers key properties needed for successful therapeutic exploitation of oligonucleotides, including (1) unprecedented binding affinity towards RNA (and DNA), (2) excellent base pairing specificity, (3) high bio-stability (resistance towards nucleolytic degradation), (4) low toxicity (at least for many LNA




Figure 1. Two representations of an LNA monomer. The structure is shown schematically to the left, whereas the structure and its locked sugar ring conformation is shown in three dimensions to the right.

oligonucleotides) in animals and (5) convenient chemistry for manufacturing and modification.

LNA in Antisense Technologies

'Antisense' is a term normally used for single stranded nucleic acid based approaches that interfere with the processing of RNA in a sequence selective manner. For effective modulation of gene expression, the advantages of synthetic oligonucleotides are exploited through binding of antisense oligonucleotides (AONs) to a specific mRNA or pre-mRNA by Watson-Crick base pairing. Upon binding, the oligonucleotide can modulate RNA processing, inhibit translation, or promote degradation. Antisense-based gene silencing strategies using LNA-containing oligonucleotides have already been the topic of detailed reviews.^{9,13,14}

LNA-antisense experiments have largely been focused on mRNA inhibition by RNase H recruitment, though non RNase H mechanisms have also been reported. These studies highlight a broad potential of LNA oligonucleotides for effective gene silencing both *in vitro* and *in vivo*. Recently, Jacobsen et al. for example showed that LNA AONs are effective inhibitors of HIV-1 expression.¹⁵

DNAzymes are catalytically active DNA molecules that are able to cleave RNA in a sequence-specific manner after binding to complementary sequences. Studies conducted using LNA-modified DNAzymes, termed LNAzymes showed an enhanced efficiency of

*Correspondence to: Jesper Wengel; Nucleic Acid Center; Department of Physics and Chemistry; University of Southern Denmark; Campusvej 55, Odense M 5230 Denmark; Email: jwe@ifk.sdu.dk

Submitted: 01/20/09; Revised: 04/15/09; Accepted: 04/18/09

Previously published online as an RNA Biology E-publication:
<http://www.landesbioscience.com/journals/rnabiology/article/8807>




Figure 2. Schematic representations of siRNA and sisiRNA constructs. In a classical siRNA construct, the sense strand complementary to the antisense strand is single stranded. In a sisiRNA construct, a nick has been introduced in the sense strand which is therefore constituted by two separate short strands complementary to the antisense strand. Both constructs are depicted with two nucleotide 3'-overhangs and with a shortened duplex region; this in typical siRNA and sisiRNA constructs contains 19 base pairs.

RNA cleavage.¹⁶ In line with previous findings, Jacobsen et al. reported efficient inhibition of HIV-1 expression by targeting LNAzymes to functionally selected binding sites,¹⁵ whereas targeting of miRNAs by using LNAzymes has recently been reported by Maiti and co-workers.¹⁷

RNA interference (RNAi) has been developed as a highly potent approach to knock down gene expression.¹⁸ RNAi mediated by small interfering RNAs (siRNAs) can target complementary mRNA and induce its degradation upon incorporation of the siRNA into the RNA-induced silencing complex (RISC). siRNAs themselves are candidates for incorporation of modified nucleotides for improved bio-stability and effective RNA targeting. In this direction, the use of LNA-modified siRNA, termed siLNA, has been investigated. Recently, Mook et al. evaluated the effect of LNA-modified siRNA both in vitro and in vivo.¹⁹ They showed that minimal LNA-modifications at the 3'-end of siRNAs are effective to stabilize the siRNA, and that multiple LNA-modifications may lead to decreased efficacy in vitro and in vivo. The study also highlighted reduced off-target gene regulation when using LNA-modified siRNA (compared to the unmodified siRNA). Very recently, Bramsen et al. introduced a three stranded siRNA construct termed 'small internally segmented interfering RNA' (sisiRNA, Fig. 2) in which the antisense strand is complexed with two short sense strands of approximately 10–12 nt in length.²⁰ In the sisiRNA approach only the antisense strand is functional as the nick completely eliminates unintended mRNA targeting by the sense strand. LNA nucleotides were incorporated to stabilize the sisiRNA constructs which proved efficient for gene silencing upon transfection into a H1299 lung carcinoma cell line.

MicroRNAs (miRNAs) constitute a class of short regulatory RNAs (~22 n) that control gene expression post-transcriptionally during development, differentiation and metabolism.^{21,22} Similar to classical AONs developed for the inhibition of coding RNAs, synthetic oligonucleotides are the only rational approach for specific inhibition of individual miRNAs and therefore have the potential to be developed into an important new class of miRNA targeting drugs. Elmén et al. have recently described effective miRNA silencing in non-human primates by short LNA-modified oligonucleotides (LNA anti-miRs; LNA/DNA mixmers) designed to target liver-expressed miR-122.²³

LNA Aptamers

Aptamers²⁴⁻²⁶ are short DNA or RNA oligonucleotide sequences which can bind tightly to a specific molecular target because of their ability to form three dimensional structures. The remarkable properties and applications of LNA oligonucleotides highlighted above substantiate the desire to explore aptamers composed of LNA-modified nucleotides to rival aptamers composed of unmodified RNA or DNA. Darfeuille et al. introduced LNA modification of an already selected RNA aptamer targeted to the TAR RNA element of HIV-1.^{27,28} Surface plasmon resonance (SPR) experiments identified LNA/DNA mixmers binding to TAR RNA with a dissociation constant in the low nanomolar range. Schmidt et al. later described the capability of LNA nucleotides to improve the in vivo stability of aptamers.²⁹ Further work showed that an aptamer modified with LNA nucleotides targeting the TAR RNA element of HIV-1 displayed good binding properties and competed with the viral protein Tat for binding to TAR.³⁰ Furthermore, the same group later reported a TAR RNA aptamer as an LNA/2'-O-methyl RNA mixmer which displayed improved HIV-1 TAR element binding.³¹ All of these results underscore the desire to develop procedures to allow evolution of LNA aptamers by SELEX-based^{32,33} processes.

Future Prospects

The remarkable properties of LNA with respect to high affinity and specificity make this analogue unique for applications in molecular biology research, biotechnology and RNA targeting. We believe that LNA constructs will be important molecules as the prospects of nucleic acid based drugs will be realized. This is underlined by the recent reports on efficient miRNA targeting using LNA probes and the fact that LNA/DNA mixmers have been forwarded into human clinical trials.³⁴ We are attempting to develop ways of evolving LNA-based aptamers and have recently published reports on enzymatic methods for synthesis of LNA containing oligonucleotides.^{35,36}

Acknowledgements

The Nucleic Acid Center is a research center of excellence funded by the Danish National Research Foundation for studies on nucleic acid chemical biology. We thank the Danish National Research Foundation for financial support.

References

- Vitravene study group. Randomized dose-comparison studies of intravitreal foscarnet for treatment of cytomegalovirus retinitis that has reactivated or is persistently active despite other therapies in patients with AIDS. *Am J Ophthalmol* 2002; 133:475-83.
- Singh SK, Nielsen P, Koshkin AA, Wengel J. LNA (locked nucleic acids): synthesis and high-affinity nucleic acid recognition. *Chem Commun* 1998; 455-6.
- Koshkin AA, Nielsen P, Meldgaard M, Rajwanshi VK, Singh SK, Wengel J. LNA (Locked Nucleic Acid): An RNA mimic forming exceedingly stable LNA:LNA duplexes. *J Am Chem Soc* 1998; 120:13252-3.
- Obika S, Nanbu D, Hari Y, Andoh J, Morio K, Doi T, et al. Stability and structural features of the duplexes containing nucleoside analogues with a fixed N-type conformation, 2'-O,4'-C-methylene ribonucleosides. *Tetrahedron Lett* 1998; 39:5401-4.
- Koshkin AA, Singh SK, Nielsen P, Rajwanshi VK, Kumar R, Meldgaard M, et al. LNA (Locked Nucleic Acids): Synthesis of the adenine, cytosine, guanine, 5-methylcytosine, thymine and uracil bicyclonucleoside monomers, oligomerisation and unprecedented nucleic acid recognition. *Tetrahedron* 1998; 54:3607-30.
- Wengel J. Synthesis of 3'-C- and 4'-C-branched oligodeoxynucleotides and the development of Locked Nucleic Acid (LNA). *Acc Chem Res* 1999; 32:301-10.
- Petersen M, Wengel J. LNA: a versatile tool for therapeutics and genomics. *Trends Biotechnol* 2003; 21:74-81.
- Vester B, Wengel J. LNA (Locked Nucleic Acid): high-affinity targeting of complementary RNA and DNA. *Biochemistry* 2004; 43:13233-41.
- Jespen JS, Sørensen MD, Wengel J. Locked Nucleic Acid: a potent nucleic acid analog in therapeutics and biotechnology. *Oligonucleotides* 2004; 14:130-46.
- Obika S, Nanbu D, Hari Y, Morio K, In Y, Ishida T, et al. Synthesis of 2'-O,4'-C-methyleneuridine and -cytidine. Novel bicyclic nucleosides having a fixed C3'-endo sugar pucker. *Tetrahedron Lett* 1997; 38:8735-8.
- Petersen M, Bondensgaard K, Wengel J, Jacobsen JP. Locked Nucleic Acid (LNA) recognition of RNA: NMR solution structures of LNA:RNA hybrids. *J Am Chem Soc* 2002; 124:5974-82.
- Nielsen KE, Rasmussen J, Kumar R, Wengel J, Jacobsen JP, Petersen M. NMR Studies of Fully Modified Locked Nucleic Acid (LNA) Hybrids: Solution structure of an LNA:RNA hybrid and characterization of an LNA:DNA hybrid. *Bioconjugate Chem* 2004; 15:449-57.
- Kauppinen S, Vester B, Wengel J. Locked Nucleic Acid (LNA): High-affinity targeting of RNA for diagnostics and therapeutics. *Drug Discovery Today: Technologies* 2005; 2:287-90.
- Ørum H, Wengel J. Locked nucleic acids (LNA): A promising molecular family for gene-function analysis and antisense drug development. *Curr Opin Mol Ther* 2001; 3:239-43.
- Jakobsen MR, Haasnoot J, Wengel J, Berkhout B, Kijms J. Efficient inhibition of HIV-1 expression by LNA modified antisense oligonucleotides and DNazymes targeted to functionally selected binding sites. *Retrovirology* 2007; 26; 4:29.
- Vester B, Lundberg LB, Sørensen MD, Babu BR, Douthwaite S, Wengel J. LNAzymes: Incorporation of LNA-type monomers into DNazymes markedly increases RNA cleavage. *J Am Chem Soc* 2002; 124:13682-3.
- Jadhav VM, Scaria V, Maiti S. Antagomirzymes: Oligonucleotide enzymes that specifically silence MicroRNA function. *Angew Chem Int Ed* 2009; 48:2557-60.
- Dorsett Y, Tuschl T. siRNAs: Applications in functional genomics and potential as therapeutics. *Nature Rev* 2004; 3:318-29.
- Mook OR, Baas F, de Wissel MB, Fluitier K. Evaluation of locked nucleic acid-modified small interfering RNA in vitro and in vivo. *Mol Cancer Ther* 2007; 6:833-43.
- Bramsen JB, Laursen MB, Damgaard CK, Lena SW, Babu BR, Wengel J, et al. Improved silencing properties using small internally segmented interfering RNAs. *Nucleic Acids Res* 2007; 35:5886-97.
- Ambros V. MicroRNAs: Tiny regulators with great potential. *Cell* 2001; 107:823-6.
- Bartel DP. MicroRNAs: Genomics, biogenesis, mechanism and function. *Cell* 2004; 116:281-97.
- Elmén J, Lindow M, Schütz S, Lawrence M, Petri A, Obad S, et al. LNA-mediated microRNA silencing in non-human primates. *Nature* 2008; 452:896-9.
- Jayasena SD. Aptamers: An emerging class of molecules that rival antibodies in diagnostics. *Clinical Chem* 1999; 45:1628-50.
- Famulok M, Mayer G, Blind M. Nucleic acid aptamers—from selection in vitro to applications in vivo. *Acc Chem Res* 2000; 33:591-9.
- Nimjee SM, Christopher PR, Sullenger BA. Aptamers: An emerging class of therapeutics. *Annu Rev Med* 2005; 56:555-83.
- Darfeuille F, Hansen JB, Ørum H, Di Primo C, Tolumé JJ. LNA/DNA chimeric oligomers mimic RNA aptamers targeted to the TAR RNA element of HIV-1. *Nucleic Acids Res* 2004; 32:3101-7.
- Darfeuille F, Hansen JB, Ørum H, Di Primo C, Tolumé JJ. Aptamers targeted to an RNA hairpin show improved specificity compared to that of complementary oligonucleotides. *Biochemistry* 2006; 45:12076-82.
- Schmidt KS, Borkowski S, Kurreck J, Stephens AW, Bald R, Hecht M, et al. Application of locked nucleic acids to improve aptamer in vivo stability and targeting function. *Nucleic Acids Res* 2004; 32:5757-65.
- Lebars I, Richard T, Di Primo C, Tolumé JJ. LNA derivatives of a kissing aptamer targeted to the trans-activating responsive RNA element of HIV-1. *Blood Cells Mol Dis* 2007; 38:204-9.
- Di Primo C, Rudloff I, Reigadas S, Arzumanov A, Gait MJ, Tolumé JJ. Systematic screening of LNA/2'-O-methyl chimeric derivatives of a TAR RNA aptamer. *FEBS Lett* 2007; 771-4.
- Ellington AD, Szostak JW. In vitro selection of RNA molecules that bind specific ligands. *Nature* 1990; 346:818-22.
- Turek C, Gold L. Systematic evolution of ligands by exponential enrichment: RNA ligands to bacteriophage T4 DNA polymerase. *Science* 1990; 249:505-10.
- <http://www.santaris.com>
- Veedu RN, Vester B, Wengel J. Polymerase chain reaction and transcription using locked nucleic acid nucleotide triphosphates. *J Am Chem Soc* 2008; 130:8124-5.
- Veedu RN, Vester B, Wengel J. Efficient enzymatic synthesis of LNA-modified DNA duplexes using KOD DNA polymerase. *Org Biomol Chem* 2009; 7:1404-9.