



**Murdoch**  
UNIVERSITY

**MURDOCH RESEARCH REPOSITORY**

<http://researchrepository.murdoch.edu.au>

*This is the author's final version of the work, as accepted for publication following peer review but without the publisher's layout or pagination.*

**Hunter, G.C., van der Merwe, N.A., Burgess, T.I., Carnegie, A.J., Wingfield, B.D., Crous, P.W. and Wingfield, M.J. (2008) Global movement and population biology of *Mycosphaerella nubilosa* infecting leaves of cold-tolerant *Eucalyptus globulus* and *E. nitens*. *Plant Pathology*, 57 (2). pp. 235-242.**

<http://researchrepository.murdoch.edu.au/2956>

Copyright © 2007 The Authors  
It is posted here for your personal use. No further distribution is permitted.

# Global movement and population biology of *Mycosphaerella nubilosa* infecting leaves of cold-tolerant *Eucalyptus globulus* and *E. nitens*

G. C. Hunter<sup>a\*†</sup>, N. A. van der Merwe<sup>b</sup>, T. I. Burgess<sup>c</sup>, A. J. Carnegie<sup>d</sup>, B. D. Wingfield<sup>b</sup>, P. W. Crous<sup>e</sup> and M. J. Wingfield<sup>a</sup>

<sup>a</sup>Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria 0002; <sup>b</sup>Department of Genetics, University of Pretoria, Pretoria 0002, South Africa; <sup>c</sup>School of Biological Sciences and Biotechnology, Murdoch University, Perth, WA 6150; <sup>d</sup>Forest Resources Research, NSW Department of Primary Industries, PO Box 100, Beecroft 2119, NSW, Australia; and <sup>e</sup>Centraalbureau voor Schimmelcultures, Fungal Biodiversity Centre, PO Box 85167, 3508 AD, Utrecht, the Netherlands

Using 10 polymorphic DNA-based microsatellite markers, the genetic diversity of eight *Mycosphaerella nubilosa* populations from *Eucalyptus*, comprising 497 isolates from five different countries, was studied using a hierarchical sampling regime. *Mycosphaerella nubilosa* from eastern Australia (New South Wales) had higher gene (0.506) and genotypic (76%) diversity than other populations, supporting the view that this represents the origin of the pathogen. It was also evident that *M. nubilosa* populations from Europe and Tanzania were clonal, with the same multilocus haplotypes occurring in South Africa, but being absent in Australia. This suggests that *M. nubilosa* may have been introduced into Europe via Africa, with a pathway of gene flow from Australia to South Africa, further into Africa and finally to Europe.

**Keywords** : cold-tolerant *Eucalyptus* spp., gene diversity, gene flow, genotypic diversity, microsatellite markers, mycosphaerella leaf blotch

## Introduction

The ascomycete genus *Mycosphaerella* includes more than 80 species of fungi that have been associated with leaf diseases of *Eucalyptus* spp. (Crous, 1998; Crous *et al.*, 2004, 2006). Infection by *Mycosphaerella* spp. can affect photosynthesis in young *Eucalyptus* leaves, resulting in premature defoliation and leading to reduced growth of trees (Park & Keane, 1982b; Carnegie & Ades, 2003; Pinkard & Mohammed, 2006).

*Mycosphaerella nubilosa* is one of the most important species on *Eucalyptus*, causing mycosphaerella leaf blotch (MLB), particularly on cold-tolerant species such as *E. globulus* and *E. nitens* (Park & Keane, 1987; Carnegie *et al.*, 1998; Carnegie & Ades, 2002; Hunter *et al.*, 2004a). This fungus typically infects juvenile leaves and susceptible trees can be severely defoliated (Park & Keane, 1982b; Lundquist & Purnell, 1987).

\*E-mail: g.hunter@cbs.knaw.nl

†Present address: Centraalbureau voor Schimmelcultures (CBS), Fungal Biodiversity Centre, PO Box 85167, 3508 AD, Utrecht, the Netherlands.

*Mycosphaerella nubilosa* was first described from *Eucalyptus* leaves in Victoria, Australia (Cooke, 1891) and subsequently in Tasmania, New South Wales, South and Western Australia (Hansford, 1956; Park & Keane, 1982a,b; Park, 1988; Maxwell *et al.*, 2001; Milgate *et al.*, 2001; Carnegie, 2007a,b). Elsewhere *M. nubilosa* occurs in the North Island of New Zealand (Dick, 1982) and in Africa, in Ethiopia, Kenya, South Africa, Tanzania and Zambia (Crous *et al.*, 2004, 2006; Hunter *et al.*, 2004a; Alemu *et al.*, 2006), and was accidentally introduced into commercial plantations of *E. globulus*, where it caused serious damage in Spain and Portugal (Crous *et al.*, 2004).

Apart from *E. globulus* and *E. nitens*, *M. nubilosa* infects *E. bridgesiana*, *E. cypellocarpa*, *E. dunnii*, *E. grandis*, *E. quadrangulata*, *E. globulus* ssp. *Bicostata* and ssp. *Maidenii* (Dick, 1982; Crous, 1998; Crous *et al.*, 2004; Hunter *et al.*, 2004a,b; Milgate *et al.*, 2005; Carnegie, 2007b). Because of their rapid growth, frost tolerance and favourable wood qualities, *E. nitens* and *E. globulus* are amongst the most popular *Eucalyptus* spp. Propagated for commercial forestry in the above countries and are particularly susceptible to *M. nubilosa*; MLB can be a serious impediment to the sustainability of *E. nitens* and *E. globulus* plantations.

*Mycosphaerella nubilosa* naturally occurs only in its sexual state, producing many small black pseudorhelia that may be amphigenous, but are predominantly hypophyllous on lesions (Crous, 1998; Crous *et al.*, 2004). When leaves are wet, ascospores are released from pseudorhelia and act as wind-dispersed primary inoculum and infect via stomata (Park & Keane, 1982b; Park, 1988). Although the anamorph state *Uwebraunia juvenis* was linked to *M. nubilosa* (Crous *et al.*, 2004), this connection has not been reconfirmed and remains doubtful.

Most research conducted on *M. nubilosa* has focussed on its epidemiology, taxonomy, host association and phylogeny, but little is known of its population biology. The aim of this study was, therefore, to investigate the population biology of *M. nubilosa* from several countries by determining the number of alleles at 10 polymorphic microsatellite loci. There were three primary objectives: (i) consideration of the genetic diversity of *M. nubilosa*, (ii) examination of this variation between different countries, and (iii) determination of the likely origin and global movement of *M. nubilosa*.

## Materials and Methods

### Isolates

Diseased *Eucalyptus* leaves infected with *M. nubilosa* were collected from Australia, Portugal, South Africa, Spain and Tanzania (Table 1).

Two populations of *M. nubilosa* were collected in Australia. From eastern Australia, one leaf was randomly collected from each of 40 *E. globulus* trees in one plantation near Bonalbo, north-eastern New South Wales (EA-MT). Similarly, in Western Australia one leaf was

randomly collected from each of 32 *E. globulus* trees in the south of the state (WA-MT). One plantation of *E. globulus* was sampled in northern Spain and one leaf was selected from each of 55 trees (S-MT). Likewise, one plantation of *E. globulus* was sampled in central Portugal, where 42 leaves were taken from an equal number of *E. globulus* trees (P-MT).

In Njombe, Tanzania, 56 *E. globulus* trees were randomly sampled, and one Leaf was collected from each tree (T-MT). In South Africa, three plantations of *E. nitens* were sampled in the Mpumalanga Province, namely Issabelladale, Wynton and Rooihogte. From Rooihogte, 48 leaves displaying spots were randomly collected from a single tree (R-1T). At this plantation, a single lesion from a single leaf was sampled and 32 isolates were collected from this lesion (R-1L). In addition, a single leaf was randomly collected from each of 56 trees within Rooihogte (R-MT). From Wynton, 64 leaves exhibiting MLB symptoms were collected from a single tree (W-1T). Furthermore, 64 trees in this plantation were randomly selected and one diseased leaf was taken from each (W-MT). From Issabelladale, 40 leaves showing MLB lesions were randomly collected from one tree (I-1T). In addition, 60 trees within the same plantation were randomly selected and one leaf was collected from each of these trees (I-MT).

#### Isolation, *in vitro* culture and DNA extraction

Isolation of *M. nubilosa* from diseased *Eucalyptus* leaves followed the protocol of Crous (1998). One ascospore per lesion, exhibiting the type-F ascospore germination pattern (Crous, 1998; Crous *et al.*, 2004), was cut from the agar surface and transferred to fresh 2% malt extract agar and incubated at 25°C for approximately 2-3 months to ensure sufficient mycelial growth. All isolates of *M. nubilosa* used in this study were deposited in the culture collection of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, South Africa.

Subsequently, mycelium from actively growing single-spore cultures of *M. nubilosa* was scraped from the surface of cultures, freeze-dried for 24 h and ground to a fine powder using

Table 1 Origins of *Mycosphaerella nubilosa* isolates used for this study

Country	Area	Plantation	Hierarchical Level	PC <sup>a</sup>	Host	Collector	Date	No. isolates <sup>b</sup>	
South Africa	Mpumalanga	Rooihogte	Single lesion	R-1L	<i>E. nitens</i>	G. C. Hunter	2005	32	
			Single tree	R-1T	<i>E. nitens</i>	G. C. Hunter	2005	48	
			Plantation	R-MT	<i>E. nitens</i>	G. C. Hunter	2005	56	
	Issabelladale	Single tree	I-1T	<i>E. nitens</i>	G. C. Hunter	2005	30		
			Plantation	I-MT	<i>E. nitens</i>	G. C. Hunter	2005	29	
			Wynton	Single tree	W-1T	<i>E. nitens</i>	G. C. Hunter	2005	49
	Plantation	W-MT		<i>E. nitens</i>	G. C. Hunter	2005	49		
	Australia	Western Australia	N/A	Plantation	WA-MT	<i>E. globulus</i>	A. Maxwell	2002	32
		New South Wales	Bonalbo	Plantation	EA-MT	<i>E. globulus</i>	A. J. Carnegie	2004	19
Spain			Plantation	S-MT	<i>E. globulus</i>	J. P. M. Vasquez	2002	55	
Portugal			Plantation	P-MT	<i>E. globulus</i>	M. J. Wingfield	2005	42	
Tanzania			Plantation	T-MT	<i>E. globulus</i>	J. Roux	2004	56	
						Total		497	

<sup>a</sup>Population code assigned to each population and subpopulation.

<sup>b</sup>Number of isolates per hierarchical sampling level.

liquid nitrogen. DNA was isolated using the phenol:chloroform (1:1) extraction protocol described by Hunter *et al.* (2004a,b). Isolated DNA was separated by electrophoresis on 1 % agarose gels (w/v) (Roche Diagnostics), stained with ethidium bromide and visualized under ultraviolet light.

#### PCR amplification and allele size determination

DNA from isolates of *M. nubilosa* served as template DNA for polymerase chain reactions (PCR). Fluorescently labelled primers used for PCR reactions were the 10 polymorphic microsatellite primers developed by Hunter *et al.* (2006). The PCR reaction mixture and conditions were the same as those described previously (Hunter *et al.*, 2006). Amplified DNA was visualized on 2% agarose gels stained with ethidium bromide and viewed under ultraviolet light. Sizes of PCR amplicons were estimated by comparison against a 100-bp molecular weight marker (O'RangeRuler™ 100-bp DNA ladder (Fermentas Life Sciences). PCR products were purified using Sephadex G-50 (Sigma Aldrich) in Centri-sep Spin Columns (Princeton Separations) as outlined by the manufacturer.

Allele sizes of amplified PCR products were determined following the method of Hunter *et al.* (2006). The number of alleles per locus and the total number of multilocus haplotypes (MLHs, hereafter referred to as haplotypes) across all the *M. nubilosa* populations were determined.

#### Gene and genetic diversity

The frequency of alleles at each locus was calculated for every *M. nubilosa* population. Allele diversity was also determined using the software program POPGINE (Yeh *et al.*, 1999) and the gene diversity ( $H$ ) of Nei (1973) was calculated at each locus and also averaged over all loci:

$$H = 1 - \sum x_k^2$$

Chi-squared tests for differences in allele frequencies between the *M. nubilosa* populations were also calculated from clone-corrected datasets (Workman & Niswander, 1970).

Every *M. nubilosa* isolate was assigned a haplotype code according to the observed allele sizes, across the 10 microsatellite loci. Genotypic diversity ( $G$ ) was calculated using the equation of Stoddart & Taylor (1988):

$$G = 1 / \sum_{x=0}^N [f_x(X/N)^2]$$

To compare genotypic diversities between the various *M. nubilosa* populations, the maximum percentage of genotypic diversity ( $\hat{G}$ ) was calculated:

$$\hat{G} = G/N \times 100$$

(Chen *et al.*, 1994). Genotypic diversity values were also compared between *M. nubilosa* populations at the plantation level for significant differences using a *t*-test (Chen *et al.*, 1994).

#### Population differentiation and gene flow

Clone-corrected populations of *M. nubilosa* were used to calculate population differentiation ( $\theta$ ) between the South African and Australian *M. nubilosa* populations using the computer software program MULTILOCUS (Agapow & Burt, 2001), using 1000 randomized datasets. The population differentiation value ( $\theta$ ), Weir's formulation of Wright's  $F_{ST}$ , was calculated (Weir, 1996; Agapow & Burt, 2001). Furthermore, gene flow ( $M$ ) was calculated from theta ( $\theta$ ) using the equation of Cockerham & Weir (1993):

$$M = [(1/\theta) - 1]/2.$$

## Results

### Isolates

A total of 497 isolates of *M. nubilosa* were collected from diseased *Eucalyptus* leaves (Table 1). These represented a wide range of geographic locations and also different hierarchical levels, and included 349 isolates from Africa, 97 from Portugal and Spain, and 51 from Australia.

### PCR amplification and allele size determination

A total of 66 alleles were observed across all 10 loci for all the *M. nubilosa* populations, with an average of 6.6 alleles per locus. The Australian *M. nubilosa* populations WA-MT and EA-MT exhibited the greatest number of alleles across all loci, namely 22 and 44, respectively (Table 2).

South African populations exhibited much lower numbers of alleles across the 10 loci than those from Australia. In total, 25 alleles were observed across all of the South African *M. nubilosa* populations at the various hierarchical levels and the majority of these alleles were commonly shared. South African populations R-MT and I-1T exhibited the most alleles (a total of 18) (Table 2). One population (R- 1L) collected from a single lesion on an *E. nitens* leaf from Rooihogte yielded a total of 13 different alleles for the 10 loci across 32 *M. nubilosa* isolates (Table 2). For the isolates collected in Spain, Portugal and Tanzania all loci were monomorphic.

Across all *M. nubilosa* populations, all of the 10 microsatellite loci were polymorphic. Both the South African and Western Australian populations exhibited 60% polymorphic loci, whilst 80% of loci were polymorphic for the population EA-MT in New South Wales.

**Table 2** Number of alleles and gene diversity at 10 microsatellite loci (MN-1–MN-14) determined using the program POPGENE for populations of *Mycosphaerella nubilosa* collected in South Africa (Rooihoogte, Issabelladale and Wynton) and Australia (Western Australia and New South Wales)

	South Africa						Australia		
	Rooihoogte			Issabelladale		Wynton		WA	NSW
	R-1L	R-1T	R-MT	I-1T	I-MT	W-1T	W-MT	WA-MT	EA-MT
N <sup>a</sup>	32	48	56	30	29	49	49	32	19
N <sup>b</sup>	4	24	24	16	17	22	23	10	16
Alleles	13	17	18	18	16	16	17	22	44
P	3	6	5	5	5	5	6	6	8
P (%)	30	60	50	50	50	50	60	60	80
H	0.149	0.240	0.248	0.222	0.211	0.231	0.250	0.242	0.506
H*	0.125	0.258	0.255	0.248	0.242	0.247	0.255	0.246	0.506

N<sup>a</sup> = number of isolates used in uncorrected dataset; N<sup>b</sup> = number of isolates used in clone-corrected dataset; P = number of polymorphic loci; P (%) = percentage of polymorphic loci; H = gene diversity of population (Nei, 1973); H\* = gene diversity of clone-corrected population.

**Table 3** Number of multilocus haplotypes (MLHs), genotypic diversity and maximum percentage of genotypic diversity of *Mycosphaerella nubilosa* isolates compiled from 10 polymorphic microsatellite markers. Haplotypes from Spain (S-MT), Portugal (P-MT) and Tanzania (T-MT) all at the plantation level. Those from three plantations in South Africa at different hierarchical levels: Rooihoogte plantation at the plantation level (R-MT), within one tree (R-1T) and within one lesion (R-1L); Issabelladale plantation at the plantation level (I-MT) and within one tree (I-1T); and Wynton plantation at the plantation level (W-MT) and within one tree (W-1T). Also two *M. nubilosa* populations, both at the plantation level, from Australia: Western Australia (WA-MT) and New South Wales (EA-MT)

	South Africa						Australia					
	Spain	Portugal	Tanzania	Rooihoogte			Issabelladale		Wynton	WA	NSW	
	S-MT	P-MT	T-MT	R-1L	R-1T	R-MT	I-1T	I-MT	W-1T	W-MT	WA-MT	EA-MT
N <sup>a</sup>	55	42	56	32	48	56	30	29	49	49	32	19
N (g) <sup>b</sup>	1	1	1	4	24	24	16	17	22	23	10	16
G <sup>c</sup>	N/A	N/A	N/A	2.26	15.78	16.88	6.63	8.73	9.61	14.82	4.86	14.49
G <sup>d</sup>	N/A	N/A	N/A	7.1	32.9	30.14	22.1	30.1	19.6	30.2	15.2	76.3

<sup>a</sup>Number of *M. nubilosa* isolates.

<sup>b</sup>Number of MLHs.

<sup>c</sup>Genotypic diversity (Stoddart & Taylor, 1988).

<sup>d</sup>Maximum percentage of genotypic diversity.

## Gene and genotypic diversity

The South African *M. nubilosa* populations had a moderate to high gene diversity, ranging from 0.149 to 0.250 for populations R-1L and W-MT, respectively (Table 2).

Gene diversity for the Western Australia population (WA-MT) (0.242) was comparable with that of the South African populations. In contrast, a substantially higher gene diversity value of 0.506 was observed for the New South Wales population (EA-MT).

A total of 68 haplotypes were observed across all *M. nubilosa* populations from the various locations considered (Table 3). Almost half (49%) of the haplotypes were unique and detected only once. The South African populations had a total of 35 haplotypes and the majority of these haplotypes were distributed among the various South African populations. However, seven haplotypes were unique to certain South African populations. None of the haplotypes that were observed for the South African populations occurred in either of the Australian populations. However, the Spanish and Portuguese populations both had only one haplotype, which was also observed in the R-MT, W-1T and W-MT *M. nubilosa* populations

of South Africa. The Tanzanian population also had only one haplotype, which was also observed in six of the seven South African populations.

A total of 26 haplotypes were observed across the two Australian populations WA-MT and EA-MT, the former with 10 unique haplotypes, the latter with 16.

Genotypic diversity (G) varied between the various populations of *M. nubilosa* from different locations (Table 3). The lowest maximum percentage of genotypic diversity ( $\hat{G}$ ), 7.1% was observed for the South African *M. nubilosa* isolates obtained from a single lesion (R-1L). However, maximum percentages of genotypic diversity for South African *M. nubilosa* populations at the plantation level were similar to each other with values of 30.14, 30.1 and 30.2% for R-MT, I-MT and W-MT, respectively (Table 3). Interestingly, the Western Australian population (WA-MT) had the second lowest value of genotypic diversity at 15.2% of the theoretical maximum value, which was lower than for all the South African populations except population R-1L. The highest genotypic diversity (76% of the theoretical maximum) was observed in the New South Wales population (EA-MT). No significant

**Table 4** Gene diversity ( $H$ ) and contingency  $\chi^2$  tests for differences in allele frequencies for the 10 microsatellite loci across the clone-corrected South African populations of *Mycosphaerella nubilosa* from different hierarchical levels. (A) Within one lesion (R-1L), within one tree (R-1T) and within one plantation (R-MT) of the Rooihooft plantation. (B) Within one tree (I-1T) and within one plantation (I-MT) of the Issabelladale plantation. (C) Within one tree (W-1T) and within one plantation (W-MT) of the Wynton plantation. (D) Comparison between three South African plantations at the plantation level (R-MT, I-MT and W-MT). Determined  $\chi^2$  values not significant

	Gene diversity ( $H$ )							A		B		C		D	
	R-1L	R-1T	R-MT	I-1T	I-MT	W-1T	W-MT	$\chi^2$	d.f.	$\chi^2$	d.f.	$\chi^2$	d.f.	$\chi^2$	d.f.
MN-3	0.00	0.49	0.49	0.43	0.46	0.48	0.50	3.0	2	0.06	1	0.22	1	2.2	2
MN-4	0.38	0.50	0.49	0.55	0.50	0.48	0.50	0.9	2	1.22	2	0.22	1	0.7	2
MN-8	0.38	0.50	0.57	0.54	0.48	0.52	0.48	3.3	6	1.10	2	1.07	2	3.6	6
MN-9	0.00	0.47	0.52	0.46	0.46	0.48	0.54	3.6	4	1.37	2	1.07	2	3.9	6
MN-10	0.00	0.08	0.00	0.00	0.00	0.00	0.08	1.2	2	–	–	0.98	1	1.8	2
MN-11	0.50	0.53	0.49	0.49	0.53	0.50	0.45	2.2	4	1.10	2	1.07	1	5.8	4
N <sup>a</sup>	4	24	24	16	17	22	23								
Mean	0.13	0.23	0.25	0.25	0.24	0.25	0.25								

<sup>a</sup>Number of *M. nubilosa* isolates from the clone-corrected dataset.

**Table 5** Gene diversity ( $H$ ) and contingency  $\chi^2$  tests for differences in allele frequencies for the 10 polymorphic microsatellite loci across the clone corrected *Mycosphaerella nubilosa* populations from (A) South Africa [RSA = (R-MT + I-MT + W-MT)] and Western Australia (WA-MT), (B) South Africa [RSA = (R-MT + I-MT + W-MT)] and eastern Australia (EA-MT), and (C) Western Australia (WA-MT) and New South Wales (EA-MT)

Locus	Gene diversity ( $H$ )			A		B		C	
	RSA	WA-MT	EA-MT	$\chi^2$	d.f.	$\chi^2$	d.f.	$\chi^2$	d.f.
MN-1	0.00	0.00	0.00	–	–	51.0*	1	26.0*	1
MN-2	0.00	0.18	0.88	3.6	1	46.5*	9	22.2*	10
MN-3	0.50	0.00	0.30	7.8*	1	15.9*	2	2.1	1
MN-4	0.50	0.00	0.69	9.4*	1	38.2*	4	22.2*	4
MN-7	0.00	0.32	0.43	7.3*	1	51.0*	2	26.0*	3
MN-8	0.54	0.84	0.63	39.6*	9	22.6*	8	20.4*	10
MN-9	0.54	0.32	0.76	5.1	3	46.4*	9	26.0*	9
MN-10	0.06	0.00	0.63	0.3	1	42.2*	5	19.0*	4
MN-11	0.52	0.62	0.00	19.0*	3	15.0*	2	9.9*	2
MN-14	0.00	0.18	0.73	3.6	1	51.0*	4	22.4*	4
N <sup>a</sup>	35	10	16						
Mean	0.27	0.25	0.51						

\*Indicates significant  $\chi^2$  values ( $P < 0.05$ ).

<sup>a</sup>Number of *M. nubilosa* isolates from the clone-corrected dataset.



Differences ( $P < 0.05$ ) in genotypic diversities were observed between *M. nubilosa* populations at the plantation level (R-MT, I-MT, W-MT, WA-MT and EA-MT) using a *t*-test (data not shown).

#### Population differentiation and gene flow

Based on contingency chi-squared tests, there were no significant differences in allele frequencies at any of the loci when the *M. nubilosa* subpopulations from the Rooihogte (R-1L, R-1T and R-MT), Issabelladale (I-1T and I-MT) and Wynton (W-1T and W-MT) *E. nitens* plantations in South Africa were considered (Table 4). Three subpopulations at the plantation level from South Africa (R-MT, I-MT and W-MT) were also compared with each other and no significant differences were observed between the allele frequencies at any of the 10 loci (Table 4). These three South African populations were, therefore, combined at the plantation level and this dataset was clone-corrected, in order to compare it to the two Australian populations (WA-MT and EA-MT).

Based on contingency chi-squared tests, significant differences ( $P < 0.05$ ) in allele frequencies were observed at nine of the 10 polymorphic loci when the two Australian *M. nubilosa* populations (WA-MT and F.A-MT) were compared against each other (Table 5). Significant differences ( $P < 0.05$ ) in allele frequencies were also observed at five loci when the combined South African *M. nubilosa* population (RSA) was compared with the Western Australian population (WA-MT), and for all 10 loci when population RSA was compared with the population from New South Wales (Table 5).

**Table 6** Population differentiation values ( $\theta$ ), for populations of *Mycosphaerella nubilosa* above the diagonal, calculated after 1000 randomizations using the program MULTILOCUS. Gene flow (number of migrants) (*M*) indicated below the diagonal

	R-MT	I-MT	W-MT	WA-MT	EA-MT
R-MT		-0.0054	-0.03034	0.25*	0.55*
I-MT	9.63		-0.0061	0.31*	0.54*
W-MT	16.93	82.47		0.29*	0.55*
WA-MT	1.54	1.13	1.24		0.49*
EA-MT	0.41	0.43	0.41	0.52	

\*Indicates significant values ( $P < 0.001$ ).

R-MT, Rooihogte; I-MT, Issabelladale; W-MT, Wynton, South Africa.

WA-MT, Western Australia; EA-MT, Eastern Australia.

Population differentiation values ( $\theta$ ) were not significantly different between any of the South African *M. nubilosa* populations at the plantation level (R-MT, I-MT and W-MT) (Table 6). Gene flow (*M*) between the three South African populations was high, with the greatest number of migrants ( $M = 82.47$ ) occurring between the W-MT and I-MT populations and the least ( $M = 9.63$ ) occurring between the I-MT and R-MT populations (Table 6). There were,

however, significant differences ( $P < 0.001$ ) at the plantation level between the three South African populations (R-MT, I-MT and W-MT) and the two Australian populations (WA-MT and EA-MT) based on  $\theta$  values (Table 6). Similarly,  $\theta$  values indicated a significant population differentiation ( $P < 0.001$ ) between the two Australian populations (Table 6), with very little gene flow ( $M = 0.52$ ) observed between the Western Australian and New South Wales populations. However, there was low gene flow ( $M = 1.13-1.54$ ) between the Western Australian population (WA-MT) and the three South African *M. nubilosa* populations (R-MT, I-MT and W-MT) and extremely limited gene flow ( $M = 0.41-0.43$ ) between the other Australian population (EA-MT) and the three South African populations (Table 6).

## Discussion

*Mycosphaerella nubilosa* is one of the most important leaf pathogens of *E. nitens* in South Africa and of *E. globulus* in Australia. In turn, these trees represent two of the most important sources of wood fibre derived from plantations. South Africa has the oldest plantation forestry programme based on non-native species in the world (Burgess & Wingfield, 2001) and *M. nubilosa* was the first pathogen of *Eucalyptus* to be reported from this country (Doidge, 1950). It has subsequently appeared in many areas of the world where *E. nitens* and *E. globulus* are grown. Although it has been assumed that the pathogen originated in Australia and that it was accidentally introduced into other parts of the world, this hypothesis has never been tested experimentally.

*Mycosphaerella nubilosa* was originally identified from Victoria in eastern Australia (Cooke, 1891). Since this initial identification, it has been hypothesized that eastern Australia would represent the centre of origin for *M. nubilosa*. Gene diversity and maximum percentage of genotypic diversity for the New South Wales *M. nubilosa* (EA-MT) population was the highest of all populations evaluated during this study. Furthermore, the (EA-MT) population had a greater number of alleles than the other *M. nubilosa* populations. It is expected that older pathogen populations representing centres of origin have higher gene diversity values as a result of the accumulation of mutations over time (McDonald, 1997). Results of this study, therefore, support the view that *M. nubilosa* originated from eastern Australia.

An interesting observation in this study was the presence of shared alleles between Western Australia and South Africa. This finding suggests that *M. nubilosa* could have been introduced into South Africa from Western Australia and not from eastern Australia. However, *M. nubilosa* was not detected in Western Australia until 1999 (Maxwell *et al.*, 2001), despite previous surveys (Carnegie *et al.*, 1997), suggesting that it was introduced relatively recently into that area. There are also no known hosts of *M. nubilosa* that naturally occur in Western Australia. Furthermore, commercial forestry with *E. globulus* (a species native to Tasmania and southern Victoria, Australia) only began in Western Australia during the 1980s. It thus seems unlikely that the fungus was introduced into South Africa from Western Australia. An alternative hypothesis suggests inoculum from eastern Australia, not sampled in this study, served as a source population for both Western Australia and South Africa. The fact that *E. globulus* seed from eastern Australia was the first *Eucalyptus* planting

stock to be planted in South Africa would provide additional support, albeit anecdotal, for this view.

Gene diversities derived in this study for the various *M. nubilosa* populations from South Africa were comparable with each other, and lack of significant population differentiation, combined with the high level of gene flow between the South African *M. nubilosa* populations, indicate that these populations are highly homogenous.

In South Africa, the gene diversity of isolates from a single tree was as great as for isolates from a single plantation, and gene flow between plantations was very high. Similar findings of genetic diversity have been found for *M. fijiensis* on banana, where the majority of genetic diversity for that pathogen exists at the plant and plantation levels (Rivas *et al.*, 2004). The results for *M. nubilosa* are also consistent with those for other *Mycosphaerella* pathosystems. For example, the majority of diversity for *M. graminicola* was observed within a single plot of a wheat field (McDonald & Martinez, 1990; Zhan *et al.*, 2003), and likewise with *M. populorum* from north-eastern America, where more than 90% of the genetic diversity was distributed within a single tree (Feau *et al.*, 2005).

Results reported here reveal that some haplotypes of *M. nubilosa* are shared between South Africa, Tanzania and Europe, indicating gene flow has occurred between these regions, resulting in the establishment and persistence of specific haplotypes in Europe and in Tanzania. Based on gene flow data between Australia and South Africa and the same haplotypes that are shared between South Africa, Tanzania and Europe, it seems likely that *M. nubilosa* was first introduced into South Africa from Australia and that it subsequently spread to other parts of Africa and into Europe. This route of movement is also consistent with the pattern of establishment of *Eucalyptus* plantations in Africa and Europe.

## **Acknowledgements**

We thank the members of the Tree Protection Co-operative Programme (TPCP), the National Research Foundation (NRF), the Mellon foundation and the THRIP initiative of the Department of Trade and Industry, South Africa for financial support. We also thank Mr Aza Mbagi of the Tanganyika Wattle Company Limited for *E. globulus* samples from Tanzania.

## References

- Agapow PM, Burt A, 2001. Indices of multilocus linkage disequilibrium. *Molecular Ecology Notes* **1**, 101–2.
- Alemu G, Roux J, Hunter GC, Wingfield MJ, 2006. *Mycosphaerella* species associated with leaf disease of *Eucalyptus globulus* in Ethiopia. *European Journal of Forest Pathology* **36**, 253–63.
- Burgess T, Wingfield MJ, 2001. Exotic pine forestry in the Southern Hemisphere: a brief history of establishment and quarantine practices. *South African Forestry Journal* **192**, 79–84.
- Carnegie AJ, 2007a. Forest health condition in New South Wales, Australia, 1996–2005. I. Fungi recorded from eucalypt plantations during forest health surveys. *Australasian Plant Pathology* **36**, 213–24.
- Carnegie AJ, 2007b. Forest health condition in New South Wales, Australia, 1996–2005. II. Fungal damage recorded in eucalypt plantations during forest health surveys and their management. *Australasian Plant Pathology* **36**, 225–39.
- Carnegie AJ, Ades PK, 2002. The proportion of leaf spots caused by *Mycosphaerella cryptica* and *M. nubilosa* on *Eucalyptus globulus*, *E. nitens* and their F1 hybrids in a family trial in Tasmania, Australia. *Australian Mycologist* **2**, 53–63.
- Carnegie AJ, Ades PK, 2003. *Mycosphaerella* leaf disease reduces growth of plantation-grown *Eucalyptus globulus*. *Australian Forestry* **66**, 113–9.
- Carnegie AJ, Keane PJ, Podger FD, 1997. The impact of three species of *Mycosphaerella* newly recorded on *Eucalyptus* in Western Australia. *Australasian Plant Pathology* **26**, 71–7.
- Carnegie AJ, Ades PK, Keane PJ, Smith IW, 1998. *Mycosphaerella* diseases of juvenile foliage in a eucalypt species and provenance trial in Victoria, Australia. *Australian Forestry* **61**, 190–4.
- Chen RS, Boeger JM, McDonald BA, 1994. Genetic stability in a population of a plant pathogenic fungus over time. *Molecular Ecology* **3**, 209–18.
- Cockerham CC, Weir BS, 1993. Estimation of gene flow from *F*-statistics. *Evolution* **47**, 855–63.
- Cooke MC, 1891. Australian fungi. *Grevillea* **19**, 60–2.
- Crous PW, 1998. *Mycosphaerella* spp. and their anamorphs associated with leaf spot diseases of *Eucalyptus*. *Mycologia Memoir* **21**, 1–170.
- Crous PW, Groenewald JZ, Mansilla JP, Hunter GC, Wingfield MJ, 2004. Phylogenetic reassessment of *Mycosphaerella* spp. and their anamorphs occurring on *Eucalyptus*. *Studies in Mycology* **50**, 195–214.
- Crous PW, Wingfield MJ, Mansilla JP, Alfenas AC, Groenewald JZ, 2006. Phylogenetic reassessment of *Mycosphaerella* spp. and their anamorphs occurring on *Eucalyptus*. II. *Studies in Mycology* **55**, 101–34.
- Dick M, 1982. Leaf inhabiting fungi of eucalypts in New Zealand. *New Zealand Journal of Forestry Science* **12**, 525–37.
- Doidge ME, 1950. The South African fungi and lichens to the end of 1945. *Bothalia* **5**, 1–1094.

- Feau N, Hamelin RC, Vandecasteele C, Stanosz GR, Bernier L, 2005. Genetic structure of *Mycosphaerella populorum* (anamorph *Septoria musiva*) populations in north-central and northeastern north America. *Phytopathology* **95**, 608–16.
- Hansford CG, 1956. Australian Fungi. III. New species and revisions (continued). *Proceedings of the Linnean Society of New South Wales* **81**, 23–51.
- Hunter GC, Crous PW, Roux J, Wingfield BD, Wingfield MJ, 2004a. Identification of *Mycosphaerella* species associated with *Eucalyptus nitens* leaf defoliation in South Africa. *Australasian Plant Pathology* **33**, 349–55.
- Hunter GC, Roux J, Wingfield BD, Crous PW, Wingfield MJ, 2004b. *Mycosphaerella* species causing leaf disease in South African *Eucalyptus* plantations. *Mycological Research* **108**, 672–81.
- Hunter GC, Cortinas MN, Wingfield BD, Crous PW, Wingfield MJ, 2006. Development of polymorphic microsatellite markers for the Eucalyptus leaf pathogen *Mycosphaerella nubilosa*. *Molecular Ecology Notes* **6**, 900–3.
- Lundquist JE, Purnell RC, 1987. Effects of *Mycosphaerella* leaf spot on growth of *Eucalyptus nitens*. *Plant Disease* **71**, 1025–9.
- Maxwell A, Hardy GESStJ, Dell B, 2001. First record of *Mycosphaerella nubilosa* in Western Australia. *Australasian Plant Pathology* **30**, 65.
- McDonald BA, 1997. The population genetics of fungi: tools and techniques. *Phytopathology* **87**, 448–53.
- McDonald BA, Martinez JP, 1990. DNA restriction fragment length polymorphisms among *Mycosphaerella graminicola* (anamorph *Septoria tritici*) isolates collected from a single wheat field. *Phytopathology* **80**, 1368–73.
- Milgate AW, Yuan ZQ, Vaillancourt RE, Mohammed C, 2001. *Mycosphaerella* species occurring on *Eucalyptus globulus* and *Eucalyptus nitens* plantations of Tasmania, Australia. *Forest Pathology* **31**, 53–63.
- Milgate AW, Potts BM, Joyce K, Mohammed C, Vaillancourt RE, 2005. Genetic variation in *Eucalyptus globulus* for susceptibility to *Mycosphaerella nubilosa* and its association with tree growth. *Australasian Plant Pathology* **34**, 11–8.
- Nei M, 1973. Analysis of gene diversity in subdivided populations. *Proceedings of the National Academy of Sciences, USA* **70**, 3321–3.
- Park RF, 1988. Epidemiology of *Mycosphaerella nubilosa* and *M. cryptica* on *Eucalyptus* spp. in south-eastern Australia. *Transactions of the British Mycological Society* **91**, 261–6.
- Park RF, Keane PJ, 1982a. Three *Mycosphaerella* species form leaf diseases of *Eucalyptus*. *Transactions of the British Mycological Society* **79**, 95–100.
- Park RF, Keane PJ, 1982b. Leaf diseases of *Eucalyptus* associated with *Mycosphaerella* species. *Transactions of the British Mycological Society* **79**, 101–15.
- Park RF, Keane PJ, 1987. Spore production by *Mycosphaerella* species causing leaf diseases of *Eucalyptus*. *Transactions of the British Mycological Society* **89**, 461–70.
- Pinkard EA, Mohammed CL, 2006. Photosynthesis of *Eucalyptus globulus* with *Mycosphaerella* leaf disease. *New Phytologist* **170**, 119–27.

- Rivas G-G, Zapater M-F, Abadie C, Carlier J, 2004. Founder effects and stochastic dispersal at the continental scale of the fungal pathogen of bananas *Mycosphaerella fijiensis*. *Molecular Ecology* **13**, 471–82.
- Stoddart JA, Taylor JF, 1988. Genotype diversity: estimation and prediction in samples. *Genetics* **118**, 705–11.
- Weir BS, 1996. *Genetic Data Analysis II*. Sunderland, MA, USA: Sinauer Associates.
- Workman PL, Niswander JD, 1970. Population studies in south-western Indian tribes. II. Local genetic differentiation in the Papago. *American Journal of Human Genetics* **22**, 24–49.
- Yeh FC, Yang RC, Boyle T, 1999. *POPGENE. Version 1.31. Microsoft Windows Based Freeware for Population Genetic Analysis*. Alberta, Canada: University of Alberta.
- Zhan J, Pettway RE, McDonald BA, 2003. The global genetic structure of the wheat pathogen *Mycosphaerella graminicola* is characterised by high nuclear diversity, low mitochondrial diversity, regular recombination, and gene flow. *Fungal Genetics and Biology* **38**, 286–97.