



Review paper

The catastrophic decline of the Sumatran rhino (*Dicerorhinus sumatrensis harrissoni*) in Sabah: Historic exploitation, reduced female reproductive performance and population viability



P. Kretzschmar^{a,*}, S. Kramer-Schadt^a, L. Ambu^b, J. Bender^a, T. Bohm^a,
M. Ernsing^a, F. Göritz^c, R. Hermes^c, J. Payne^d, N. Schaffer^e, S.T. Thayaparan^f,
Z.Z. Zainal^d, T.B. Hildebrandt^{c,1}, H. Hofer^{a,1}

^a Department of Evolutionary Ecology, Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Str. 17, 10315 Berlin, Germany

^b Sabah Wildlife Department, Tingkat 5, Wisma Muis, 88100 Kota Kinabalu, Malaysia

^c Department of Reproductive Biology, Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Str. 17, 10315 Berlin, Germany

^d Borneo Rhino Alliance, c/o Institute for Tropical Biology and Conservation, University Malaysia Sabah, 88999 Kota Kinabalu, Malaysia

^e SOS Rhino, 680 N. Lake Shore Drive, Suite 807, Chicago, IL 60611, USA

^f College of Veterinary Medicine, School of Veterinary and Life Sciences, Murdoch University, 90 South Street, Murdoch, WA 6150, Australia

HIGHLIGHTS

- The reason for the recent decline of the Sumatran rhino is poorly understood.
- The remaining small isolated rhino populations are deemed to go extinct.
- Historic exploitation led to low numbers of Sumatran rhinos on Borneo.
- PVA analyses identify the fertility of females as key factor for population growth.
- A resource selection study identifies habitat characteristics preferred by the rhinos in Sabah.

ARTICLE INFO

Article history:

Received 27 November 2015

Received in revised form 23 February 2016

Accepted 23 February 2016

Available online 7 April 2016

Keywords:

Megaherbivore

Sumatran rhino

Conservation

Extinction

Resource selection function

Population viability analyses

ABSTRACT

The reasons for catastrophic declines of Sumatran rhinos are far from clear and data necessary to improve decisions for conservation management are often lacking. We reviewed literature and assembled a comprehensive data set on surveys of the Sumatran rhino subspecies (*Dicerorhinus sumatrensis harrissoni*) in the Malaysian state of Sabah on Borneo to chart the historical development of the population in Sabah and its exploitation until the present day. We fitted resource selection functions to identify habitat features preferred by a remnant population of rhinos living in the Tabin Wildlife Reserve in Sabah, and ran a series of population viability analyses (PVAs) to extract the key demographic parameters most likely to affect population dynamics. We show that as preferred habitat, the individuals in the reserve were most likely encountered in elevated areas away from roads, in close distance to mud-volcanoes, with a low presence of human trespassers and a wallow on site, and within a neighbourhood of dense forest and grassland patches

* Corresponding author. Tel.: +49 305168513.

E-mail address: petra.kretzschmar@gmx.de (P. Kretzschmar).

¹ Contributed equally to this work.

preferably on Fluvisols and Acrisols. Our population viability analyses identified the percentage of breeding females and female lifetime reproductive period as the crucial parameters driving population dynamics, in combination with total protection even moderate improvements could elevate population viability substantially. The analysis also indicates that unrestrained hunting between 1930 and 1950 drastically reduced the historical rhino population in Sabah and that the remnant population could be rescued by combining the effort of total protection and stimulation of breeding activity. Based on our results, we recommend to translocate isolated reproductively healthy individuals to protected locations and to undertake measures to maximise conceptions, or running state-of-the-art reproductive management with assisted reproduction techniques. Our study demonstrates that a judicious combination of techniques can do much to illuminate causes of population declines, improve decision making for conservation management and possibly prevent similar developments in populations of other species of similar ecological standing.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

1. Introduction.....	258
2. Methods.....	259
2.1. Study area.....	259
2.2. Data collection.....	259
2.3. Environmental predictors.....	260
2.4. Statistical analysis.....	261
2.5. Literature search.....	263
2.6. Population viability analysis.....	263
2.7. Sensitivity analysis.....	263
3. Results.....	263
3.1. Resource selection.....	263
3.2. Historical distribution of Sumatran rhinos on Borneo.....	265
3.3. PVA model results.....	268
4. Discussion.....	269
4.1. Resource selection and preferred habitat features.....	269
4.2. Population viability analysis.....	272
4.3. Conclusions.....	272
Acknowledgements.....	273
Appendix A. Supplementary data.....	273
References.....	273

1. Introduction

The conservation of highly endangered species is challenging. Basic data about their ecological needs and geographic distribution are often lacking (Funk and Richardson, 2002; Harris et al., 2009; Bland et al., 2015; Abram et al., 2015) and the question arises of how to use the limited and uncertain data to improve decisions for conservation management (Tear et al., 1995; Karanth et al., 2003; Huettmann, 2005). Conservation measures such as forest corridors connecting insular environments and the translocation of individual animals into protected areas have been proposed for endangered species suffering from habitat loss and fragmentation (Griffith et al., 1989; Bennett, 2003; Rayan and Linkie, 2015). These measures require an understanding which environmental variables, such as terrain, climate, substrate or land cover determine the distribution of a species (Ferrier et al., 2002; Rushton et al., 2004). The rise of habitat modelling techniques combined with Geographical Information Systems (GIS) has led to the development of new methods (Guisan and Zimmermann, 2000) to identify relevant habitat characteristics for the occurrence of a species. Habitat suitability models identify areas with the best resources for the conservation of the respective species and habitats (Leblond et al., 2014).

In addition to habitat loss and fragmentation, the size of a population, its vulnerability to demographic and environmental stochasticity and negative impacts of human activity also influence its viability (Mace and Lande, 1991; Melbourne and Hastings, 2008; Meek et al., 2015). Population viability analysis (PVA) methods help to identify factors important for the persistence or decline of a species. They improve decisions by reducing and quantifying uncertainty and thereby improve decision making for conservation efforts (Beissinger and McCullough, 2002; Robinson et al., 2015), or at least highlight the key gaps for which further scientific research should provide essential data.

The Sumatran rhino (*Dicerorhinus sumatrensis*) is a good example to illustrate the challenges in the conservation of a highly endangered species. The two still extant subspecies, *D.s. sumatrensis* in Sumatra, Indonesia, and *D.s. harrissoni*

(hereafter called Borneo rhino), in Borneo in the states of Sabah, Malaysia, and Kalimantan, Indonesia, are split into very small and highly fragmented populations whose numbers are continuously decreasing. Around one hundred or so animals of the Sumatran subspecies are estimated to exist in three isolated populations in the wild whereas the Borneo rhino is nearly extinct (IUCN, 2015). Its disappearance occurred almost undetected and the reasons for its recent decline are little understood. The Sumatran rhino is an example of a large-bodied megaherbivore (Owen-Smith, 1988). It occurs in a tropical rainforest without any serious predators and may have always lived at low densities. Learning about the causes of the decline is an essential part in order to develop targeted conservation measures for its continued existence and that of its Sumatran sister subspecies. It may also be a model to understand similar past, current or future declines in species with a similar ecological position within tropical and subtropical habitats.

Several conservation strategies have been proposed to stop the decline and improve reproduction of the Sumatran rhino (Clements et al., 2010; Zafir et al., 2011; Nardelli, 2014). In terms of ecological approaches, translocation of individual animals into a small designated area to increase the chance of breeding encounters and to support natural breeding with artificial reproduction techniques is one key measure. However data suitable and relevant for such measures, such as the habitat requirements of the Sumatran rhino, hardly exist, as demonstrated by a comprehensive literature review of the last 20 years (Rookmaaker, 2014, see Appendix A).

This study reports and analyses an extensive data set from the yearly and monthly rhino surveys (SOS Rhino and BORA between 2000–2013, unpublished survey reports) in the Tabin Wildlife Reserve (TWR) in Sabah, Malaysia. We use a use-versus-availability approach (Boyce et al., 2002; Johnson et al., 2006) in order to assess resource selection of the population in this reserve. With the help of high-resolution remote sensing data, several environmental predictors were derived which indicate vegetation density at different scales. Additionally, we conducted a literature search to identify the main factors responsible for the decline of the Borneo rhino since 1874. We combine all information in a novel way with a PVA analysis to assess the probability of extinction for a hypothetical rhino population in Sabah in the 20th century, as well as for a locally restricted population in the TWR.

2. Methods

2.1. Study area

The Tabin Wildlife Reserve (TWR), at 5° 15′–5° 10′N, 118°30′–118°45′E, is a medium-sized (1205 km²) protected area in the eastern part of the Malaysian State of Sabah (Fig. 1(A)). It consists mainly of lowland forest (*Dipterocarpus* sp.) and is surrounded by palm oil (*Elaeis guineensis*) plantations (Dawson, 1993). More than 80% of the reserve has been selectively logged between 1960 and 1984 (Sabah Government, 2014). Only the so-called “core area” in the centre and several virgin jungle reserves remain unlogged. Logging activities outside the reserve continued until 1989 (Sabah Government, 2014). The TWR holds several natural mineral sources, including four mud volcanoes and small patches of grassland which developed mainly as a consequence of logging between 1960 and 1984. It has tropical climate with temperatures ranging between 22° C and 32° C (Dawson, 1993). The annual rainfall amounts up to 3000 mm; the wettest period is between October and March (WWF-Malaysia, 1986, cited in Dawson, 1993). The topography of the area is characterised by steep hills and deep valleys which covers an altitudinal range from sea level to about 500 m. Tabin is well drained by a number of seasonal and perennial streams which originate in the hills and form the five major rivers of the area (Dawson, 1993).

2.2. Data collection

Data on rhino occurrence in the TWR were gathered from unpublished reports kindly provided by SOS Rhino and the Borneo Rhino Alliance, and by conducting a systematic camera trap study at the end of the study period. Both organisations conducted regular surveys on foot between October 2000 and February 2013. The surveys were mainly concentrated in the western and central area of the TWR (Fig. 1(B)) and lasted on average up to 8 days (Table A1). Rhino presence was either detected by direct sightings or more frequently by indirect signs such as footprints, scratch-marks, feeding-marks, faeces and wallows. Indirect signs such as wallows were often created by other animals and were therefore only used in combination with footprints in order to avoid false-positive detection. Individual animals could not be identified with this method. The location of rhinos as well as the position of human footprints, cartridge cases or tree logs, used to identify trespassers and disturbance, was measured with a portable Geographical Positioning System (GPS). In case of low GPS coverage, which occurred more frequently between 2000 and 2005, the location of rhino presence indicators was either estimated from a nearby area with better GPS coverage, such as rivers, clearings or hills, or estimated with the help of a topographic map. The estimated error of non-GPS locations averaged approximately 250 m.

During the surveys, GPS locations of the survey route were collected sporadically between October 2000 and July 2010 and systematically between June 2012 and February 2013 at intervals of 100 m, as part of an intensive survey accompanying the camera trap study (see below).

Between 2003 and 2005, the time and the position when and where the survey started, ended or interrupted was noted. This enabled us to sum up the time spent walking and the distance walked per survey (Table A1). For the years 2000 until 2002, information on survey length and duration was lacking, for the years from 2006 until 2008 this information




Fig. 1. Study site in the north east of Sabah on the island of Borneo (A), distribution of survey routes in the study site between 2000 and 2013 (B) and locations of rhinos indicated by footprints in combination with scratch-marks, feeding-marks, faeces and wallows (C).

was fragmentary. In the latter cases the duration of a survey was estimated using either the kilometres walked within a day or the duration of the following days as a reference. The distance walked per day was calculated as the shortest distance between the last coordinate of the day, the starting point and the consecutive coordinates. Additionally, the search intensity was estimated using the walking speed and the direction of a survey as a reference. The year of the survey, its duration, searching intensity and distance walked were used to estimate search effort within the statistical analysis.

Between June 2012 and November 2013, we conducted a camera trap survey in areas with known rhino presence in previous years (Figure A1). The survey was carried out in the central and northern part of the TWR and ranged over an area of 301 km². A total of 57 camera trap stations were placed along a square grid of 6 km². The grid size was chosen according to the minimum home range size of Sumatran rhinos of 10 km² (Strien, 1986). Cameras were spaced at an average of 2.4 km at locations preferred by rhinos, such as game trails, mud wallows or hill crest. This set up enabled us to sample all rhinos possibly ranging in the survey area. Two remotely triggered infrared cameras (Reconyx PC 800 Hyperfire Professional IR, Reconyx, Inc, Wisconsin, USA) were placed at each camera trap station. The cameras set to operate for 24 h each day. Stations were checked every two to three months and the number of days each camera trap station was functional over the total survey period was recorded.

2.3. Environmental predictors

We used seven high resolution satellite maps (RapidEye, BlackBridge, Berlin, Germany) covering the whole TWR to derive fine-scale land cover maps. The satellite maps were provided in L3A-format with a spatial resolution of 5 m. We applied radiometric corrections established for multi-temporal and multi-sensor data applications (Chavez, 1996; Chander et al., 2009) to reduce scene-to-scene variability. The image-based atmospheric corrections included 'dark object subtractions' and exo-atmospheric reflectance conversions (TOA).

We defined six land cover categories of biological relevance to the rhinos: dense forest, open forest, grassland, plantations, bare land and streams and performed a remote-sensing multispectral classification using ERDAS Imagine (Intergraph Corporation, Madison, USA) and ArcMap 10 (ESRI Inc., Redlands, USA). Clouds and cloud shadows were eliminated (Langner et al., 2012). The classifications were ground truthed with data for grassland, plantations and dense forest.

Table 1
Predictors used for assessing habitat use of the rhino population in the TWR.

Variable	Abbreviation	Type
Rhino presence (yes/no)	rhpr	Categorical
Year (year)	year	Continuous
Sampling effort		
Time spend walking (min)	dur	Continuous
Distance walked (km)	dist	Continuous
Search intensity (high/low)	surv	Categorical
Human disturbance variables		
Trespasser (yes/no)	hum	Categorical
Distance to logging road (m)	d_road	Continuous
Human population density in a square of 30'' (arc seconds)	hpd	Continuous
Topographic variables		
Elevation (m)	dgm	Continuous
Slope (%)	sl	Continuous
Land cover variables		
Dense forest ^a	f_dfo_x	Continuous
Open forest ^a	f_ofo_x	Continuous
Grassland ^a	f_gr_x	Continuous
Oil palm plantations ^a	f_pl_x	Continuous
Barren land ^a	f_bl_x	Continuous
Classify (1–5)	clas	Categorical
Food, shelter		
Distance to streams (m)	d_st	Continuous
Soil type (12, 13, 23, 32)	sp	Categorical
Distance to mudvolcano (m)	d_mvno	Continuous
Mud wallow (yes/no)	wal	Categorical

^a = 100 m²; 250 m²; 500 m²; 1000 m².

We deployed a focal or so-called neighbourhood analysis to include land cover connectivity in the surroundings of a rhino location (ArcGIS 10, ESRI Inc., Redlands, USA). This analysis was employed for five land cover categories excluding streams and encompassed the area sizes of 100 m², 250 m², 500 m² and 1000 m² of potential relevance for daily movement distances of rhinos whilst foraging. Additional environmental variables of potential biological relevance for habitat use of the rhinos such as climatic, topographical variables or predictors indicating human impact were included in the analysis (Table 1, A2 and Table A4). Distances were calculated from each rhino location as well as for survey and systematic locations (see below) to the nearest border of these features.

2.4. Statistical analysis

We employed a use-versus-availability approach based on resource selection functions (RSF) (Boyce et al., 2002; Johnson et al., 2006) in a binomial generalised linear modelling (GLM) framework. Environmental covariates at the location where the rhino was present (use) were contrasted with covariates at locations taken from an area deemed to be available for selection (available). The size and spatial extent of the available sample was defined by the border of the TWR. In order to avoid multi-collinearity, only environmental variables were included in the analysis which were not highly correlated (Spearman rank $p < 0.75$; Table A2) and only one size scale of neighbourhood variables were included in each model. Two sets of availability locations were included in the analysis: (1) evenly distributed data points across the study site with a distance of 1 km to each other ("systematic"); (2) GPS locations collected during surveys ("survey") where rhinos were assumed to be present but no presence sign was detected. The inclusion of survey locations enabled us to incorporate variables indicating survey effort in the statistical analysis, such as year of the survey, its duration, searching intensity and distance walked. Spatial autocorrelation was addressed by using spatial filtering to reduce the number of occurrence and survey records in oversampled regions (Dormann et al., 2007; Veloz, 2009). We filtered all locations within a year by selecting only data points with more than 1 km distance to each other. This resulted in a data set of 109 rhino locations (use), 814 "systematic" (availability) and 1068 "survey" (availability) locations. We designed two sets of candidate models (defined by the two availability data sets) for Borneo rhino occurrence guided by three general hypotheses: (1) Human disturbance is the only driver for Borneo rhino presence or absence. (2) Habitat use is linked to food availability, access to safe areas and access to other ecological resources. (3) Human disturbance, food availability, access to safe areas and access to other ecological resources are all important in terms of habitat use of rhinos. A null model containing only the intercept was also included in the candidate set for comparison as well as models only containing the variables indicating survey effort. A set of 23 candidate models was designed for the systematic data and a set of 37 candidate models for the survey data (Table 2). In order to avoid linearity assumptions, we preliminarily explored the shape of the response for each predictor variable before fitting them to the final equations (Austin, 2002). To this end, we built binomial Generalised Additive Models (GAMs, R package 'mgcv', Wood, 2011) using rhino locations (use) versus systematic or survey points (availability), respectively, and

Table 2

Summary of models used to establish habitat use of Borneo rhinos employing two different sampling sets: rhino presence versus random/survey points. The models are organised in groups corresponding to different hypothesis of landscape factors potentially affecting habitat use of the species.

No Rhino locations versus random points	No rhino locations versus survey points
0 Null model	0 Null model
Human disturbance	Controlling variables
1a d_road+hpd+f_pl_x+f_dfo_x+f_gr_x	1a year
1b d_road+hpd+f_pl_x+f_gr_x	1b year+dur+dist+surv
1c hpd+f_pl_x+f_dfo_x+f_gr_x	1c year+dist
1d d_road+hpd+f_pl_x+f_gr_x+f_bl_x	1d surv
1e f_pl_x+f_gr_x+f_bl_x	1e dist+surv
Food, terrain and other ecological resources	1f dur
2a d_st+d_mvov+f_dfo_x+f_gr_x+dgm+sl+sp	1g dur+surv
2b d_st+d_mvov+f_dfo_x+f_gr_x+dgm +sp	Human disturbance
2c f_gr_x+f_ofo_x+d_st+f_dfo_x+dgm+sl+sp	2a year+dur+surv+d_road+hpd+f_pl_x+hum+f_gr_x+f_bl_x
2d f_gr_x+f_ofo_x+d_st+f_dfo_x+dgm+sl	2b year+dur+surv+d_road+hpd+f_pl_x+hum+f_gr_x
2e f_gr_x+f_ofo_x+d_st+f_dfo_x+dgm	2c year+dur+surv+hpd+f_pl_x+hum+f_gr_x
2f f_dfo_x+dgm	2d year+dur+dist+surv+hpd+hum
2g f_dfo_x+dgm+sl	2e year+dur+dist+surv+hum+f_gr_x
2h f_dfo_x+dgm+sl+d_mvov	2f dur+dist+surv+d_road+hpd+f_pl_x+hum+f_gr_x
2i f_dfo_x+dgm+d_mvov	2g dur+surv+d_road+hpd+f_pl_x+hum+f_gr_x
2j f_dfo_x+dgm+d_mvov +f_gr_x	2h dur+dist+surv+hpd+f_pl_x+hum+f_gr_x
2k f_ofo_x+d_st+clas	2i dur+surv+hpd+f_pl_x+hum+f_gr_x
2l f_dfo_x+d_mvov	2j dur+dist+surv+hpd+hum
2m f_dfo_x+d_mvov+sp	2k dur+surv+hpd+hum
2n clas	Food availability, access to safe areas and other ecological resources
Human disturbance, food availability, access to safe areas and other ecological resources	3a year+dur+surv+d_st+wal+d_mvov+f_dfo_x+f_gr_x+dgm+sl+so
3a d_road+hpd+f_pl_x+f_gr_x+f_dfo_x+d_st+d_mvov+dgm+sl+sp	3b year+dur+surv+d_st+wal+d_mvov+f_dfo_x+f_gr_x+dgm+sl+sp
3b hpd+f_pl_x+f_gr_x+f_dfo_x+d_st+d_mvov+dgm+sl	3c year+dur+surv+d_st+wal+d_mvov+f_dfo_x+f_gr_x+dgm+sp
3c hpd+f_gr_x+f_dfo_x+d_st+d_mvov+dgm+sl	3d year+dur+surv+d_st+wal+d_mvov+f_dfo_x+f_gr_x+dgm
	3e year+dur+surv+wal+d_mvov+dgm
	3f year+dur+dist+surv+d_st+f_gr_x+f_ofo_x+f_dfo_x+dgm+sl+sp
	3g year+dur+dist+surv+f_dfo_x+dgm+sl
	3h year+dur+dist+surv+f_ofo_x+d_st+clas
	3i year+dur+dist+surv+f_dfo_x+d_mvov
	3j year+dur+dist+surv+f_dfo_x+d_mvov+sp
	3k dur+surv+d_st+wal+d_mvov+f_dfo_x+f_gr_x+dgm+sl+sp
	3l dist+dur+surv+d_st+wal+d_mvov+f_dfo_x+f_gr_x+dgm+sl+sp
	3m dur+surv+wal+d_mvov+dgm
	3n dist+dur+surv+wal+d_mvov+dgm
	Human disturbance, food availability, access to safe areas and other ecological resources
	4a year+dur+surv+hpd+hum+f_pl_x+f_gr_x+d_st+wal+d_mvov+f_dfo_x+dgm+sl
	4b year+dur+surv+hpd+hum+f_gr_x+d_st+wal+d_mvov+f_dfo_x+dgm+sl+sp
	4c year+dur+surv+hpd+hum+d_road+f_gr_x+d_st+wal+d_mvov+f_dfo_x+dgm+sp
	4d year+dur+surv+hpd+hum+f_gr_x+d_st+wal+d_mvov+f_dfo_x+dgm

fitted smoothing splines with 3 degrees of freedom on the full model. Then, linearity was assessed by visual inspection of the partial residual plots and where appropriate, logarithmic or quadratic transformations of predictor variables were applied (Crawley, 2007). GLMs were then fitted and models were compared and hierarchically ordered using Akaike's Information Criterion (AICc) corrected for small sample size. We repeated model fitting using a randomisation method by selecting 100 rhino locations and an equal amount as well as a fourfold amount of systematic or survey points with replacement from the rhino locations and systematic or survey points, respectively. The size of the availability sample was carefully chosen to avoid a bias in our estimates (Northrup et al., 2013). This routine was iterated 1000 times. Each time, we fitted the set of candidate models, calculated the AICc and recorded the model with the lowest AICc value. The probability of model selection was estimated counting the number of randomisations in which each candidate model was scored as best. To then test the contribution of each variable within the selected model, we fitted the best model with the highest score probability to the respective data set and selected final models within $<2\Delta\text{AICc}$ (package 'MuMIn'), i.e. the difference to the AICc of the best ranked model, as all of them should be considered competitive for interpretation (Burnham and Anderson, 2002). We then used model averaging to derive final coefficient values. All statistical analyses were performed using the R statistical software R 3.1.1 (R Development Core Team, 2014).

2.5. Literature search

We reviewed the literature on Sumatran rhinos (Rookmaaker, 2014) from 1874 onwards focusing on information about the number and distribution of rhinos on Borneo. Particular interest was paid to the year of the census, the quality of the data and the information provided about hunting of wildlife by local people or Europeans in the area. Additionally, the literature on surveys conducted in the TWR was reviewed and the estimated rhino numbers as well as the number of survey days, kilometres walked and methods analysed.

2.6. Population viability analysis

We used the stochastic population model VORTEX (Version 10, Lacy and Pollak, 2014) to identify possible factors responsible for the decline of the Borneo rhino in Sabah, Malaysia. We analysed two different scenarios, firstly a hypothetical rhino population in the 20th century with a yearly hunting quota of 120 animals between 1930 and 1950 (scenario (a)) and secondly, a population known to exist in the TWR (scenario (b)). Since population data on free-ranging Borneo rhinos do not exist, the model is based on information available from the Sumatran subspecies combined with data on other rhino species (Appendix B). Based on this information, we defined a standard set of parameters for both scenarios (Table 3) and modified individual parameters thereon. Population size (ps) and carrying capacity (cc) of scenario (a) were estimated based on a natural distribution of 10 rhinos per 100 km² (Strien, 1986) assuming a theoretical utilisation of 25% (ps) and 50% (cc) of the total land area of Sabah of 74,398 km² (Encyclopaedia of the Nations, 2015). This resulted in an initial population size of 3720 rhinos and a carrying capacity of 5580 rhinos. Yearly hunting was set to 60 adult males and 60 adult females between 1930 and 1950, using information from historical reports (see 3.2). For scenario (b) an initial population size of 15 individuals was chosen based on surveys conducted in TWR between 1980 and 1989 (Table A3). The carrying capacity was calculated on the basis of the size of TWR (1205 km²) as described above. Three different models were calculated with different levels of harvesting to investigate the impact of hunting on the population: for scenario (b1) no harvesting was included in the model; for scenario (b2) a single harvesting event was included, which reflects the real situation where an adult male and an adult female were captured for breeding purposes; for scenario (b3) an adult male and an adult female were harvested every ten years. Results were predicted over 115 years (scenario (a)) and 35 years (scenario (b)) respectively. Output of the PVA included mean stochastic growth rate (\pm SD), probability of population extinction (\pm SD), mean time to extinction (\pm SD), mean population size (\pm SD) and decline in genetic variability expressed as the expected genetic diversity (Gatti et al., 2011).

2.7. Sensitivity analysis

To evaluate the sensitivity of the simulation results to variation in specific parameters, we calculated a sensitivity index (S_y) as following:

$$S_y = (\Delta y/y)/(\Delta x/x)$$

where $\Delta y/y$ is the observed relative change in the predicted variable y resulting from the relative change $\Delta x/x$ in the predictor parameter x (Pulliam et al., 1992). We examined model sensitivity to variations in population size, lifetime reproductive period, length of harvesting period, number of males/females harvested, percentage of female breeding, carrying capacity, juvenile mortality (female/male), adult mortality (female/male) and percentage of mate monopolisation. Variation of predictor variables were limited to a moderate \pm 10%.

3. Results

3.1. Resource selection

Comparing rhino locations ($n = 100$) with survey data ($n = 400$, 1000 repetitions) revealed that rhino presence in the TWR was best described by the four candidate models 4c, 4a, 4b and 3n -, in decreasing order of importance (Table 2). The best model with a selection probability of 34% (4c) and the two adjacent models (4a, 4b) are consistent with the hypothesis that human disturbance, food availability, access to safe areas and access to other ecological resources predict habitat use. Fitting the candidate models to the data set resulted in 3 best models with a $\Delta AICc < 2$, each consisting of ten and eleven predictor variables respectively (Table 4). Model averaging revealed that the search effort variables 'year', 'duration of survey' and 'low search intensity' were all negatively correlated with rhino presence (Table 5). Rhino presence was thus highest in the first years of the study period (from 2000 onwards), during short surveys and when search intensity was high. The environmental predictors indicated that rhinos preferred elevated areas in the TWR with a low presence of human trespassers and a wallow on site, away from roads and in close distance to mud-volcanoes. Within a neighbourhood of 1000 m², rhinos prefer a high proportion of dense forest and grassland and favour fine textured thionic Fluvisols category 13, and orthic Acrisols category 23 and 32 (Table 5, Table A4). The environmental predictor 'distance to streams' was not included in the model anymore while the predictor variable 'distance to mudvolcanoes' was only present in two of the three models.

Table 3

Life history parameters used for population viability analysis of the historic rhino population scenario (a) and the rhino population living in the TWR scenario (b).

Parameter	Baseline values
Number of populations	1
Number of iterations	1000
Number of years scenario a/scenario b	115/35
Duration of each year (days)	365
Extinction definition	1 sex remains
Inbreeding depression ^a	6.29
Percent due to recessive lethal alleles (%)	50
Polygynous	yes
Age of first offspring female/male (years)	6/7
Maximum lifespan (years)	30
Maximum lifetime reproductive period of females/males (years)	20/25
Maximum number of broods/progeny per year	1/1
Sex ratio of males at birth (%)	50
Density dependent reproduction	none
Breeding females (%)	25
SD of breeding females due to environment variation (%)	0
Distribution of broods per year: 0/1 broods (%)	0/100
<i>Mortality of females/males from</i>	
age 0–1 years (%) scenario a/scenario b	20/20
age 1–2 years (%) scenario a/scenario b	5/5
age 2–3 years (%) scenario a/scenario b	5/10
age 3–4 years (%) scenario a/scenario b	5/10
age 4–5 years (%) scenario a/scenario b	5/5
age 5–6 years (%) scenario a/scenario b	4/10
age 6–7 years (%) scenario a/scenario b	4/10
> 7 years (%) scenario a/scenario b	–/5
Catastrophes	none
Males in breeding pool (%)	80
Initial population size scenario a/scenario b	3720/15
Age distribution	stable age dist.
Carrying capacity (K) scenario a/scenario b	5580/120
SD (K)	0
Population harvested?	yes
First year of harvest scenario a/scenario b1/b2 (year)	30/32/0
Last year of harvest scenario a/scenario b1/b2 (year)	50/33/35
Interval between harvests scenario a/scenario b (year)	1/30
Optional criteria for harvest scenario a/scenario b	1/1
Optional criteria for individuals	1/1
Number of adult females (6 + years) harvested scenario a/scenario b	60/1
Number of adult males (6 + years) harvested scenario a/scenario b	60/1
Supplementation	none

^a Mean effect of inbreeding on fecundity and first year survival (O'Grady et al., 2006).

Table 4

Summary of the best models for predicting habitat use of the Borneo rhino using the survey data to describe availability: 100 presence points were compared with 400 survey points, running 1000 repetitions.

	Int	sp	hum	surv	wal	d_mvo	d_st	d_road	f_dfo_1000	f_gr_1000	dur	dgm	year	df	logLik	AICc	delta	weight
Model 1	1342	+	+	+	+			+	+	+	+	+	+	13	–70.279	167.8	0	0.419
Model 2	1266	+	+	+	+	+		+	+	+	+	+	+	14	–69.224	167.8	0.08	0.403
Model 3	1326	+	+	+	+		+	+	+	+	+	+	+	14	–70.046	169.5	1.72	0.177

A comparison of rhino locations ($n = 100$) with systematic data ($n = 400$, 1000 repetitions) indicate that rhino presence in the TWR was best described by model 2c (Table 2) predicting that habitat use of Sabah rhinos was linked to food availability, access to safe areas and access to other ecological resources (model selection probability 48%). The second best model, with a selection probability of 35%, presumed that human disturbance, food availability, terrain and access to other ecological resources are equally important (3a, Table 2). Fitting the best candidate model (2c) to the data set resulted in 11 best models with a $\Delta AICc < 2$ (Table 6). Model averaging resulted in three predictor variables which were consistently present in all models (soiltypes, focal grassland 100 m² and elevation) while the remaining variables were only represented in four models (Table 7). Habitat suitability for rhinos increased with increasing elevation and declined with increasing distance to streams and increasing gradient. Within an area of 100 m², rhinos preferred more densely forested areas and avoided areas with a high proportion of grassland and open forest. Locations with fine textured Thionic Fluvisols were more frequently used than areas of a soil category 23 and 32, in decreasing order of importance (Table 7).

Table 5

Results of the model averaging analysis using the survey data to describe absence of rhinos: 100 rhino presence points were compared with 400 survey (rhino absence) points running 1000 repetitions.

Abbreviation	Variable description	Averaged model coefficients	2.50%	97.50%	Present in model
Intercept		1305	802	1807	1, 2, 3
year	Year	−0.66	−0.91	−0.41	1, 2, 3
dur	Duration (min)	−0.0008	−0.0012	−0.0004	1, 2, 3
surv (low)	Search intensity low	−1.67	−3.38	0.05	1, 2, 3
sp (13)	Soil category 13	5.84	2.41	9.26	1, 2, 3
sp (23)	Soil category 23	7.26	4.09	10.44	1, 2, 3
sp (32)	Soil category 32	3.18	0.42	5.95	1, 2, 3
f_gr_1000	Focal grassland 1000 m ²	12.60	1.96	23.20	1, 2, 3
f_dfo_1000	Focal dense forest 1000 m ²	10.22	4.45	15.99	1, 2, 3
hum (yes)	Human trespassers	−18.84	−1952	1915	1, 2, 3
wal (yes)	Presence of wallow	3.85	1.74	5.96	1, 2, 3
d_road	Distance road (m)	0.0001	0.00004	0.0002	1, 2, 3
dgm	Elevation (m)	0.02	0.01	0.02	1, 2, 3
d_mv0	Distance mudvolcano (m)	−0.0001	−0.0003	0.00005	2

Table 6

Summary of the best models for predicting habitat use of the Borneo rhino using the systematic data to describe availability: 100 presence points were compared with 400 random points, running 1000 repetitions.

	Int	sp	d_st	f_dfo_100	f_gr_100	f_ofo_100	sl	dgm	df	logLik	AICc	delta AICc	weight
Model 1	−2.898	+			+			+	6	−169.992	352.2	0.00	0.141
Model 2	−2.654	+			+	+		+	7	−168.997	352.3	0.08	0.136
Model 3	−3.569	+		+	+			+	7	−169.088	352.5	0.26	0.124
Model 4	−2.858	+	+		+			+	7	−169.187	352.7	0.46	0.112
Model 5	−2.642	+	+		+	+		+	8	−168.401	353.2	0.97	0.087
Model 6	−3.459	+	+	+	+			+	8	−168.508	353.4	1.19	0.078
Model 7	−2.512	+			+	+	+	+	8	−168.603	353.6	1.38	0.071
Model 8	−2.763	+			+		+	+	7	−169.671	353.6	1.43	0.069
Model 9	−3.481	+		+	+		+	+	8	−168.646	353.7	1.46	0.068
Model 10	−2.711	+	+		+		+	+	8	−168.810	354.0	1.79	0.058
Model 11	−3.099	+		+	+	+		+	8	−168.822	354.0	1.82	0.057

Table 7

Results of the model averaging analysis using the systematic data to describe absence of rhinos: 100 rhino presence points were compared with 400 random (rhino absence) points running 1000 repetitions.

Abbreviation	Variable description	Averaged model coefficients	2.50%	97.50%	Present in model
Intercept		−2.96	−4.76	−1.16	
sp (13)	Thionic Fluvisols fine textured ^a	2.77	0.69	4.85	1–10
sp (23)	Orthic Acrisols medium to fine textured with Ferric Acrisols ^a	1.83	0.20	3.45	1–10
sp (32)	Orthic Acrisols medium to fine textured with Humic Acrisols ^a	0.78	−0.75	2.30	1–10
f_gr_100	Focal grassland 100 m ²	−5995	−570700	558709	1–10
f_ofo_100	Open forest 100 m ²	−1.15	−2.82	0.52	2, 5, 7, 10
f_dfo_100	Dense forest 100 m ²	0.91	−0.47	2.29	3, 6, 9, 11
d_st	Distance stream (m)	−0.001	−0.00169	0.0004	4, 5, 6, 10
dgm	Elevation (m)	0.01	0.003	0.01	1–10
sl	Slope (%)	−0.03	−0.08	0.03	7–10

^a For more details see Table A4.

3.2. Historical distribution of Sumatran rhinos on Borneo

There is a wide range of historical information on Borneo rhinos from early explorers visiting the island as well as from residents and local inhabitants. However estimates about the abundance and distribution of rhinos on Borneo are vague. They reported that rhinos were common in Sabah and in Kalimantan (Indonesia) and less abundant in Sarawak (Malaysia) (Bartlett, 1891, Table 8). The reports are either based on hearsay or on estimates resulting from field trips in a restricted area. The rapid decrease of rhino numbers was most likely induced by the pacification of indigenous people in the interior of the island of Borneo by the respective governments at the beginning of the 20th century. Trade started between the interior and the coast and Chinese traders at the coast sourced rhino horn and rhino nails as aphrodisiacs and for medicinal purposes (Harrisson, 1946, Appendix C). From 1930 onwards firearms became numerous and cheap. With such arms, the indigenous people in the interior of Borneo began a wholesale slaughter of rhinos (Banks, 1937; Harrisson, 1946). On the basis of the reports by Harrisson (1956) it can be estimated that a minimum of 100 animals were killed per year between 1925 and 1931 in Sarawak alone (Appendix C). There are no data on the number of rhinos hunted in Sabah before 1961, but reports suggest

Table 8
Historic and recent estimates of the number and distribution of the Sumatran rhino in Borneo.

Source	Census year	Sabah	Sarawak	Kalimantan	Quality of data	Hunting evidence
Wallace (1874)	/	not abundant	not abundant	not abundant	guesstimate	rhino horn sold frequently in Kalimantan
Pryer (1881)	1881	frequent	/	/	estimate	/
Bock (1881)	/	/	/	exists	guesstimate	natives ask high price for rhino horn frequently shot by natives
Anonymous (1890)	1883–1888	frequent	/	/	guesstimate	/
Bartlett (1891)	/	numerous	rare	common	guesstimate	/
Jentink (1897)	/	/	/	common	estimate	/
Shelford (1916)	/	common	/	/	estimate	high price for rhino horn
Ridley (1916)	/	common	/	/	estimate	/
Mjoeberg (1929)	1925	less common	common	/	estimate	numerous rhinos are killed every year
Banks (1931)	/	/	becoming scarce	/	estimate	36 trophies in Beluga, Sarawak in 2 years
Banks (1935)	1934/35	/	decreased	/	estimate	/
Heynsius Viruly and Heurn (1935)	/	/	/	scarce and decreasing	guesstimate	/
Banks (1937)	/	/	a few individuals	/	estimate	price for rhino between \$300–\$400
Comyn-Platt (1937)	/	/	/	few	guesstimate	Chinese will pay any price for the horn
Schneeberger (1945)	/	/	/	fairly large numbers	estimate	rhinos frequently hunted by Iban Dayaks
Harrison (1946)	1945/1946	5	none	2	Sarawak: loc. Surv; Kalimantan, Sabah: estimate	16 rhinos killed by a native
Harrison (1948)	1945/46	considerable number	none—few scattered individuals	some	estimate	/
Banks (1949)	/	/	/	/	guesstimate	native hunters have driven rhinos into remote areas
Harrison (1949)	1946/46	many	none	some	estimate	many hundred rhinos killed in Sarawak during this century
Harrison (1955)	/	some	2–3	handful	guesstimate	/
Harrison (1956)	1956	very few	2	a few	guesstimate	/
Harrison (1961)	/	~10	/	/	localised surveys	/
Burgess (1961)	1985	20–30	/	/	guesstimate	evidence of rhino poaching in Sabah
Bruton (1963)	/	20–30	/	/	status report	/
Harrison (1965)	1961–1965	11–13	/	10 (5)	guesstimate	sea smuggling to Singapore stopped
Anderson (1968)	/	/	virtually extinct	/	status report	/
Silva (1968)	/	nearly extinct	/	/	guesstimate	/
Carson (1968)	/	extinct soon	/	/	status report	/
Mackinnon (1970)	1969/1970	<100	/	/	estimate	hunters get 2000 MS out of a rhino.

(continued on next page)

Table 8 (continued)

Source	Census year	Sabah	Sarawak	Kalimantan	Quality of data	Hunting evidence
Strien (1974)	/	few	extinct?	few	estimate	/
Harrison (1975)	/	10–20	none	1–2	guestimate	/
Rookmaaker (1977)	1976	10–20	0–3	5	estimate	/
Zon (1977)	/	/	2	/	estimate	/
Strien (1979)	1979	/	/	on verge of extinction	localised survey	/
Davies and Payne (1982)	1979–1981	15–30	/	/	survey	5+ rhinos have been killed during last 5 years
Flynn (1981)	1980	S: 8–10+	/	/	localised survey	2 rhinos poached in Silabukan since 1975.
Strien (1981)	1980 / 81	/	/	probably extinct	survey	1 young rhino trapped 1981 in Silabukan
Andau and Payne (1982)	1982	S: 5–10	/	/	localised survey	/
Payne (1986)	/	rhinos in central Sabah	/	/	status report	2 rhinos poached between 1981 and 1984.
Andau (1987)	1982	T: 7	/	/	localised survey	/
Andau (1987)	1986	T: 6, M, B: 3, few isolated rhinos	/	/	survey	/
Nor et al. (1989)	1988	T: 3	/	/	localised survey	/
Khan (1989)	/	> 38 (T: 20+, K: 8, DV: 10)	5–15	perhaps near border to Sabah	status report	/
Payne (1990)	1989	T: 15–20+, non-viable populations in other regions	/	/	survey	illegal hunting still occurs
Boonratana (1997)	1988	11–33	/	/	survey	/
Martin (1989)	1988	30–100	very few	/	survey in horn trade	Sabah: A minimum of 10 pairs of horn are annually exported. Sarawak: no hunting since 1960
Rabinowitz (1992)	1992	DV: 13–23	/	/	localised survey	hunting evidence in the area
Khan (1993)	1995	40–60	/	/	status report	/
Tajuddin Abdullah et al. (1995)	1995	DV: no fresh tracks found	/	/	localised survey	hunting activity in the area
Meijaard (1996)	1994	/	/	20 sightings since 1985	survey	/
Khan et al. (1999)	/	50–70	/	/	status report	/
Strien (2002)	/	~50	/	/	status report	/
Our camera trap study	2012/2013	T: 0	/	/	localised survey	/

Guestimate = based on little evidence, estimate = based on a field trip without systematic survey, localised survey = systematic survey in a certain area, survey = systematic survey in several parts of the state, status report = information gathered from unknown origin.

T = Tabin, DV = Danum Valley, S = Silabukan, M = Malua, B = Biling, K = Kretam.

Table 9

Stochastic growth rate and extinction risk of 23 population viability analysis scenarios for the historic rhino population in Sabah (means \pm SD), using the computer program VORTEX. Growth rates are specified prior to carrying capacity truncation.

Number	Description	Growth rate	Probability of extinction (%)	Years until first extinction	Population size for extant populations	Genetic diversity (%)
1	Basic	-0.021 ± 0.047	0	50 ± 0	391 ± 186	98
2	Basic + 10% higher no. of males harvested	-0.022 ± 0.055	3 ± 2	64 ± 25	369 ± 191	98
3	Basic + 10% lower no. of males harvested	-0.018 ± 0.036	0	\pm	508 ± 179	83
4	Basic + 10% higher no. of females harvested	-0.024 ± 0.06	0	\pm	270 ± 139	98
5	Basic + 10% lower no. of females harvested	-0.018 ± 0.038	0	\pm	511 ± 193	99
6	Basic + 10% higher reproductive age of females	-0.009 ± 0.026	0	\pm	1415 ± 368	99
7	Basic + 10% lower reproductive age of females	-0.080 ± 0.16	89 ± 3	52 ± 12	41 ± 27	88
8	Basic + 10% higher duration of harvesting	-0.027 ± 0.068	7	54 ± 6	242 ± 135	97
9	Basic + 10% lower duration of harvesting	-0.017 ± 0.036	0	\pm	541 ± 178	99
10	Basic + 10% higher percentage of breeding females	-0.005 ± 0.022	0	\pm	2039 ± 415	100
11	Basic + 10% lower percentage of breeding females	-0.093 ± 0.179	98 ± 1	49 ± 6	12 ± 9	82
12	Basic + 10% higher mortality of juvenile females	-0.030 ± 0.072	8	65 ± 29	212 ± 141	97
13	Basic + 10% lower mortality of juvenile females	-0.016 ± 0.037	0	\pm	618 ± 198	99
14	Basic + 10% higher mortality of juvenile males	-0.021 ± 0.047	0	\pm	382 ± 186	98
15	Basic + 10% lower mortality of juvenile males	-0.021 ± 0.047	0.01	50 ± 0	395 ± 182	98
16	Basic + 10% higher mortality of adult females	-0.056 ± 0.125	49 ± 5	63 ± 24	60 ± 62	88
17	Basic + 10% lower mortality of adult females	-0.010 ± 0.026	0	\pm	1251 ± 306	99
18	Basic + 10% higher mortality of adult males	-0.021 ± 0.048	1 ± 1	50 ± 0	375 ± 176	98
19	Basic + 10% lower mortality of adult males	-0.021 ± 0.047	1 ± 1	50 ± 0	384 ± 166	99
20	Basic + 10% higher perc. of males in breeding pool	-0.021 ± 0.05	1 ± 1	86 ± 0	411 ± 200	98
21	Basic + 10% lower perc. of males in breeding pool	-0.022 ± 0.05	1 ± 1	50 ± 0	369 ± 169	98
22	Basic + 10% higher initial population size	-0.018 ± 0.035	0	\pm	580 ± 192	99
23	Basic + 10% lower initial population size	-0.031 ± 0.083	12 ± 3.2	62 ± 23	216 ± 160	96

that hunting pressure in Sabah was high (Comyn-Platt, 1937; Loch, 1937) and that after virtually exterminating rhinos in Sarawak, the indigenous people of Sarawak crossed the state borders to hunt rhinos in other states or countries of Borneo (Harrisson, 1946). In 1946, rhinos were still thought to exist at a considerable numbers in Sabah, mainly in the north east (Harrisson, 1949). Their numbers appeared to have dropped to very few in 1956 (Harrisson, 1956).

Following the first state wide faunal survey by Davies and Payne (1982), two isolated populations of 13–23 animals at a maximum each were rediscovered in the TWR and Danum Valley together with a few scattered individuals (Davies and Payne, 1982; Andau and Payne, 1982; Boonratana, 1997). Numerous surveys were carried out, especially in the Tabin Wildlife Reserve (Table A3). Resulting population estimates range from three to 20 individuals without an obvious trend in population development. Hunting of rhinos in Sabah continued until recent times, but at a much lower rate. Payne (1980) described four incidences of hunting between 1976 and 1980. The last evidence of rhino poaching was in the year 2001 (New Straits Times, Malaysia April 19, 2001) near the border to Sarawak. Since then no dead rhino has been reported but it cannot be assumed that this equated to zero hunting.

Our resource selection study reveals a clear trend in population development with high number of rhino sightings at the beginning of the study period in the year 2000 and low numbers at the end in 2013. Our camera trap study in the TWR between June 2012 and November 2013 confirms this result. A total of 781.9 km of surveys and a total of 10,316 trapping days (counting only functional camera traps) provided no evidence of Borneo rhino presence in TWR anymore.

Currently, the known population of Sumatran rhinos in Sabah consists of 3 animals in captivity. There is evidence of a few individuals in Kalimantan (WWF, 2013) and no evidence of Sumatran rhinos in Sarawak and Brunei.

3.3. PVA model results

Under scenario (a), the model predicted on average a negative population growth of $r = -0.021 \pm 0.047$ of the historical population of Sabah since the late 19th century with a mean population size of 391 ± 186 individuals after 115 years (Table 9, number 1). The probability of extinction of this population was zero and the mean time to first extinction 50 years. The sensitivity analysis revealed that the percentage of breeding females and the lifetime reproductive period of females – in decreasing order of importance – were the key factors in the population dynamics of Borneo rhinos (Table 10). A decrease in the percentage of breeding females and the lifetime reproductive period of females almost always led to an extinction of the population whereas an increase of both parameters had only little effect on population growth. This suggests a high sensitivity of the population to a reduction in female fertility or reproduction. In comparison to these biological parameters, anthropogenic parameters such as the duration of harvesting or the number of harvested males and females had a less pronounced influence on growth rate. When considering all analysed scenarios (Table 9), the chance of extinction of the

Table 10
Sensitivity indices for changes in parameters of $\pm 10\%$ in the model of the historical rhino population in Sabah.

Parameter	Relative influence on growth rate	
	Parameter +10%	Parameter -10%
Percentage of breeding females	6.35	29.00
Reproductive period of females	5.71	28.10
Total mortality of adult females	6.67	2.10
Total mortality of juvenile females	4.29	2.38
No. of males harvested	0.48	1.43
Duration of harvesting	2.86	1.90
No. of females harvested	1.43	1.43
Percentage of males in breeding pool	0.00	0.48
Total mortality of juvenile males	0.00	0.00
Total mortality of adult males	0.00	0.00
Initial population size	1.07	3.57

historical population was very low, but a decrease in the percentage of breeding females and in the lifetime reproductive period of females severely increased the risk of population extinction.

The model of the population in the TWR without hunting (scenario (b1)) predicted a mean population size of 10 ± 6 individuals after 35 years (Table 11, number 1). The probability of extinction was 20% and the mean time to first extinction 28 ± 6 years. The capture of a male and female for breeding purposes increased the probability of extinction of the population to 31% (Table 11, number 2). Hunting of additional adult males and females (scenario b3) further increased the probability of extinction to 47% and reduced the growth rate of the population to -0.033 ± 0.123 . Similar to the historical population, the sensitivity analysis indicated that percentage of females breeding and lifetime reproductive period of females are the key factors in population dynamics (Table 12), independent of the level of hunting. Additional parameters influencing the population growth rate, but to a much lesser extent, were mortality of adult and juvenile females. The growth rate for such a population with a lower percentage of breeding females and a lower lifetime reproductive period of females combined with moderate hunting activity (scenario b3, Table 11, number 63), resulted in a very high probability of extinction of $75\% \pm 1\%$ after 25 ± 6 years. Management measures preventing poaching and aiming to increase the percentage of breeding females and lifetime reproductive period of females would result in a very low probability of extinction of $9\% \pm 1\%$ and a population size slightly above the size of the initial population of 16 ± 9 (Table 11, number 58).

4. Discussion

We charted the historical development of the Sumatran rhino population in Borneo, and identified the main drivers responsible for its extinction in Sabah and the key demographic parameters which have the strongest influence on Borneo rhino population dynamics. Additionally we documented the resource selection functions of the population in the Tabin Wildlife Reserve and its preferred habitat features.

4.1. Resource selection and preferred habitat features

Our study on habitat use of the rhinos in the Tabin Wildlife Reserve revealed that human disturbance, food availability, access to safe areas and access to other ecological resources are the main factors influencing habitat selection of this population. Our hypotheses that habitat use is only linked to human disturbance or exclusively linked to food availability, terrain and access to other ecological resources were rejected.

A recent study on all remaining Sumatran rhino populations in Sumatra confirms our results, showing that habitat factors and human disturbance are important predictors for Sumatran rhinos, while preferences differed between the different sites (Pusparini et al., 2015).

Six environmental predictors influenced Borneo rhino habitat use in our study area: the soil type, the availability of grassland and dense forest patches within a vicinity of 1 km surrounding a rhino location, elevation, presence of wallows and distance to mud-volcanoes. The importance of the last four factors was already described by Strien (1974, 1986) from his studies in the early 1980s on the basis of opportunistic observations. Our study provides the first statistical quantification of this knowledge within a small scale or second order selection study (Johnson, 1980) on habitat use.

The strongest predictor of rhino presence in our analysis was the year of surveys. Rhino sightings were highest at the beginning of the study period in the year 2000 and lowest at the end of the study period in 2013. This result is being supported by the camera trap study conducted in the year 2012/2013 in the TWR which showed no evidence of rhinos. These results reveal that the rhino population has decreased continuously from 2000 onwards until it became locally extinct. The development of the TWR population therefore followed exactly the predictions of our PVA model for small isolated populations. A small population size combined with a low reproductive performance and an increased mortality rate lead to the extinction of the population in the TWR (see 4.2).

Table 11

Stochastic growth rate and extinction risk of 63 population viability analysis scenarios for the rhino population in the TWR (mean \pm SD). Growth rates are specified prior to carrying capacity truncation.

Number	Description	Growth rate	Probability of extinction (%)	Years until first extinction	Population size for extant populations	Genetic diversity (%)
1	Basic b1 = no harvest	-0.021 ± 0.101	20 ± 1	28 ± 6	10 ± 6	80
2	Basic b2 = one time harvest of ♂ and ♀	-0.025 ± 0.109	31 ± 1	29 ± 5	10 ± 6	80
3	Basic b3 = harvesting of ♂ and ♀ every 10 years	-0.033 ± 0.123	47 ± 1	27 ± 5	9 ± 5	78
Impact of reproductive age of females						
4	Basic b1 + 10% higher rep. age of females	-0.013 ± 0.097	12 ± 1	28 ± 6	13 ± 7	82
5	Basic b2 + 10% higher rep. age of females	-0.020 ± 0.107	23 ± 1	30 ± 5	9 ± 7	81
6	Basic b3 + 10% higher rep. age of females	-0.027 ± 0.120	37 ± 2	27 ± 6	10 ± 6	79
7	Basic b1 + 10% lower rep. age of females	-0.028 ± 0.105	28 ± 1	29 ± 5	8 ± 5	78
8	Basic b2 + 10% lower rep. age of females	-0.033 ± 0.113	43 ± 2	29 ± 4	8 ± 5	78
9	Basic b3 + 10% lower rep. age of females	-0.039 ± 0.129	58 ± 2	26 ± 6	8 ± 5	78
Impact of percentage of breeding females						
10	Basic b1 + 10% higher perc. of breeding females	-0.011 ± 0.097	10 ± 1	28 ± 5	13 ± 8	82
11	Basic b2 + 10% higher perc. of breeding females	-0.017 ± 0.103	21 ± 1	30 ± 4	12 ± 7	82
12	Basic b3 + 10% higher perc. of breeding females	-0.026 ± 0.118	36 ± 2	27 ± 6	10 ± 7	80
13	Basic b1 + 10% lower perc. of breeding females	-0.030 ± 0.108	30 ± 1	28 ± 5	8 ± 5	78
14	Basic b2 + 10% lower perc. of breeding females	-0.033 ± 0.114	41 ± 5	30 ± 4	7 ± 5	77
15	Basic b3 + 10% lower perc. of breeding females	-0.041 ± 0.128	58 ± 2	26 ± 6	7 ± 4	76
Impact of mortality of juvenile females						
16	Basic b1 + 10% higher mort. of juv. females	-0.022 ± 0.102	19 ± 1	28 ± 6	10 ± 5	80
17	Basic b2 + 10% higher mort. of juv. females	-0.026 ± 0.110	23 ± 1	30 ± 5	11 ± 7	79
18	Basic b3 + 10% higher mort. of juv. females	-0.038 ± 0.135	61 ± 2	25 ± 6	7 ± 4	75
19	Basic b1 + 10% lower mort. of juv. females	-0.020 ± 0.10	18 ± 1	29 ± 5	11 ± 6	80
20	Basic b2 + 10% lower mort. of juv. females	-0.025 ± 0.109	29 ± 5	29 ± 5	7 ± 6	82
21	Basic b3 + 10% lower mort. of juv. females	-0.032 ± 0.132	45 ± 2	26 ± 6	9 ± 5	78
Impact of mortality of juvenile males						
22	Basic b1 + 10% higher mort. of juvenile males	-0.020 ± 0.101	20 ± 1	29 ± 5	10 ± 6	80
23	Basic b2 + 10% higher mort. of juvenile males	-0.026 ± 0.106	41 ± 5	30 ± 5	6 ± 5	78
24	Basic b3 + 10% higher mort. of juvenile males	-0.033 ± 0.125	48 ± 2	26 ± 6	9 ± 5	79
25	Basic b1 + 10% lower mort. of juvenile males	-0.021 ± 0.102	18 ± 1	29 ± 5	10 ± 6	80
26	Basic b2 + 10% lower mort. of juvenile males	-0.024 ± 0.108	32 ± 5	29 ± 5	10 ± 5	80
27	Basic b3 + 10% lower mort. of juvenile males	-0.033 ± 0.125	46 ± 2	26 ± 5	8 ± 5	78
Impact of mortality of adult females						
28	Basic b1 + 10% higher mort. of adult females	-0.025 ± 0.106	26 ± 1	28 ± 6	9 ± 6	80
29	Basic b2 + 10% higher mort. of adult females	-0.033 ± 0.114	47 ± 5	28 ± 6	5 ± 5	79
30	Basic b3 + 10% higher mort. of adult females	-0.039 ± 0.130	55 ± 2	26 ± 6	8 ± 5	77
31	Basic b1 + 10% lower mort. of adult females	-0.015 ± 0.096	14 ± 1	29 ± 5	12 ± 7	81
32	Basic b2 + 10% lower mort. of adult females	-0.020 ± 0.100	31 ± 5	30 ± 3	9 ± 8	82
33	Basic b3 + 10% lower mort. of adult females	-0.028 ± 0.119	39 ± 2	27 ± 6	10 ± 6	79
Impact of mortality of adult males						
34	Basic b1 + 10% higher mort. of adult males	-0.022 ± 0.105	22 ± 1	28 ± 5	10 ± 6	79
35	Basic b2 + 10% higher mort. of adult males	-0.024 ± 0.110	34 ± 5	29 ± 6	7 ± 6	79
36	Basic b3 + 10% higher mort. of adult males	-0.034 ± 0.124	51 ± 2	26 ± 6	8 ± 5	77
37	Basic b1 + 10% lower mort. of adult males	-0.019 ± 0.098	15 ± 1	28 ± 5	11 ± 6	81
38	Basic b2 + 10% lower mort. of adult males	-0.022 ± 0.106	27 ± 4	30 ± 5	10 ± 5	81
39	Basic b3 + 10% lower mort. of adult males	-0.032 ± 0.123	43 ± 2	27 ± 5	8 ± 5	78
Impact of percentage of males in breeding pool						
40	Basic b1 + 10% higher perc. of males in br. pool	-0.021 ± 0.102	20 ± 1	29 ± 5	10 ± 6	80
41	Basic b2 + 10% higher perc. of males in br. pool	-0.024 ± 0.107	37 ± 5	30 ± 4	11 ± 6	82
42	Basic b3 + 10% higher perc. of males in br. pool	-0.033 ± 0.124	49 ± 2	27 ± 6	9 ± 5	79
43	Basic b1 + 10% lower perc. of males in br. pool	-0.021 ± 0.102	20 ± 1	28 ± 5	10 ± 6	79
44	Basic b2 + 10% lower perc. of males in br. pool	-0.027 ± 0.114	34 ± 5	30 ± 4	10 ± 6	80
45	Basic b3 + 10% lower perc. of males in br. pool	-0.034 ± 0.124	47 ± 2	26 ± 5	8 ± 5	78
Impact of initial population size						
46	Basic b1 + 10% higher initial pop. size	-0.018 ± 0.095	13 ± 1	29 ± 5	12 ± 7	82
47	Basic b2 + 10% higher initial pop. size	-0.023 ± 0.102	20 ± 4	30 ± 3	9 ± 8	82
48	Basic b3 + 10% higher initial pop. size	-0.031 ± 0.118	37 ± 2	27 ± 5	10 ± 6	80
49	Basic b1 + 10% lower initial pop. size	-0.021 ± 0.108	25 ± 1	28 ± 5	9 ± 6	78
50	Basic b2 + 10% lower initial pop. size	-0.030 ± 0.116	54 ± 5	29 ± 5	5 ± 5	79
51	Basic b3 + 10% lower initial pop. size	-0.035 ± 0.132	61 ± 2	25 ± 6	7 ± 5	75

(continued on next page)

Table 11 (continued)

Number	Description	Growth rate	Probability of extinction (%)	Years until first extinction	Population size for extant populations	Genetic diversity (%)
Impact of harvest interval						
52	Basic b3 + 10% higher harvest int.	-0.033 ± 0.122	49 ± 2	27 ± 6	8 ± 5	79
53	Basic b3 + 10% lower harvest int.	-0.034 ± 0.123	47 ± 2	25 ± 5	8 ± 5	78
Impact of number of harvested females						
54	Basic b3 + 10% higher har. females	-0.040 ± 0.141	63 ± 2	25 ± 6	7 ± 4	77
55	Basic b3 + 10% lower har females	-0.026 ± 0.109	35 ± 2	26 ± 6	10 ± 6	80
Impact of number of harvested males						
56	Basic b3 + 10% higher har. Males	-0.037 ± 0.138	61 ± 2	25 ± 6	8 ± 5	77
57	Basic b3 + 10% lower har. males	-0.031 ± 0.111	41 ± 2	27 ± 5	9 ± 5	79
Impact of reproductive age and perc of breeding females						
58	4 + 10% higher percentage of breeding females	-0.006 ± 0.094	9 ± 1	28 ± 6	16 ± 9	83
59	5 + 10% higher percentage of breeding females	-0.011 ± 0.100	13 ± 1	30 ± 5	14 ± 8	82
60	6 + 10% higher percentage of breeding females	-0.019 ± 0.113	26 ± 1	27 ± 6	12 ± 8	80
61	7 + 10% lower percentage of breeding females	-0.035 ± 0.109	39 ± 2	28 ± 6	7 ± 4	77
62	8 + 10% lower percentage of breeding females	-0.039 ± 0.117	58 ± 2	29 ± 5	7 ± 4	77
63	9 + 10% lower percentage of breeding females	-0.05 ± 0.131	75 ± 1	25 ± 6	6 ± 3	75

Table 12

Relative influence on growth rate for changes in parameters of $\pm 10\%$ for the rhino population in the TWR with different hunting scenarios.

Parameter	Scenario b1		Scenario b2		Scenario b3	
	Sensitivity indices		Sensitivity indices		Sensitivity indices	
	+10%	-10%	+10%	-10%	+10%	-10%
Reproductive age of females	3.81	3.33	2.00	3.20	1.82	1.82
Percentage of females breeding	3.97	3.57	2.67	2.67	1.77	2.02
Total mortality of juvenile females	0.48	0.48	0.40	0.00	1.52	0.3
Total mortality of juvenile males	0.48	0.00	0.40	0.40	0.00	0.00
Total mortality of adult females	0.76	1.14	1.28	0.80	0.73	0.61
Total mortality of adult males	0.24	0.48	0.20	0.60	0.15	0.15
Percentage of males in breeding pool	0.00	0.30	0.40	0.80	0.00	0.00
Initial population size	1.07	0.00	0.60	1.50	0.45	0.45
Females harvested	/	/	/	/	0.21	0.21
Males harvested	/	/	/	/	0.12	0.06
Harvest interval	/	/	/	/	0.00	0.30

To our surprise this decline has not been detected by the regular surveys conducted on a yearly and even monthly basis within the reserve. The lack of a rigid scientific sampling scheme as well as the shortage of information about the effort expended by the field teams in terms of search routes, distance covered and time spent searching are part of the problem. Repeated information about presence/absence data and individual identification, as required by an occupancy modelling framework (MacKenzie and Royle, 2005), is not available. The absence of such data is a regular problem in conservation studies, pointing towards serious shortcomings in the design or execution of management-oriented or conservation-oriented surveys (Karanth et al., 2003; Pusparini and Wibisono, 2013). Large amounts of money were spent with very few results, often leading to poor conservation practices (Karanth et al., 2003). We therefore highly recommend for all future studies on Sumatran rhinos as well as other endangered animals to conduct regular surveys in an appropriate season and in a habitat which is representative for the whole population. The survey team need to carefully record sampling effort and georeference the area. Decisions based on little evidence can be harmful to wildlife, habitat and the survival of both (Huettmann, 2005).

We studied resource selection and habitat use within the Tabin wildlife reserve. The study represents the only existing data on habitat use of the Borneo rhino. Our study utilised data collected during 277 surveys conducted over a study period of 13 years. It thus covers a long time period and includes data collected during both wet and dry seasons, providing a comprehensive summary of the range of conditions over which rhinos preferentially occur in time and space (Wisiz et al., 2008). Our study area is comparably small and it has been modified by human activity. The assessments therefore need to be treated cautiously and we cannot assume that it reflects the preferences of rhinos in undisturbed situations. But such situations hardly exist for Sumatran rhinos anymore. In Sabah, less than 51% of the land area is covered with forest, 32% of these forests have been logged several times leaving extensive areas in a highly damaged condition and only 1% of undisturbed lowland forest remained (Reynolds et al., 2011). A similar situation exists in Kalimantan (Gaveau et al., 2014). Our resource selection study thus describes habitat preferences of a species that lived for more than 30 years in a human altered landscape in one of the few remaining lowland forests of Sabah. We applied a use-vs-availability design to identify relevant habitat covariates within this set of given choices. The exact number of rhinos included in this study is not known. The data on surveys conducted in the TWR (Boonratana, 1997: estimated 3–9 rhinos in 1997 and Thayaparan, 2008: 5 rhinos in 2005) in combination with the results of our PVA analysis for the TWR population suggest that there was a population

of at least 5–9 individuals at the beginning of our study period. This assessment is consistent with a rough estimate of the number of rhinos which emerged when we grouped GPS locations of rhinos within a certain year. Here we assume that the maximum size of a male home range was 60 km² (Strien, 1986) and that home ranges did not overlap.

We studied resource selection and habitat use within a relatively homogenous forest reserve with comparably little variation in habitat factors. The high resolution satellite maps (5 m) used for the vegetation interpretation provided us with the best possible data currently available to detect small scale differences in vegetation (Niedballa et al., 2015). However, satellite data will not provide information on variation in the undergrowth of a forest where the main food of Sumatran rhinos is located. Vegetation assessments on the ground are therefore essential to complement vegetation assessments based on remote-sensing data. The data used in our study were not evenly spread across the whole study area and some areas were more frequently surveyed than others. Such a spatial bias can lead to an 'environmental bias' in the assessment of resource selection functions because of the overrepresentation of frequently represented environmental features (Kramer-Schadt et al., 2013). We addressed this problem by reducing the number of occurrence records in oversampled regions using spatial filtering (Dormann et al., 2007; Veloz, 2009; Northrup et al., 2013). This method has the apparent disadvantage that highly preferred habitats, which may be of great importance for the Sabah rhinos, will be downgraded in their apparent importance (Boyce et al., 2002). Such downgrading is less likely to happen if spatial filtering is applied for only a limited time frame. The covariates identified with such a filtering approach have thus been identified with a conservative method.

4.2. Population viability analysis

The historical population model predicted a low chance of extinction of the rhino population in Sabah after 115 years, despite a yearly harvesting quota of 120 animals between 1930 and 1950. The model indicates that population growth was mainly influenced by the percentage of breeding females and the lifetime reproductive period of females. A small decrease of 10% in these parameters resulted already in the near extinction of the entire rhino population.

Recent findings suggest that reproduction of the remnant Sumatran rhinos is being constrained by reproductive pathologies in females. In Sabah, two adult females caught in the past four years had large tumours in their reproductive tract which made it impossible for them to successfully conceive and deliver a young (TB Hildebrandt et al., unpublished data), and in peninsular Malaysia, 6 out of 9 rhinos captured in the 1980s had reproductive tract pathologies (Schaffer et al., 1994; Ahmad et al., 2013). In most cases, the tumours were present at time of capture and did not develop in captivity. Reproductive tract tumours in rhinos develop after long non-reproductive periods and prolonged exposure to sex steroids (Hermes et al., 2004, 2006, 2014), which eventually results in a shortening of the lifetime reproductive period of the females. Such non-reproductive periods can occur in small remnant populations of low population size and in areas of low population densities where mating partners have difficulties to find each other.

The hunting of rhinos most probably contributed towards such a low population density, and logging as well as agricultural and other human activities have likely fostered this trend. The first large-scale deforestation activities in Borneo started with the colonialisation. At that time forests on flat terrain were logged manually to meet the local and global demand for rice, rubber, palm oil and coconuts (Davies and Payne, 1982; Langner, 2009). An increasing demand for timber in the mid 1950s set off the beginning of commercial logging (Davies and Payne, 1982). From then on logging increased at an accelerating rate. From 1973 until 2010 around 30% of the primary forest area in Borneo was converted into oil palm plantations and agricultural land, with the highest losses of rainforest recorded in Sabah (40%) and Kalimantan (31%) (Gaveau et al., 2014).

The disturbance caused by logging and its associated human activities had a strong impact on the Sumatran rhinos. Individuals were scattered (Flynn, 1981) and rhino populations were isolated in patches of rainforest surrounded by agricultural land (Davies and Payne, 1982). One of these populations occurred in the TWR, the other in Danum Valley, Sabah. The PVA model of the TWR population showed that such small and isolated populations have a negative growth rate, even in cases of zero hunting and a moderate (zero hunting) to high (moderate level of hunting) probability of extinction within 30 years. The model indicated that the combined efforts of total protection and an increase in the reproductive performance of females could have saved the Tabin population. Capturing of individuals in areas of low population densities and assembling them into suitable areas with high levels of protection could have been such a possible measure.

The PVA model of the TWR population also indicates that a species with such a low reproductive output and with proven reproductive health problems needs further measures to guarantee its long term persistence. These measures should aim to increase the chance of successful mating and conception and shortening the intercalving interval. Assisted reproduction techniques could be such possible measures.

The predictions made by the PVA model need to be interpreted cautiously because of the limited data available (Ellner et al., 2002; Brook et al., 2002). As some of the crucial biological information was unavailable, we chose conservative estimates for the parameters in order to ensure that any factor likely to emerge as important would do so despite these conservative assumptions.

4.3. Conclusions

Small isolated populations of Sumatran rhinos are deemed to go extinct unless measures designed to prevent hunting, increase fertility, enhance the chance of encounters of potential mating partners in the wild or shorten the intercalving

interval are undertaken. The predominant conservation practice of governments and NGOs to avoid interfering with free-ranging populations and to rely on natural breeding will not solve the problem. A scientifically rigorous surveying regime ought to be applied in order to identify small populations, monitor their development and forecast any potentially negative population developments. The experiences with the rhinos in Sabah show that a remnant population with less than 15 individuals and with its inherent reproductive specifics have a high chance of extinction if left to their own devices. Our PVA analysis suggests that a combination of zero hunting and improvements in reproductive performance of females will make a crucial difference to the viability of such a remnant population. Improving reproductive performance usually requires comprehensive checks on the reproductive health of all members of the population and intensive efforts to maximise fertility and successful encounters with potential mating partners. This will require an intensive conservation management, which may require rhinos which occur in areas with high human encroachment as well as in areas with low nutrient and mineral concentration to be translocated to more suitable areas to provide optimal conditions for them to breed. It may also include the capture and reproductive management of key individuals within a confined space where the animals are easily accessible and where the monitoring of their health, fertility and pregnancies can be ensured. Such efforts will be too late for the Borneo rhinos in Sabah, but the Sumatran subspecies can still be saved. The success of the coordinated captive breeding program of the Iberian lynx (*Lynx pardinus*) (Vargas et al., 2009), the most endangered felid worldwide, is an example which demonstrates how this could be done.

Acknowledgements

The authors thank the Sabah Wildlife Department and the Forestry Department for the permission to conduct their research activities in the TWR and their local partner BORA for the comprehensive support. Financial support was provided by the Federal Ministry of Education and Research (BMBF) FKZ 033L045, Zoo Leipzig and SOS Rhino. We thank the anonymous reviewers whose suggestions helped improve this manuscript. The publication of this article was funded by the Open Access fund of the Leibniz Association.

Appendix A. Supplementary data

Supplementary material related to this article can be found online at <http://dx.doi.org/10.1016/j.gecco.2016.02.006>.

References

- Abram, N.K., Meijaard, E., Wells, J.A., Ancrenaz, M., Pellicier, A.-S., Runting, R.K., Gaveau, D., Wich, S., Nardiyono, Tjiu, A., Nurcahyo, A., Mengersen, K., 2015. Mapping perceptions of species' threats and population trends to inform conservation efforts: the Bornean orangutan case study. *Divers. Distrib.* 21, 487–499.
- Ahmad, A.H., Payne, J., Zainuddin, Z.Z., 2013. Preventing the extinction of the Sumatran rhinoceros. *J. Indon. Nat. Hist.* 1, 11–22.
- Andau, P.M., 1987. Conservation of the Sumatran rhinoceros in Sabah, Malaysia. *Rimba Indones.* 21, 39–47.
- Andau, P.M., Payne, J., 1982. The plight of the Sumatran rhinoceros in Sabah. Report Presented at the 8th Malaysia Forestry Conference, Sandakan-Sabah, pp. 1–13.
- Anderson, J.A.R., 1968. Conservation in the state of Sarawak, Malaysia. In: Talbot, L.M., Talbot, M.H. (Eds.), *Conservation in Tropical South East Asia*. Gland, Vol. 10. IUCN Publications, NS, pp. 498–502.
- Anonymous, 1890. *Handbook of British North Borneo*. William Clowes & Sons, London.
- Austin, M.P., 2002. Spatial prediction of species distribution: an interface between ecological theory and statistical modelling. *Ecol. Model.* 157, 101–118.
- Banks, E., 1931. A popular account of the mammals of Borneo. *J. Malays. Branch R. Asiatic Soc.* 9, 1–139.
- Banks, E., 1935. A collection of montane mammals and birds from Mulu in Sarawak. *Sarawak Museum J.* 4, 327–341.
- Banks, E., 1937. Rum and rhinos. *Sarawak Gazette* 163–164. 1937 August 2.
- Banks, E., 1949. *Bornean Mammals*. Kuching Press, Kuching, pp. 1–83.
- Bartlett, E., 1891. Notes on the Bornean rhinoceros. *Proc. Zool. Soc. Lond.* 654–655.
- Beissinger, S.R., McCullough, D.R., 2002. *Population Viability Analysis*. The University of Chicago Press, Chicago, USA.
- Bennett, A.F., 2003. *Linkages in the Landscape. The Role of Corridors and Connectivity in Wildlife Conservation*, second ed. IUCN, Gland, Switzerland, Cambridge, UK.
- Bland, L.M., Collen, B., Orme, C.D.L., Bielby, J., 2015. Predicting the conservation status of data-deficient species. *Conserv. Biol.* 29, 250–259.
- Bock, C., 1881. The head-hunters of Borneo: a narrative of travel up to Mahakkam and down the Barito; also, journeying in Sumatra. London, Sampson Low, Marston, Searle and Rivington, pp. 1–344.
- Boonratana, R., 1997. A state-wide survey to estimate the distribution and density of the Sumatran rhinoceros, Asian elephant and banteng in Sabah, Malaysia. Report to Wildlife Conservation Society, New York, pp. 1–76.
- Boyce, M.S., Vernier, P.R., Nielsen, S.E., Schmiegelow, F.K.A., 2002. Evaluating resource selection functions. *Ecol. Model.* 157, 281–300.
- Brook, B.W., Burgman, M.A., Akçakaya, H.R., O'Grady, J.J., Frankham, R., 2002. Critiques of PVA ask the wrong questions: Throwing the heuristic baby out with the numerical bath water. *Conserv. Biol.* 16, 262–263.
- Brunton, R., 1963. The rhinos of South-East Asia. *Conserv. News (August)*, 5–10.
- Burgess, P.F., 1961. Wildlife conservation in North Borneo. In: Wyatt-Smith, J., Wycherley, P.R. (Eds.), *Nature Conservation in Western Malaysia*. Nature Soc., pp. 143–151. Special issue of the Malay. Nature J. Kuala Lumpur, Malay. 1–260.
- Burnham, K.P., Anderson, C., 2002. *Model Selection and Multi-model Inference: A Practical Information-Theoretic Approach*. Springer-Verlag, New York.
- Carson, G.L., 1968. Conservation in Sabah, Malaysia. In: Talbot, L.M., Talbot, M.H. (Eds.), *Conservation in Tropical South East Asia*. Gland. IUCN Publications, NS, pp. 492–497. 10.
- Chander, G., Markham, B.L., Helder, D.L., 2009. Summary of current radiometric calibration coefficients for Landsat MSS, TM, ETM+ and EO-1 ALI sensors. *Remote Sens. Environ.* 113, 893–903.
- Chavez Jr., P.S., 1996. Image-based atmospheric corrections—revisited and revised. *Photogramm. Eng. Remote Sens.* 62, 1025–1036.
- Clements, R., Rayan, D.M., Zafir, A.W.A., Venkataraman, A., Alfred, R., Payne, J., Ambu, L., Sharma, D.S.K., 2010. Trio under threat: can we secure the future of rhinos, elephants and tigers in Malaysia. *Biodivers. Conserv.* 19, 1115–1136.
- Comyn-Platt, T., 1937. A report on fauna preservation in British North Borneo. *J. Soc. Preserv. Fauna Emp.* 30, 52–55.

- Crawley, M.J., 2007. *Statistics. An Introduction Using R*. John Wiley & Sons, Ltd., West Sussex, England, pp. 1–327.
- Davies, G., Payne, J., 1982. A faunal survey of Sabah. Report. Forest Dept., Sabah, pp. 1–294.
- Dawson, S., 1993. Estimating elephant numbers in Tabin Wildlife Reserve, Sabah, Malaysia Gajah 11, 16–28.
- Dormann, C.F., McPherson, J.M., Araújo, M.B., Bivand, R., Bolliger, J., Carl, G., Davies, R.G., Hirzel, A., Jetz, W., Daniel Kissling, W., Kühn, I., Ohlemüller, R., Peres-Neto, P.R., Reineking, B., Schröder, B., Schurr, F.M., Wilson, R., 2007. Methods to account for spatial autocorrelation in the analysis of species distributional data: a review. *Ecography* 30, 609–628.
- Ellner, S., Fieberg, J., Ludwig, D., Wilcox, C., 2002. Precision of population viability analysis. *Conserv. Biol.* 16, 258–261.
- Encyclopaedia of the Nations, 2015. <http://www.nationsencyclopedia.com/Asia-and-Oceania/Malaysia-LOCATION-SIZE-AND-EXTENT.html>.
- Ferrier, S., Drielsma, M., Manion, G., Watson, G., 2002. Extended statistical approaches to modelling spatial pattern in biodiversity in northeast New South Wales. II. Community-level modelling. *Biodivers. Conserv.* 11, 2309–2338.
- Flynn, R.W., 1981. Operation rhino Sabah, phase I. Report WWF Malaysia. pp. 1–14.
- Funk, V.A., Richardson, K.S., 2002. Systematic data in biodiversity studies: Use it or lose it. *Syst. Biodivers.* 51, 303–316.
- Gaveau, D.L.A., Sloan, S., Molidena, E., Yaen, H., Sheil, D., Abram, N.K., Ancrenaz, M., Nasi, R., Quinones, M., Wielaaard, N., Meijaard, E., 2014. Four decades of forest persistence, clearance and logging on Borneo. *PLoS One* 9, e101654. <http://dx.doi.org/10.1371/journal.pone.0101654>.
- Gatti, A., Brito, D., Mendes, S.L., 2011. How many lowland tapirs (*Tapirus terrestris*) are needed in Atlantic forest fragments to ensure long-term persistence? *Stud. Neotrop. Fauna Environ.* 46, 77–84.
- Griffith, B., Scott, J.M., Carpenter, J.W., Reed, C., 1989. Translocation as a species conservation tool: status and strategy. *Science* 4, 477–480.
- Guisan, A., Zimmermann, N.E., 2000. Habitat distribution models in ecology. *Ecol. Model.* 135, 147–186.
- Harris, G., Thirgood, S., Hopcraft, J.G.C., Cromsigt, J.P.G.M., Berger, J., 2009. Global decline in aggregated migrations of large terrestrial mammals. *Endanger. Species Res.* 7, 55–76.
- Harrisson, T., 1946. Pygmy rhinoceros in Borneo. *Times (London)* (September), 5.
- Harrisson, T., 1948. The large mammals of Borneo. *Sarawak Gazette* 1948 Dec. 1: 254–261.
- Harrisson, T., 1949. The large mammals of Borneo. *Malayan Nat. J.* 4, 70–76.
- Harrisson, T., 1955. Borneo fauna anxieties. *Oryx* 3, 134–137.
- Harrisson, T., 1956. Rhinoceros in Borneo: and traded to China. *Sarawak Museum J.* 7, 263–274.
- Harrisson, T., 1961. The threat to rare animals in Borneo. *Oryx* 6, 126–128.
- Harrisson, T., 1965. A future for Borneo's wildlife? *Oryx* 8, 99–104.
- Harrisson, T., 1975. The rhinoceros- and mammal extinction in general. *Borneo Res. Bull.* 7, 71–72.
- Hermes, R., Göritz, F., Saragusty, J., Hildebrandt, T.B., 2014. Reproductive tract tumours: the scourge of woman reproduction ails Indian rhinoceroses. *PLoS One* <http://dx.doi.org/10.1371/journal.pone.0092595>.
- Hermes, R., Hildebrandt, T., Göritz, F., 2004. Reproductive problems directly attributable to long-term captivity-asymmetric reproductive ageing. *Anim. Reprod. Sci.* 82–83, 49–60.
- Hermes, R., Hildebrandt, T.B., Walzer, C., Göritz, F., Patton, M.L., Silinski, S., Anderson, M.J., Reid, C.E., Wibbelt, G., Tomasova, K., Schwarzenberger, F., 2006. The effect of long non-reproductive periods on the genital health in captive female white rhinoceroses (*Ceratotherium simum simum*, *C.s. cottoni*). *Theriogenology* 65, 1492–1515.
- Heynsius Viruly, A., Heurn, F.C.van, 1935. Overzicht van de uit Nederlandsch-Indie ontvangen gegevens, met biologische aantekeningen omtrent de betreffende diersoorten. *Meded. Ned. Commissie voor Int. Natuurbescherming* 10 (Supplement), 25–77. pls. 1–4.
- Huettmann, F., 2005. Databases and science-based management in the context of wildlife and habitat: toward a certified ISO standard for objective decision-making for the global community by using the Internet. *J. Wildl. Manage.* 69, 466–472.
- International Union for Conservation of Nature (IUCN) 2015. Sumatran rhino likely go extinct unless action is taken urgently, warns IUCN. http://www.iucn.org/news_homepage/?21904/Sumatran-Rhino-likely-to-go-extinct-unless-action-is-taken-urgently-warns-IUCN.
- Jentink, F.A., 1897. Zoological results of the Dutch Scientific Expedition to Central Borneo: The mammals. *Notes Leyden Mus.* 19, 29–66.
- Johnson, D.H., 1980. The comparison of usage and availability measurements for evaluating resource preference. *Ecology* 61, 65–71.
- Johnson, C.J., Nielsen, S.E., Merrill, E.H., McDougal, T.L., Boyce, M.S., 2006. Resource selection functions based on use-availability data: theoretical motivation and evaluation methods. *J. Wildl. Manage.* 70, 347–357.
- Karant, K.U., Nichols, J.D., Seidenstricker, J., Dinerstein, E., Smith, J.L.D., McDougal, C., Johnsingh, A.J.T., Chundawat, R.S., Thapar, V., 2003. Science deficiency in conservation practice: the monitoring of tiger populations in India. *Anim. Conserv.* 6, 141–146.
- Khan, M., 1989. Asian Rhinos: An Action Plan for their Conservation. IUCN, Gland, pp. 1–23.
- Khan, M., 1993. Asian Rhino specialist group. *Pachyderm* 18, 3–8.
- Khan, M., Foote, T.J., Strien, N.J.van, 1999. Asian Rhino Specialist Group (AsRSG). *Pachyderm* 27, 16–21.
- Kramer-Schadt, S., Niedballa, J., Pilgrim, J.D., Schröder, B., Lindenborn, J., Reinfelder, V., Stillfried, M., Heckmann, I., Scharf, A.K., Augeri, D.M., Cheyne, S.M., Hearn, A.J., Ross, J., Macdonald, D.W., Mathai, J., Eaton, J., Marshall, A.J., Semiadi, G., Rustam, R., Bernard, H., Alfred, R., Samejima, H., Duckworth, J.W., Breitenmoser-Wuersten, C., Belant, J.L., Hofer, H., Wilting, A., 2013. The importance of correcting for sampling bias in MaxEnt species distribution models. *Divers. Distrib.* 19, 1366–1379.
- Lacy, R.C., Pollak, J.P., 2014. *Vortex: A Stochastic Simulation of the Extinction Process. Version 10.0*. Chicago Zoological Society, Brookfield, Illinois, USA.
- Langner, A.J., 2009. Monitoring tropical forest degradation and deforestation in Borneo, Southeast Asia. *Diss. LMU Munich*, pp. 1–179.
- Langner, A., Samejima, H., Ong, R.C., Titin, J., Kitayama, K., 2012. Integration of carbon conservation into sustainable forest management using high resolution satellite imagery: A case study in Sabah, Malaysian Borneo. *Int. J. Appl. Earth Obs. Geoinf.* 18, 305–312.
- Leblond, M., Dussault, C., St-Laurent, M.-H., 2014. Development and validation of an expert-based habitat suitability model to support boreal caribou conservation. *Biol. Cons.* 177, 100–108.
- Loch, C.W., 1937. Rhinoceros sondaicus: the Javan or lesser one-horned rhinoceros and its geographical distribution. *J. Malays. Branch R. Asiatic Soc.* 15, 130–149.
- Mace, G.M., Lande, R., 1991. Assessing Extinction Threats: Toward a reevaluation of IUCN threatened species categories. *Conserv. Biol.* 5, 148–157.
- MacKenzie, D.I., Royle, J.A., 2005. Designing occupancy studies: general advice and allocating survey effort. *J. Appl. Ecol.* 42, 1105–1114.
- MacKinnon, J., 1970. Stop the rhino horn merchants. *Oryx* 10, 347–348.
- Martin, E.B., 1989. The rhino product trade in Northern and Western Borneo. *Pachyderm* 12, 38–41.
- Meek, M.H., Wells, C., Tomalty, K.M., Ashander, J., Cole, E.M., Gille, D.A., Putman, B.J., Rose, J.P., Savoca, M.S., Yamane, L., Hull, J.M., Rogers, D.L., Rosenblum, E.B., Shogren, J.F., Swaisgood, R.R., May, B., 2015. Fear of failure in conservation: The problem and potential solutions to aid conservation of extremely small populations. *Biol. Cons.* 184, 209–217.
- Meijaard, E., 1996. The Sumatran rhinoceros in Kalimantan, Indonesia: its possible distribution and conservation prospects. *Pachyderm* 21, 15–23.
- Melbourne, B.A., Hastings, A., 2008. Extinction risk depends strongly on factors contributing to stochasticity. *Nature* 454, 100–103.
- Mjoeberg, E., 1929. Durch die Insel der Kopfjaeger: Abenteuer im Innern von Borneo. Leipzig, F.A. Brockhaus, pp. 1–332.
- Nardelli, F., 2014. The last chance for the Sumatran rhinoceros? *Pachyderm* 55, 43–53.
- Niedballa, J., Sollmann, R., bin Mohamed, A., Bender, J., Wilting, A., 2015. Defining habitat covariates in camera-trap based occupancy studies. *Sci. Rep.* 5, <http://dx.doi.org/10.1038/srep17041>.
- Nor, S.M., Ambu, L.N., Tuuga, A., 1989. Rhino survey in Tabin Wildlife Reserve. In: Lahad Datu. Sabah Mus. Monog., vol. 3. pp. 77–82.
- Northrup, J.M., Hooten, M.B., Anderson Jr., C.R., Wittemyer, G., 2013. Practical guidance on characterizing availability in resource selection functions under a use-availability design. *Ecology* 94, 1456–1463.
- O'Grady, J.J., Brook, B.W., Reed, D.H., Ballou, J.D., Tonkyn, D.W., Frankham, R., 2006. Realistic levels of inbreeding depression strongly affect extinction risk in wild populations. *Biol. Cons.* 133, 42–51.
- Owen-Smith, R.N., 1988. *Megaherbivores. The Influence of Very Large Body Size on Ecology*. Cambridge University Press, 369.

- Payne, J., 1980. Report on rhinoceros (*Dicerorhinus sumatrensis*) in Silabukan Forest Reserve, Sabah: a recommendation to catch and translocate them. Draft report. published by Borneo Rhino Alliance.
- Payne, J., 1986. Development of conservation areas in Sabah, North Borneo. WWF Yb. 1985–1986, pp. 482–484.
- Payne, J., 1990. The distribution and status of the Asian two-horned rhinoceros (*Dicerorhinus sumatrensis harrissoni*) in Sabah, Malaysia. Kuala Lumpur, WWF Malay. Rep. pp. 1–33.
- Pryer, W.B., 1881. Animal life in Borneo. Zoologist 5, 393–398.
- Pulliam, H.R., Dunning, J.B., Liu, L., 1992. Population dynamics in complex landscapes: a case study. Ecol. Appl. 2, 165–177;
- Pusparini, W., Wahyudi, T., Ritonga, H., Rumapea, E., 2013. Re-estimating Sumatran rhinoceros population in Gunung Leuser National Park, Sumatran, Indonesia. Report. pp. 1–35.
- Pusparini, W., Sievert, P.R., Fuller, T.K., Randhier, T.O., Andayani, N., 2015. Rhinos in the Parks: An Island-wide survey of the last wild population of the sumatran rhinoceros. PLoS One 10, <http://dx.doi.org/10.1371/journal.pone.0136643>.
- Pusparini, W., Wibisono, H.T., 2013. Landscape-level assessment of the distribution of the Sumatran rhinoceros in Bukit Barisan Selatan National Park. Sumatra. Pachyd. 53, 59–65.
- R Development Core Team, 2014. R: A language and environment for statistical computing. Vienna, Austria. R Foundation for Statistical Computing. Retrieved from <http://R-project.org>.
- Rabinowitz, A., 1992. Rhino survey in the Greater Danum Valley conservation area. pp. 1–54. 14–28 September 1992. Rep.
- Rayan, D.M., Linkie, M., 2015. Conserving tigers in Malaysia: A science-driven approach for eliciting conservation policy change. Biol. Cons. 184, 18–26.
- Reynolds, G., Payne, J., Sinun, W., Mosigil, G., Walsh, R.P.D., 2011. Changes in forest land use and management in Sabah, Malaysian Borneo, 1990–2010, with a focus on the Danum Valley region. Philos. Trans. R. Soc. Lond. B Biol. Sci. 366, 3168–3176.
- Ridley, H.N., 1916. Notes in Shelford, R.W.C., 1916. A naturalist in Borneo. London, T. Fisher Unwin, pp. 1–331.
- Robinson, O.J., Lockwood, J.L., Stringham, O.C., Fefferman, N.H., 2015. A novel tool for making policy recommendations based on PVA: helping theory become practice. Conserv. Lett. 8, 190–198.
- Rookmaaker, L.C., 1977. The distribution and status of the rhinoceros, *Dicerorhinus sumatrensis*, in Borneo—a review. Bijdragen tot de Dierkunde. Amsterdam 47, 197–204.
- Rookmaaker, K., 2014. www.rhinoresourcecenter.com (accessed June 2014).
- Rushton, S.P., Ormerod, S.J., Kerby, G., 2004. New paradigms for modeling species distributions? J. Appl. Ecol. 41, 193–200.
- Sabah Government, 2014. http://ww2.sabah.gov.my/htan_caims/Level%20%20frame%20pgs/Class%20%20Frames/tabin_fr.htm.
- Schaffer, N.E., Zahari, Z.Z., Suri, M.S.M., Jainudeen, M.R., Jeyendran, R.S., 1994. Ultrasonography of the reproductive anatomy in the Sumatran rhinoceros (*Dicerorhinus sumatrensis*). J. Zoo Wildl. Med. 25, 337–348.
- Schneeberger, W.F., 1945. The Kerayan-Kelabit highland of Central Northeast Borneo. Geogr. Rev. 35, 544–562.
- Shelford, R.W.C., 1916. A naturalist in Borneo. Edited with a biographical introduction by Edward B. Poulton. London, T. Fisher Unwin, pp. 1–331.
- Silva, G.S. de, 1968. Wildlife conservation in the state of Sabah. In: Talbot, L.M., Talbot, M.H. (Eds.), Conservation in Tropical South East Asia, Gland. IUCN Publications, N.S., pp. 144–150. 10.
- Strien, N.J. van, 1974. *Dicerorhinus sumatrensis* (Fischer), the Sumatran or two-horned rhinoceros: a study of literature. Meded. Landbouwhogeschool Wageningen 74, 1–82.
- Strien, N.J. van, 1979. Report on a survey of the proposed Banumuda extension of Kutai, East Kalimantan. Rep.
- Strien, N.J. van, 1981. Survey of Sumatran rhinos, Banumuda area, East Kalimantan. WWF Yb. 1980–1981, 459–460.
- Strien, N.J. van, 1986. The Sumatran rhinoceros, *Dicerorhinus sumatrensis* in the Gunung Leuser National Park, Sumatra, Indonesia: its distribution, ecology and conservation. Mamalia Depicta 12, Paul Parey, Hamburg, pp. 1–200.
- Strien, N.J. van, 2002. Conservation proram for Sumatran and Javan rhino in Indonesia and Malaysia. In: Schwammer, H.M., Foose, T.J., Olson, D. Schöling Verlag Münster. pp. 231–236.
- Tajuddin Abdullah, M., Sinun, Waidi, Mathew, D., Gadas, S., Bernard, H., Gumpil, J., Murray, L., Ola, B., 1995. Sumatran rhinoceros and wildlife survey East of Segama, Danum Valley, Sabah, Malaysia. Report to SWD/MTED/YS/WCS. pp. 1–43.
- Tear, T.H., Scott, J.M., Hayward, P.H., Griffith, B., 1995. Recovery Plans and the Endangered Species Act: are criticisms supported by data? Conserv. Biol. 9, 182–195.
- Thayaparan, S.T., 2008. Final Field Report. SOS Rhino Borneo, Sabah, Malaysia.
- Vargas, A., Breitenmoser, C., Breitenmoser, U., 2009. Iberian lynx ex-situ conservation: an interdisciplinary approach. Fundación Biodiversidad, Madrid, Spain. pp. 1–534.
- Veloz, S.D., 2009. Spatially autocorrelated sampling falsely inflates measures of accuracy for presence-only niche models. J. Biogeogr. 36, 2290–2299.
- Wallace, A.R., 1874. On the rhinoceros of Borneo. Proc. Zool. Soc. Lond. 498–499.
- Wisz, M.S., Hijmans, R.J., Peterson, A.T., Graham, C.H., Guisan, A., 2008. Effects of sample size on the performance of species distribution models. Divers. Distrib. 14, 763–773.
- Wood, S.N., 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. J. R. Stat. Soc. Ser. B 73, 3–36.
- WWF, 2013. <http://www.worldwildlife.org/press-releases/footprints-of-sumatran-rhinos-discovered-in-borneo>.
- WWF-Malaysia, 1986. Tabin Wildlife Reserve, Sabah. A preliminary management plan. Report IUCN/WWF Project No. 3050. WWF, Malaysia, Kuala Lumpur.
- Zafir, A.W.A., Payne, J., Mohamed, A., Lau, C.F., Sharma, D.S.K., Alfred, R., Williams, A.C., Nathan, S., Ramono, W.S., Clements, G.R., 2011. Now or never: what will it take to save the Sumatran rhinoceros *Dicerorhinus sumatrensis* from extinction? Oryx 45, 225–233.
- Zon, A.P.M. van der, 1977. Sumatran rhino in Kalimantan (Borneo). Tiger Paper 4, 12.