



RESEARCH REPOSITORY

*This is the author's final version of the work, as accepted for publication following peer review but without the publisher's layout or pagination.
The definitive version is available at:*

<https://doi.org/10.1016/j.eng.2018.06.005>

Dowla, M.A.N.N.U., Edwards, I., O'Hara, G., Islam, S. and Ma, W. (2018) Developing wheat for improved yield and adaptation under a changing climate: Optimization of a few key genes. Engineering

<http://researchrepository.murdoch.edu.au/id/eprint/41317/>

Copyright: © 2018 THE AUTHORS
It is posted here for your personal use. No further distribution is permitted.

Accepted Manuscript

Review

Developing Wheat for Improved Yield and Adaptation under a Changing Climate: Optimization of a Few Key Genes

M.A.N. Nazim Ud Dowla, Ian Edwards, Graham O'Hara, Shahidul Islam, Wujun Ma

PII: S2095-8099(17)30248-5
DOI: <https://doi.org/10.1016/j.eng.2018.06.005>
Reference: ENG 88

To appear in: *Engineering*

Received Date: 26 April 2017
Revised Date: 25 August 2017
Accepted Date: 20 November 2017

Please cite this article as: N.N.U. Dowla, I. Edwards, G. O'Hara, S. Islam, W. Ma, Developing Wheat for Improved Yield and Adaptation under a Changing Climate: Optimization of a Few Key Genes, *Engineering* (2018), doi: <https://doi.org/10.1016/j.eng.2018.06.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Research
Crop Genetics and Breeding—Review

Developing Wheat for Improved Yield and Adaptation under a Changing Climate: Optimization of a Few Key Genes

M.A.N. Nazim Ud Dowla ^a, Ian Edwards ^{a,b}, Graham O'Hara ^a, Shahidul Islam ^a, Wujun Ma ^{a,*}

^a School of Veterinary and Life Sciences, Murdoch University, Perth, WA 6150, Australia

^b Edstar Genetics Pty. Ltd., Perth, WA 6150, Australia

* Corresponding author.

E-mail address: w.ma@murdoch.edu.au

ARTICLE INFO

Article history:

Received 26 April 2017

Revised 25 Aug 2017

Accepted 20 November 2017

Available online

Keywords:

Phenology

Wheat

Vernalization

Photoperiod

Drought

Climate change

ABSTRACT

Wheat grown under rain-fed conditions is often affected by drought worldwide. Future projections from a climate simulation model predict that the combined effects of increasing temperature and changing rainfall patterns will aggravate this drought scenario and may significantly reduce wheat yields unless appropriate varieties are adopted. Wheat is adapted to a wide range of environments due to the diversity in its phenology genes. Wheat phenology offers the opportunity to fight against drought by modifying crop developmental phases according to water availability in target environments. This review summarizes recent advances in wheat phenology research, including vernalization (*Vrn*), photoperiod (*Ppd*), and also dwarfing (*Rht*) genes. The alleles, haplotypes, and copy number variation identified for *Vrn* and *Ppd* genes respond differently in different climatic conditions, and thus could alter not only the development phases but also the yield. Compared with the model plant *Arabidopsis*, more phenology genes have not yet been identified in wheat; quantifying their effects in target environments would benefit the breeding of wheat for improved drought tolerance. Hence, there is scope to maximize yields in water-limited environments by deploying appropriate phenology gene combinations along with *Rht* genes and other important physiological traits that are associated with drought resistance.

1. Introduction

Cereals constitute a prime global human food source. Among them, wheat (*Triticum aestivum* L.) ranks as the second most important food after rice, and is the most widely cultivated cereal in the world. It is one of the central pillars of food security, supplying 20% of total calories and a similar portion of total protein to the world's population [1]. The average global wheat yield is $3.3 \text{ t}\cdot\text{hm}^{-2}$; however, it varies widely, with regional averages ranging from $1.7 \text{ t}\cdot\text{hm}^{-2}$ in Australia to a potential of up to $9 \text{ t}\cdot\text{hm}^{-2}$ in other parts of the world (data from FAOSTAT database 2015 [2]). The yield penalty is usually due to different environmental stresses that reduce yield potential by 69.1% [3]. In most developed countries, wheat is mainly grown in rain-fed marginal land, where inadequate and erratic rainfall limits the yield (Table 1) [4]. Drought is a key stress that constrains wheat production on about $6.5 \times 10^7 \text{ hm}^2$ of land worldwide [5] and reduces yield by up to 50% [6]. Modeling exercises have revealed that water stress in marginal wheat-growing environments reduces 50%–90% of their yield potential under irrigated conditions [7]. In 2012, the overall global wheat production decreased by 1.4%, mainly due to severe drought in the United States, Europe, and central Asia (data from FAOSTAT database 2013 [2]). The Australian wheat yield dropped by 46% in 2006 compared with the yield trend of the previous 50 years, resulting in billion dollar losses for the wheat industry (data from FAOSTAT database 2012 [2]).

Table 1

Major wheat producers of the world: five year (2008–2012) averages of production, area harvested, and area irrigated for wheat [4].

Region	Production (t)	Area harvested (10^7 m^2)	Area irrigated (10^7 m^2)
Australia	24.51	13 650	—
Canada	26.22	9200	—
China (mainland)	116.15	24 110	—
France	38.37	5540	30.24
India	84.36	28 640	—
Pakistan	23.40	8860	7335.00
Russian	52.19	24 090	—
Turkey	19.99	7980	—
Ukraine	20.34	6480	46.90
USA	60.91	20 060	1662.00
World	678.02	220 400	13 241.50

The impact of future drought episodes on wheat production is expected to increase due to the effects of climate change on temperature and precipitation. It is estimated that the $1 \text{ }^\circ\text{C}$ increase of temperature that has occurred during the last 29 years has resulted in a 6% reduction of wheat yield compared with the expected yield without global warming effects [8]. According to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC), the global mean temperature will increase by $3.7 \text{ }^\circ\text{C}$ by the end of this century, with incidents of hottest days and coolest nights occurring 50% more frequently than at present [9]. Changes in the precipitation pattern coupled with increasing temperature would affect the major crop production of the world, and wheat production could decline by 23.2%–27.2% by 2050, unless protective measures for limiting global warming or appropriate cultivars and crop management practices are adopted [10]. Wheat production in low-latitude sites would be more vulnerable with the rise of a $3\text{--}5 \text{ }^\circ\text{C}$ temperature scenario, compared with production in high latitude regions, and yields could decline by up to 40% with an increase of $2 \text{ }^\circ\text{C}$ in temperature [11]. In Australia, wheat belts are typical of those in a Mediterranean climate: most precipitation occurs in winter, followed by less-frequent rain in spring and hot dry summer. Thus, water stress in spring is the major factor limiting yield improvement in these regions and often coincides with stem elongation, flowering, and grain filling [12]. In these environments, terminal heat often combines with drought during the grain-filling period and further limits grain yield [13]. In their Fifth Assessment Report, the IPCC predicted that an increase in mean annual temperature by $2.2\text{--}5 \text{ }^\circ\text{C}$ with +5% to –30% change in precipitation patterns will result in the expansion of drought-affected areas by 5.4%, 4.6%, and 3.8% by 2030, 2050, and 2070, respectively [9]. In this situation, plant breeders must be well-prepared to embrace the challenges of climate change and to feed the world by developing varieties that are better adapted to water-limited environments. Better utilization of the available genetic resources of wheat is essential in order to maintain and maximize wheat yield potential in water-limited environments, and the optimization of phenology is one of the most effective ways to achieve this goal.

Phenology is the key factor for crop adaptation to a particular environment. A proper understanding of the genetic control of phenological traits will enable breeders to develop crops that are better adapted to a specific environment. It is well documented that yield loss due to drought depends on the growth stage at drought

occurrence, as well as the duration and intensity of the stress [14,15]. Spike development, from terminal spikelet initiation to anthesis, is the most important phase in determining grain yield, as it has been observed that a heavier spike at anthesis is positively correlated with grain yield [16] and can be manipulated without affecting other phases [17]. Therefore, adverse effects of drought could be minimized by ensuring that the most sensitive developmental stages do not occur during stress periods [18]. Hence, fine-tuning of flowering and the duration of developmental phases are advocated for better adaptation of wheat in water-limited environments or to escape from these constraints [19–22]. Phenology genes also regulate the physiological development of wheat [23], and some morpho-physiological traits have been identified as effective in breeding drought-adaptive varieties [24,25]. Taking account of many important traits and their interactions in stress environments, a sound understanding of the genetic control and physiological basis of drought tolerance would facilitate the improvement of yield in water-limited environments. We acknowledge the importance of good agronomic practices, that is, management, and several other traits involved in physiological mechanisms to reduce adverse effects of drought; however, this review focuses on the phenology genes as one of the most important measures to avoid drought stress (Fig. 1). Therefore, an effort has been made to summarize the progress achieved to date on key phenology genes and the integration of this knowledge in breeding new varieties that are adapted to future climate change. Moreover, an attempt has been taken to compile information on molecular markers for the identified alleles of vernalization (*Vrn*) and photoperiod (*Ppd*) genes that will help in evaluating the cultivars that are adapted to target environments as well as marker-assisted breeding.

2. Phenology genes

Wheat is adapted to a wide range of agricultural environments [26]. The synchrony of flowering to a wider range of climatic conditions is largely controlled by ① *Vrn* genes (exposure to cold temperature requirement), ② *Ppd* genes (photoperiod sensitivity), and ③ autonomous earliness *per se* (*Eps*) genes [27]. Hence, the adaptation of a genotype to a particular environment depends on the interaction of these three groups of genes.




Fig. 1. Drought shield: complementary approaches to sustain wheat yield in water-limited environments. WUE: water-use efficiency; RUE: radiation use efficiency).

2.1. Vernalization genes

Vernalization promotes the switching of the plant vegetative phase to the reproductive phase by inducing floral primordia from leaf primordia in the shoot apical meristem [28,29]. In wheat, three genes determine the vernalization requirement: *Vrn1*, *Vrn2*, and *Vrn3* [30–32]. The three orthologous *Vrn1* genes—*Vrn-A1*, *Vrn-B1*, and *Vrn-D1*—are located on the long arms of the homoeologous chromosomes 5A, 5B, and 5D, respectively, in

common wheat, and mainly control the vernalization requirement [30,31,33,34]. The *Vrn2* gene is also located on the long arm of 5A; and *Vrn3* is located on the short arm of chromosome 7B [32,35–37]. Winter wheat varieties require a certain period of cold to induce flowering, whereas varieties that flower without vernalization are referred to as spring types. The dominant alleles of *Vrn-A1*, *Vrn-B1*, *Vrn-D1*, and *Vrn3* are responsible for the spring growth habit; thus, a dominant allele at any of the three *Vrn1* loci confers a spring type. On the other hand, *Vrn2* is dominant for the winter type and is epistatic to dominant alleles of *Vrn1* [31,37–39]. *Vrn2* is a floral repressor that delays flowering, but vernalization under long days suppresses the expression of *Vrn2* and enhances the expression of *Vrn1* [40]. Multiple alleles of *Vrn1* with different levels of responses to vernalization and effects on flowering have been identified (Table 2) [41–55], and have an adaptive value [56–61]. The extent of flowering depends on the basal level of *Vrn1* expression [62]; some alleles of *Vrn1* are expressed without prior cold treatment, thus allowing flowering without vernalization [36,62,63]. Mutations in the promoter or deletion in the first intron of the *Vrn1* gene cause expression of *Vrn1* without vernalization, and the alleles lacking the larger section are more active during earlier flowering without vernalization [36,44,45,64,65]. On the other hand, varieties of wheat and barley flower early without vernalization when they lack a functional copy of the *Vrn2* gene [31,37]. The *Vrn3* gene also expresses at a high level when *Vrn2* is absent, and active alleles of *Vrn3* accelerate flowering irrespective of day length or vernalization [32]. Thus, five loci of *Vrn* genes influence flowering by controlling the vernalization requirement of wheat cultivars in different parts of the world [30,32,66,67].

Table 2Current status of the identified alleles for *Vrn1* and *Ppd1* loci.

Gene	Allele	Sequence variation from wild type	Response to light/temperature	Reference
<i>Vrn-A1</i>	<i>Vrn-A1a</i>	231-bp and 140-bp insertions in the promoter region of common wheat	Insensitive	[44]
	<i>Vrn-A1b</i>	20-bp deletion in the promoter region of common wheat	Insensitive	[44]
	<i>Vrn-A1c</i>	7222 bp deletion in intron 1	Insensitive	[45]
	<i>Vrn-A1d</i>	32-bp deletion in the promoter region of tetraploid wheat	Insensitive	[44]
	<i>Vrn-A1e</i>	54-bp deletion in the promoter region of tetraploid wheat	Insensitive	[44]
<i>Vrn-B1</i>	<i>vrn-A1</i>	Wild type	Sensitive	[44]
	<i>Vrn-B1a</i>	6850 bp deletion in intron 1	Insensitive	[45]
	<i>Vrn-B1b</i>	6850 bp and 36 bp deletion in intron 1	Insensitive	[46]
	<i>Vrn-B1c</i>	817 bp deletion and 432 bp duplication in intron 1	Insensitive	[47]
	<i>vrn-B1</i>	Wild type	Sensitive	[45]
<i>Vrn-D1</i>	<i>Vrn-D1a</i>	4235 bp deletion in intron 1	Insensitive	[45]
	<i>Vrn-D1b</i>	C replaced by A at translation site in CARG-box of wild type	Facultative	[48]
	<i>vrn-D1</i>	Wild type	Sensitive	[45]
<i>Ppd-A1</i>	<i>Ppd-A1a1</i>	1085 bp deletion in the promoter region	Insensitive	[49]
	<i>Ppd-A1a2</i>	1027 bp deletion in the promoter region	Insensitive	[20]
	<i>Ppd-A1a3</i>	1117 bp deletion in the promoter region	Insensitive	[50]
	<i>Ppd-A1a4</i>	684 bp deletion in the promoter region	Insensitive	[51]
	<i>ppd-A1b</i>	Wild type	Sensitive	[50]
<i>Ppd-B1</i>	<i>Ppd-B1a.1</i>	308 bp insertion in the promoter region	Insensitive	[49]
	<i>Ppd-B1a.2</i>	Four copy of <i>Ppd-B1</i>	Insensitive	[52]
	<i>Ppd-B1a.3</i>	3 copy of <i>Ppd-B1</i>	Insensitive	[52]
	<i>Ppd-B1a.4</i>	2 copy of <i>Ppd-B1</i>	Insensitive	[52]
	<i>Ppd-B1e</i>	—	—	[53]
	<i>ppd-B1b</i>	Wild type	Sensitive	[52]
<i>Ppd-D1</i>	<i>Ppd-D1a.1</i>	2089 bp deletion in the promoter region	Insensitive	[54]
	<i>Ppd-D1a.2</i>	5 bp deletion in exon 7	Intermediate	[55]
	<i>ppd-D1b.1</i>	Wild type	Sensitive	[54]
	<i>ppd-D1b.2</i>	Insertion of transposable element in the intron 1	Sensitive	[55]

2.2. Photoperiod genes

Wheat is a long day plant, requiring exposure to long days (> 14 h light) for flowering, whereas photoperiod-insensitive varieties flower early in short days (10 h or less light) [54,68,69]. This photoperiod sensitivity is controlled by the semi-dominant homoeologous *Ppd1* gene on the short arm of chromosome group 2; as is the case with *Vrn1*, the dominant allele confers photoperiod insensitivity [70–74]. The effects of the photoperiod-insensitive allele *Ppd* were studied thoroughly by Worland [22] over a 14 year period in different wheat-growing regions; their work revealed that insensitive *Ppd1* advances flowering time by 9–15 days, and that this earliness can be utilized to obtain yield advantages in water-limited environments by drought avoidance. The early *Ppd* gene also has some pleiotropic effects including reduced plant height and number of tillers, and fewer spikelets per ear [73]. However, an increase in spikelet fertility can compensate for the yield penalty [74]. It is clear that *Ppd* insensitivity brings forward the time of terminal spikelet formation, thus advancing the flowering time by reducing the number of spikelets in the ear. However, it does not influence the rates of leaf and flower primordial production. There is also variation among the potency of three *Ppd1a* loci, where plants with *Ppd-A1a* and *Ppd-D1a* are earlier in flowering than plants with *Ppd-B1a* [52]. In the same way that a number of *Vrn1* alleles have been identified, a number of alleles and their haplotypes have also been identified recently for all three homoeologous loci of the *Ppd* gene

(Table 2) in both bread and durum wheat [49,51,53–55,75]. These findings have a great agronomic importance for deployment in breeding programs.

2.3. Earliness per se (*Eps*) genes

The *Eps* genes control flowering time independent of temperature and photoperiod. To date, very few *Eps* genes have been identified in wheat, but several quantitative trait locus (QTL) studies revealed that most of the chromosome groups carry such genes and that they are present as QTL effects rather than as major genes in the *Ppd* and *Vrn* pathways [74,76–81]. The *Eps* genes are involved in the fine-tuning of flowering time [82], and hence can be utilized for adaptation to specific climatic conditions.

3. Molecular intervention of phenology genes

In concordance with studies on *Arabidopsis*, important progress has been made at the molecular level to elucidate the flowering pathway in wheat. Molecular and sequence analysis revealed that *Vrn1* encodes a MADS-box transcription factor similar to the *Arabidopsis* meristem identity genes *APETALA1* (*API*), *CAULIFLOWER* (*CAL*), and *FRUITFULL* (*FRU*), which regulate the shoot apical meristem to determine the transition from vegetative to reproductive development [36]. Insertions, deletions, and mutations in the promoter region are associated with allelic variation of *Vrn1* [44]. Following this finding, a series of molecular markers have been developed (Table 3) [32,44,45,48,52,54,55,83,84] and successfully utilized to identify allele frequency of the local wheat cultivars as well as these from the CIMMYT (Centro Internacional de Mejoramiento de Maíz y Trigo) collection [45,83,85,86]. The *Vrn2* encodes a zinc finger-CCT domain transcription factor and is a floral repressor, down-regulated by both vernalization treatment and short day length [37]. *Vrn2* plays a very similar role to that of *FLOWERING LOCUS C* (*FLC*) in *Arabidopsis* but actually has no orthologues, suggesting an independent evolution of the vernalization pathways [87]. Vernalization gene *Vrn3* is similar to *Arabidopsis* *FLOWERING LOCUS T* (*FT*), and the dominant allele is associated with a retro element insertion in the *Triticum aestivum* L. (*TaFT*) promoter, results in early flowering [32]. Recent screening of a set of Chinese wheat cultivars led to the discovery of two more dominant alleles of *Vrn3*, and 80 days variation in heading has been observed due to their action [83]. Allelic variations of these *Vrn1* genes quantify the vernalization effects, and determine flowering time by interacting with photoperiod gene *Ppd1*. The latter is a member of a pseudo response regulator (PRR) gene family in which insensitivity is associated with deletion or transposon insertion within the promoter region, as well as with copy number variation [52,54]. In wheat, *Ppd-1* directly regulates the *FLOWERING LOCUS T1* (*FT1*); mutants with promoter deletions result in the overexpression of *FT1*, causing early flowering [51]. Markers have been developed to identify the *Ppd* mutants with different promoter deletions (Table 2) that will facilitate their effects on the flowering time of wheat [49,51,53,54].

Table 3
Polymerase chain reaction (PCR) markers for the different vernalization and photoperiod response alleles.

Allele	Primer	Primer sequence (5'-3')	Annealing temperature (°C)	Product size	Reference
<i>Vrn-A1a</i>	VRNA1F	GAAAGGAAAAATTCTGCTCG	50	965 and 876	[44]
<i>Vrn-A1b</i>				714	
<i>Vrn-A1c</i>	VRN1-INT1R	TGCACCTTCCC(C/G)CGCCCAT		734	
<i>vrn-A1</i>				734	
<i>Vrn-A1</i>	BT706	CATTGTTCCCTCCTGTCCACCC	63	1431	[84]
	BT750	ATTACTCGTACAGCCATCTCAGCC			
<i>Vrn-A1c</i> (Langdon)	Ex1/C/F	GTTTCTCCACCGAGTCATGGT	55.6	522	[45]
	Intr1/A/R3	AAGTAAGACAACACGAATGTGAGA			
<i>Vrn-A1c</i> (IL 369)	Intr1/A/F2	AGCTCCACGGTTTGAAAGTAA	58.9	1170	
	Intr1/A/R3	AAGTAAGACAACACGAATGTGAGA			
<i>vrn-A1</i>	Intr1/C/F	GCACTCCTAACCCACTAACC	56	1068	
	Intr1/AB/R	TCATCCATCATCAAGCAAA			
<i>Vrn-B1a</i>	Intr1/B/F	CAAGTGGAACGGTTAGGACA	58	709	
	Intr1/B/R3	CTCATGCCAAAAATTGAAGATGA			
<i>vrn-B1</i>	Intr1/B/F	CAAGTGGAACGGTTAGGACA	56.4	1149	
	Intr1/B/R4	CAAATGAAAAGGAATGAGAGCA			
<i>Vrn-D1</i>	Intr1/D/F	GTTGTCTGCCTCATCAAATCC	61	1671	
	Intr1/D/R3	GGTCACTGGTGGTCTGTGC			
<i>vrn-D1</i>	Intr1/D/F	GTTGTCTGCCTCATCAAATCC	61	997	
	Intr1/D/R4	AAATGAAAAGGAACGGAGCG			
<i>Vrn-D1a</i>	VRN1DF	CGACCCGGGCGGCACGAGTG	65	612	[48]
	VRN1-SNP161CR	AGGATGGCCAGGCCAAAACG			
<i>Vrn-D1b</i>	VRN1DF	CGACCCGGGCGGCACGAGTG			
	VRN1-SNP161AR	AGGATGGCCAGGCCAAAACG			
<i>Vrn-3</i>	FT-B-INS-F	CATAATGCCAAGCCGGTGAGTAC	63	1200	[32]
	FT-B-INS-R	ATGTCTGCCAATTAGCTAGC			
<i>vrn-3</i>	FT-B-NOINS-F or FT-B-NOINS-F2	ATGCTTTCGCTTGCCATCC or GCTGTGTGATCTTGCTCTCC	57	1140 or 691	
	FT-B-NOINS-R	CTATCCCTACGGCCATTAG			
<i>Ppd-D1a</i>	Ppd-D1_F	ACGCCTCCCCTACTACTG	54	288	[54]
	Ppd-D1_R2	AND CACTGGTGGTAGCTGAGATT			
<i>ppd-D1b</i>	Ppd-D1_F	ACGCCTCCCCTACTACTG	54	414	
	Ppd-D1_R1 and	GTTGGTTCAAACAGAGAGC			
16 bp deletion in exon 8	Ppd-D1exon8_F1	GATGAACATGAAACGGG	52	320 or 326 and 22, 257, 69, and 22	
	Ppd-D1exon8_R1	GTCTAAATAGTAGGTACTAGG			
<i>Ppd-B1</i>	Ppd-B1exon3SNP_F1	AGACGATTCATTCGGCTCC	55	471, 328 and 155	
	Ppd-B1exon3SNP_R1	TCTGAATGATGATACACCATG			
	Ppd-B1_2ndcopy_F1	TAACTGCTCGTCACAAGTGC	55	425 and 475	
	Ppd-B1_2ndcopy_R1	CCGGAACCTGAGGATCATC			

Allele	Primer	Primer sequence (5'-3')	Annealing temperature (°C)	Product size	Reference
5bp deletion in exon 7	D5-1F	GAATGGCTTCCTGCTC	50	1032 or 1027	[55]
	D5-1R	GATGGCGAAACCTTATT			
	D5-2F	GTGTCCTTTGCGAATCCTT	53	184 or 179	
	D5-2R	TTGGAGCCTTGCTTCATCT			
A TE insertion in intron 1	D520F	AGGTCCTTACTCATCTCAATCTCA	50	2612	
	D520R	CTCCATTGTTGGTGTGTTA			
	D78F	CCATTCGAGGAGACGATTTCAT	55	1005	
	D78R	CTGAGAAAGAACAGAGTCAA			
Truncated <i>Ppd-B1</i> gene in the "Chinese Spring" allele	219H05F2	TAAGTCTCCTCACAAAGTGC	56	425	[52]
	97J10R2	CCGGAACCTGAGGATCATC			
Intact <i>Ppd-B1</i> copies in the "Chinese Spring" allele	Ppd-B1_F25	AAAACATTATGCATATAGCTTGTGTC	58	994	
	Ppd-B1_R70	CAGACATGGACTCGGAACAC			
Intact <i>Ppd-B1</i> copies in the "Sonora64"/"Timstein" allele	Ppd-B1_F31	CCAGGCGAGTGATTACACA	58	223	
	Ppd-B1_R36	GGGCACGTAAACACACCTTT			
<i>Vrn-A1b</i>	<i>Vrn-P2</i>	CCTGCCGGAATCCTCGTTTT	63	147 or 167	[83]
		CTACGCCCTACCCTCCAACA			
<i>Vrn-B1b</i>	<i>Vrn-P7</i>	CCAATCTCACATGCCTCCAA	59	215 or 252	
		ATGCGCCATGAACAACAAAG			
<i>Vrn-B3c</i>	<i>Vrn-P14</i>	GCTTTGAACTCCAAGGAGAA ATAATCAGCAGGTGAACCAG	52	1401	
<i>vrn-B3/Vrn-B3</i>	<i>Vrn-P15</i>	ACTCATCATCACCTTCCT TAATGCTTAATTCGTGGCTG	51	1499	
<i>Vrn-B3</i> promoter	<i>Vrn-P16</i>	GTCCATACAAATCATGCCAC TTCTGACAGTTTTAGTTGCG	51	491	
<i>Vrn-B3</i> promoter	<i>Vrn-P17</i>	GCTTTCGCTTGCCATCCCAT GCGGGAACGCTAATCTCCTG	62	898	
<i>Vrn-B3</i> promoter	—	TTTGAGACAGGAGATTAGCG	53	1131	
		ACCATCATGAGGCACCATTA	52	1425	
		GCTTTGAACTCCAAGGAGAA			
		ATAATCAGCAGGTGAACCAG			
		CCGTTCAACATCTATTGCTC	55	1259	
CACCCAAATCCTTCATCTCA					
<i>Vrn-B3-RT</i>	—	GGAGGTGATGTGCTACGAGA TTGTAGAGCTCGGCGAAGTC	55	147	

The complicated interaction of these phenology genes has resulted in two conflicting models of the flowering regulatory network: The first model, designated as *Vrn2* to *FT* [88], recommends that *Vrn2* represses *FT* expression but that vernalization during winter slightly up-regulates *Vrn1*, causing down-regulation of *Vrn2* and the release of *FT* expression. This *FT* then interacts with *Ppd1* and again up-regulates the *Vrn1* beyond the threshold to initiate flowering under long day length [89]. By contrast, the second model, known as *FT* to *Vrn2*, was proposed by Shimada et al. [90], who suggested that *Vrn1* promotes *FT* transcription, which down-regulates *Vrn2* to initiate flowering based on the fact that the maintained vegetative phase (*mvp*) mutants lacking *Vrn1* fail to up-regulate *FT*. Subsequent detailed experimentation by Distelfeld and Dubcovsky [88] with the *mvp* mutants segregating for *Vrn1* and *Vrn2* deletions resulted in evidence to contradict both of the previously proposed models; we therefore suggest that more investigation should be conducted to elucidate the flowering network of wheat, and that doing so may lead to the identification of more genes that interact in the flowering pathway.

4. Dwarfing genes

The introduction of dwarfing (*Rht*) genes into cereals, including wheat, was a key driver of the green revolution. Since then, *Rht-B1b* and *Rht-D1b* (previously known as *Rht1* and *Rht2*, respectively) are the most commonly adopted *Rht* genes in wheat-breeding programs throughout the world [91]. Together, these two semi-dwarfing genes produce the dwarf phenotype, whereas alone in combination with their counterpart *Rht-B1a* or *Rht-D1a*, they produce semi-dwarf plants in nature. The plants with these genes are less prone to lodging and are more effective in partitioning assimilates to the grain. Some researchers have suggested that the improved yield potential of such varieties is only limited to a favorable growth environment [92,93]. However, these specific *Rht* genes are insensitive to endogenous gibberellins, and produce shorter plants with smaller cells [94]. These smaller size cells are consequently responsible for the shorter coleoptile length, less early vigor, smaller leaf area, lower water-use efficiencies, and poor seedling establishment, especially in water-limited environments [95–99]. The insensitivity to gibberellins of both the *Rht-B1b* and *Rht-D1b* alleles is due to single nucleotide substitutions that create a translational stop codon, TGA, reducing the plant's ability to respond to gibberellins [100].

Most of the world's wheat is grown without irrigation; because of the dependence on seasonal rainfall, the potential yield is often hampered by water scarcity. About 50% of rainwater can be lost directly through soil evaporation, whereas early vigor can increase water-use efficiency by 25% and thus improve yield [101–104]. Again, deep sowing of longer season varieties is often recommended to obtain yield benefits in dry areas, such as those occurring in southern Australia, but seedling establishment is impaired when dwarf/semi-dwarf varieties are sown more than 5 cm deep [96]. Consequently, farmers wait until the first rains before sowing, resulting in between 140–330 kg yield loss per week per hectare being reported in Australian wheat crops [105,106]. Wheat varieties with longer coleoptiles are able to emerge sooner when sown deep, and have greater early vigor [107,108]. Moreover, early vigor and longer coleoptiles help plants to avoid the phytotoxic effects of residual herbicides, compete against weeds, and reduce evaporative water loss by shading. Hence, breeding for vigorous seedling growth and breeding for longer coleoptiles are the prime objectives for the better adaptation of wheat in water-limited environments [109–111]. A project with these objectives is currently underway at the International Maize and Wheat Improvement Center (CIMMYT) in Mexico[†].

On the other hand, a number of *Rht* genes such as *Rht 7*, *Rht 8*, *Rht 9*, *Rht 13*, and *Rht 14* have been reported, which have potential in reducing plant height without affecting seedling vigor and tissue response to gibberellins [112–114]. Under stress conditions, taller varieties store assimilates in the stem and do not depend entirely on current assimilation for grain filling [115]. Several studies across many favorable and unfavorable environments demonstrated that plants with heights of 70–100 cm are better yielders than those that are taller or shorter than this range [97,116]. Therefore, the accumulation of minor *Rht* genes or combination with one of the gibberellins insensitive genes for shorter plant height are desirable [116,117], as shown by different studies that used *Rht8* and/or *Rht13* alleles with *Rht1* and/or *Rht2* to maximize yields compared with other dwarf/semi-dwarf varieties [91]. Markers linked to these *Rht* alleles make it easier to select both alleles simultaneously across a large population [118,119].

5. Physiological aspects of phenology and dwarfing genes

Grain yield is strongly influenced by the timing of developmental stages in a particular environment, making crop phenology a critical component for yield physiology [120]. Moisture stress at the reproductive stage, especially that period from a few weeks before anthesis to a few days after anthesis, has the most critical effect on crop yields in water-limited environments [25,121]. Passioura [122] emphasized the importance of water use, water-use efficiency, and harvest index (HI) for crop yields in dry areas. In dry environments, an important portion of soil moisture that could be available for transpiration is evaporated from a barren soil surface, thus indirectly affecting dry matter accumulation by limiting water availability to roots, and modifying canopy

[†] Edwards IB—personal communication.

temperature [123]. In this situation, faster early seedling growth is beneficial to prevent evaporation by shading. Moreover, late-flowering cultivars continue to produce tillers until they receive the signal for reproductive development, and many of them cannot produce fertile spikes, but put pressure on the available soil moisture through normal transpiration. In this regard, heading date and effective tiller number should be additional considerations for improving water-use efficiency in varieties being developed for drought environments. Moreover, water requirement varies throughout the growth period and is higher during seed setting and development stages. Hence, there is an opportunity to improve yield through changes in crop development. The synchronization of crop developmental stages by phenological adjustment with seasonal moisture availability should be the most important target for new wheat varieties being developed for water-limited environments such as those that occur in Mediterranean climate regions.

HI and ultimately final grain yield largely depend on pre- and post-anthesis biomass production, mobilization of assimilates to florets, and the pattern of water supply during the life cycle [111,124]. One strategy for raising the HI may be increasing the assimilate movement to developing florets, which will prevent floret abortion before anthesis. This can be done by increasing the duration of spike growth with a reduction in the earlier period for larger ear development [16]. Moreover, this larger ear will also contribute more photosynthate during grain filling along with the flag leaf, thereby increasing the HI. Studies on two alternative spring alleles of *Vrn-A1* have shown their significant influence on the variation in root and vegetative morphology such as rosette growth habit, plant height, and leaf length [125]. A significant relation has recently been observed between the duration of pre-anthesis growth phases and the tillering and dry matter accumulation [126]. A detailed study of Australian wheat cultivars over several years and a wide range of locations has revealed that cultivars with one spring allele in any of the three *Vrn1* loci are the earliest in heading when compared with cultivars having two spring alleles [127]. Again, spring alleles in all three *Vrn1* loci have very small effects in forwarding the heading date, which suggests the presence of epistatic or overdose effects. This study also showed that *Vrn-B1* has a weaker effect on the reduction of heading time compared with *Vrn-A1* or *Vrn-D1*. Recently, however, it has been shown that *Vrn-B1* has the greatest effect on grain yield [128].

Semi-dwarf varieties with *Rht-D1b* are advantageous over *Rht-B1b* in environments with high maximum temperatures and lower rainfall during the flowering and grain-filling periods, as *Rht-D1b* is associated with less leaf porosity in plants relative to *Rht-B1b*, leading to slow transpiration before heading and leaving more soil moisture for later use [128,129]. Plants reduce their water use during drought stress by means of accelerated leaf desiccation and death, which causes a reduction of current photosynthate [130–133]. As a result, stem reserves become an important source of carbohydrate for grain filling [134–136]. However, *Rht-B1b* and *Rht-D1b* genes reduce stem reserves by 35% and 39%, respectively [137]; hence, taller varieties often perform better in stress environments when there is a shortage of assimilates, compared with the modern dwarf cultivars.

6. Future strategies

Difficulties in the identification and precise measurement of key physiological determinants of yield is the bottleneck in the improvement of drought tolerance in plants, and its complex genetic control makes progress more difficult [138–140]. Hence, the improvement of plant traits at both physiological and molecular levels is vital in order to address this complex issue.

The current use of automated high-throughput plant-phenotyping facilities greatly assists researchers in phenotyping plants more precisely and accurately. An in-depth understanding of plant physiology will help dissecting the genetic components of drought tolerance, while molecular and genomic tools will help to identify candidate genes and QTLs for drought-tolerance traits. The integration of physiology with molecular tools will provide new insights into gene function. To optimize output in drought research, a detailed knowledge of the growing environment and of genotype-environment interactions is essential. Fine-tuning a genotype to a specific environment is possible by combining the best-suited alleles of phenology genes to adapt better in the existing environment.

Recent studies have revealed that copy number variation of phenology genes also plays a vital role in crop adaptation. An increased copy number of *Ppd-B1* confers earlier flowering, and an increased copy number of *Vrn-A1* requires a longer vernalization period and is thus associated with late flowering [52]. A similar investigation for the copy number variation of *Ppd-B1* in Australian wheat cultivars resulted in the identification of five alleles with one to four copies as well as a null copy of *Ppd-B*, where plants with an allele with a lower copy number were the latest in heading relative to plants with an allele having more copies [53]. In addition, the haplotypes variation identified in other studies for these genes was found to affect several yield-contributing parameters, and thus adaptation to different environments [51,55]. As a result, several attempts have been made to determine the value of the alleles of *Vrn1* and *Ppd1* genes over the past few years in the local environments of different countries [52,55,68,73,83,84,86,127,128,141]. In most cases, the plant material did not cover all the available alleles present in nature, or even the same allele in different genetic backgrounds. Thus, obtaining the true effect of an allele in breeding a variety will warrant the development of appropriate near-isogenic line (NIL) populations of the locally adapted cultivars with different alleles, which will require significant effort. An earlier

example of a successful attempt is Triple Dirk, which was developed by Pugsley [30] to study the alleles of the *Vrn1* gene; at present, the John Innes Center in the UK is performing substantial research in developing this type of population for different *Ppd1* alleles. Such efforts will certainly advance research into phenology genes in optimizing plant development and productivity in local water-limited environments. Therefore, intensive and thorough research to optimize the effect of each allele/haplotype of the phenology genes would enable plant breeders to determine the basic genetic architecture of wheat in each key growing environment, for better yields under stress conditions. Thus, once molecular and physiological tools are used to identify and prove the efficiency of different traits and their regulating genes for drought tolerance, these various useful traits could be aggregated in the base population through a marker-assisted gene pyramiding scheme, as demonstrated by Servin et al. [142]. In summary, success against the adverse effects of climate change relies on the consequences of proper characterization of target environments (i.e., soil properties, precipitation pattern, drought severity, and etc.); and then on designing an appropriate crop ideotype that combines useful phenology with other drought-tolerance-attributing genes, along with good management practices.

7. Conclusion

Drought is a major threat to world agriculture, and is predicted to worsen in the near future due to climate change. Wheat is the most widely grown cereal crop in the world and is vital for global food security. Altering the developmental stages and maturity of wheat is one of the best ways to combat drought without compromising yield. However, current knowledge about the number of genes that control flowering and maturity in wheat is limited. Based on knowledge obtained from the model plant species *Arabidopsis*, in which more than 80 genes have been reported to control flowering, it is logical to conclude that many new genes and genetic pathways for wheat flowering and maturity are yet to be discovered. However, the fact is that different genetic pathways finally converge, interact, and ultimately lead to the activation of floral identity genes in the floral primordia [143], and these interacting networks that promote flowering are yet to be unveiled. As significant research efforts are currently underway, knowledge on wheat-flowering genes and pathways will increase over time and will require advances in computational biology in order to integrate and interpret this information. In addition, future international collaboration will help to combine the cumulative efforts of the research underway in different research groups and disciplines; the challenge for the breeders will be to integrate this work into new genetic combinations.

Compliance with ethics guidelines

M.A.N. Nazim Ud Dowla, Ian Edwards, Graham O'Hara, Shahidul Islam, and Wujun Ma declare that they have no conflict of interest or financial conflicts to disclose.

References

- [1] Braun HJ, Atlin G, Payne T. Multi-location testing as a tool to identify plant response to global climate change. In: Reynolds MP, editor. Climate change and crop production. Wallingford: Center for Agriculture and Biosciences International; 2010. p. 115–38.
- [2] FAOSTAT database collections [Internet]. Rome: Food and Agriculture Organization of the United Nations; c2017 [cited 2015 Oct 23]. Available from: <http://www.fao.org/faostat/en/#data/QC>.
- [3] Boyer JS. Plant productivity and environment. *Science* 1982;218(4571):443–8.
- [4] AQUASTAT main database (2015) [Internet]. Rome: Food and Agriculture Organization of the United Nations; c2017 [cited 2015 Oct 22]. Available from: <http://www.fao.org/nr/water/aquastat/data/query/index.html?lang=en>.
- [5] Zaynali Nezhad K, Weber WE, Röder MS, Sharma S, Lohwasser U, Meyer RC, et al. QTL analysis for thousand-grain weight under terminal drought stress in bread wheat (*Triticum aestivum* L.). *Euphytica* 2012;186(1): 127–38.
- [6] Byerlee D, Morris M. Research for marginal environments: are we underinvested? *Food Policy* 1993;18(5):381–93. doi:10.1016/0306-9192(93)90061-F.
- [7] Morris ML, Belaid A, Byerlee D. Part 1: wheat and barley production in rainfed marginal environments of the developing world. In: 1990–91 CIMMYT world wheat facts and trends: wheat and barley production in rainfed marginal environments of the developing world. Mexico: International Maize and Wheat Improvement Center; 1991. p. 1–28.
- [8] Lobell DB, Gourdji SM. The influence of climate change on global crop productivity. *Plant Physiol* 2012;160(4):1686–97. doi:10.1104/pp.112.208298. PMID:23054565
- [9] Intergovernmental Panel on Climate Change. Summary for policymakers. In: Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, et al., editors. Climate change 2014: impacts, adaptation, and vulnerability. Part A: global and sectoral aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press; 2014. p. 1–32.
- [10] Nelson GC, Rosegrant MW, Koo J, Robertson R, Sulser T, Zhu T, et al. Climate change: impact on agriculture and costs of adaptation. Washington, DC: International Food Policy Research Institute; 2009.
- [11] Reynolds MP, editor. Climate change and crop production. Wallingford: Center for Agriculture and Biosciences International; 2010.
- [12] Turner NC. Sustainable production of crops and pastures under drought in a Mediterranean environment. *Ann Appl Biol* 2004;144(2):139–47. doi:10.1111/j.1744-7348.2004.tb00327.x.
- [13] Nachit MM. Durum breeding research to improve dryland productivity in the Mediterranean region. In: Nachit MM, Baum M, Porceddu E, Monneveux P, Picard E, editors. SEWANA (South Europe, West Asia and North Africa) durum research network: Proceedings of the SEWANA Durum Network Workshop; 1995 Mar 20–23; Aleppo, Syria. Aleppo: International Center for Agricultural Research in the Dry Areas; 1998. p. 1–15.
- [14] Lopez CG, Banowitz GM, Peterson CJ, Kronstad WE. Dehydrin expression and drought tolerance in seven wheat cultivars. *Crop Sci* 2003;43(2):577–82.

- [15] Serraj R, Hash CT, Rizvi SMH, Sharma A, Yadav RS, Bidinger FR. Recent advances in marker-assisted selection for drought tolerance in pearl millet. *Plant Prod Sci* 2005;8(3):334–7. doi:10.1626/pss.8.334.
- [16] Slafer GA, Abeledo LG, Miralles DJ, Gonzalez FG, Whitechurch EM. Photoperiod sensitivity during stem elongation as an avenue to raise potential yield in wheat. *Euphytica* 2001;119(1–2):191–7. doi:10.1023/A:1017535632171.
- [17] Whitechurch EM, Slafer GA. Contrasting *Ppd* alleles in wheat: Effects on sensitivity to photoperiod in different phases. *Field Crops Res* 2002;73(2–3):95–105. doi:10.1016/S0378-4290(01)00188-5.
- [18] Saini HS, Westgate ME. Reproductive development in grain crops during drought. *Adv Agron* 1999;68:59–96.
- [19] Richards RA. Crop improvement for temperate Australia: Future opportunities. *Field Crops Res* 1991;26(2):141–69. doi:10.1016/0378-4290(91)90033-R.
- [20] Worland AJ. The influence of flowering time genes on environmental adaptability in European wheats. *Euphytica* 1996;89(1):49–57. doi:10.1007/BF00015718.
- [21] Debaeke P. Scenario analysis for cereal management in water-limited conditions by the means of a crop simulation model (STICS). *Agronomie* 2004;24(6–7):315–26. doi:10.1051/agro:2004035.
- [22] Cockram J, Jones H, Leigh FJ, O’Sullivan D, Powell W, Laurie DA, et al. Control of flowering time in temperate cereals: genes, domestication, and sustainable productivity. *J Exp Bot* 2007;58(6):1231–44. doi:10.1093/jxb/erm042. PMID:17420173
- [23] Barrett B, Bayram M, Kidwell K, Weber WE. Identifying AFLP and microsatellite markers for vernalization response gene *Vrn-B1* in hexaploid wheat using reciprocal mapping populations. *Plant Breed* 2002;121(5):400–6. doi:10.1046/j.1439-0523.2002.732319.x.
- [24] Acevedo E. Assessing crop and plant attributes for cereal improvement in water-limited Mediterranean environments. In: Srivastava JP, Porceddu E, Acevedo E, Varma S, editors. Drought tolerance in winter cereals. Chichester: John Wiley and Sons; 1987. p. 303–20.
- [25] Reynolds M, Foulkes MJ, Slafer GA, Berry P, Parry MA, Snape JW, et al. Raising yield potential in wheat. *J Exp Bot* 2009;60(7):1899–918. doi:10.1093/jxb/erp016. PMID:19363203
- [26] Curtis BC, Rajaram S, Gómez Macpherson H, editors. Bread wheat: improvement and production. Rome: Food and Agriculture Organization of the United Nations; 2002.
- [27] Kato K, Yamagata H. Method for evaluation of chilling requirement and narrow-sense earliness of wheat cultivars. *Jpn J Breed* 1988;38(2):172–86. Japanese. doi:10.1270/jsbbs1951.38.172.
- [28] Trevisk B. The central role of the *VERNALIZATION1* gene in the vernalization response of cereals. *Funct Plant Biol* 2010;37(6):479–87. doi:10.1071/FP10056.
- [29] Le Gouis J, Bordes J, Ravel C, Heumez E, Faure S, Praud S, et al. Genome-wide association analysis to identify chromosomal regions determining components of earliness in wheat. *Theor Appl Genet* 2012;124(3):597–611. doi:10.1007/s00122-011-1732-3. PMID:22065067
- [30] Pugsley AT. A genetic analysis of the spring-winter habit of growth in wheat. *Aust J Agric Res* 1971;22(1):21–31. doi:10.1071/AR9710021.
- [31] Dubcovsky J, Lijavetzky D, Appendino L, Tranquilli G. Comparative RFLP mapping of *Triticum monococcum* genes controlling vernalization requirement. *Theor Appl Genet* 1998;97(5–6):968–75. doi:10.1007/s001220050978.
- [32] Yan L, Fu D, Li C, Blechl A, Tranquilli G, Bonafede M, et al. The wheat and barley vernalization gene *VRN3* is an orthologue of *FT*. *Proc Natl Acad Sci USA* 2006;103(51):19581–6. doi:10.1073/pnas.0607142103. PMID:17158798
- [33] Law CN, Worland AJ, Giorgi B. The genetic control of ear-emergence time by chromosomes 5A and 5D of wheat. *Heredity* 1976;36(1):49–58. doi:10.1038/hdy.1976.5.
- [34] Galiba G, Quarrie SA, Sutka J, Morgounov A, Snape JW. RFLP mapping of the vernalization (*Vrn1*) and frost resistance (*Frl*) genes on chromosome 5A of wheat. *Theor Appl Genet* 1995;90(7–8):1174–9. doi:10.1007/BF00222940. PMID:24173081
- [35] Law CN, Wolfe MS. Location of genetic factors for mildew resistance and ear emergence time on chromosome 7B of wheat. *Can J Genet Cytol* 1966;8(3):462–70. doi:10.1139/g66-056.
- [36] Yan L, Loukoianov A, Tranquilli G, Helguera M, Fahima T, Dubcovsky J. Positional cloning of the wheat vernalization gene *VRN1*. *Proc Natl Acad Sci USA* 2003;100(10):6263–8. doi:10.1073/pnas.0937399100. PMID:12730378
- [37] Yan L, Loukoianov A, Blechl A, Tranquilli G, Ramakrishna W, SanMiguel P, et al. The wheat *VRN2* gene is a flowering repressor down-regulated by vernalization. *Science* 2004;303(5664):1640–4. doi:10.1126/science.1094305. PMID:15016992
- [38] Stelmakh AF. Genetic effects of the *Vrn1-3* loci and specific action of the dominant *Vrn3* allele in common bread wheat. *Cytol Genet* 1987;21(4):278–86. Russian.
- [39] Tranquilli G, Dubcovsky J. Epistatic interaction between vernalization genes *Vrn-A^m1* and *Vrn-A^m2* in diploid wheat. *J Hered* 2000;91(4):304–6. doi:10.1093/jhered/91.4.304. PMID:10912677
- [40] Trevisk B, Hemming MN, Dennis ES, Peacock WJ. The molecular basis of vernalization-induced flowering in cereals. *Trends Plant Sci* 2007;12(8):352–7. doi:10.1016/j.tplants.2007.06.010. PMID:17629542
- [41] Tsunewaki K, Jenkins BC. Monosomic and conventional gene analyses in common wheat. II. Growth habit and awnedness. *Jpn J Genet* 1961;36(11–12):428–43. Japanese. doi:10.1266/jjg.36.428.
- [42] Roberts DMA, MacDonald MD. Evidence for the multiplicity of alleles at *Vrn1*, the winter–spring habit locus in common wheat. *Can J Genet Cytol* 1984;26(2):191–3. doi:10.1139/g84-032.
- [43] Koval SF, Goncharov NP. Multiple allelism at the *VRN1* locus of common wheat. *Acta Agron Hung* 1998;46(2):113–9.
- [44] Yan L, Helguera M, Kato K, Fukuyama S, Sherman J, Dubcovsky J. Allelic variation at the *VRN-1* promoter region in polyploid wheat. *Theor Appl Genet* 2004;109(8):1677–86. doi:10.1007/s00122-004-1796-4. PMID:15480533
- [45] Fu D, Szűcs P, Yan L, Helguera M, Skinner JS, von Zitzewitz J, et al. Large deletions within the first intron in *VRN-1* are associated with spring growth habit in barley and wheat. *Mol Genet Genomics* 2005;273(1):54–65. doi:10.1007/s00438-004-1095-4. PMID:15690172
- [46] Santra DK, Santra M, Allan RE, Campbell KG, Kidwell KK. Genetic and molecular characterization of vernalization genes *Vrn-A1*, *Vrn-B1*, and *Vrn-D1* in spring wheat germplasm from the Pacific Northwest region of the USA. *Plant Breed* 2009;128(6):576–84. doi:10.1111/j.1439-0523.2009.01681.x.
- [47] Mílec Z, Tomková L, Sumíková T, Pánková K. A new multiplex PCR test for the determination of *Vrn-B1* alleles in bread wheat (*Triticum aestivum* L.). *Mol Breed* 2012;30(1):317–23. doi:10.1007/s11032-011-9621-7.
- [48] Zhang J, Wang Y, Wu S, Yang J, Liu H, Zhou Y. A single nucleotide polymorphism at the *Vrn-D1* promoter region in common wheat is associated with vernalization response. *Theor Appl Genet* 2012;125(8):1697–704. doi:10.1007/s00122-012-1946-z. PMID:22875177
- [49] Nishida H, Yoshida T, Kawakami K, Fujita M, Long B, Akashi Y, et al. Structural variation in the 5' upstream region of photoperiod-insensitive alleles *Ppd-A1a* and *Ppd-B1a* identified in hexaploid wheat (*Triticum aestivum* L.), and their effect on heading time. *Mol Breed* 2013;31(1):27–37. doi:10.1007/s11032-012-9765-0.
- [50] Wilhelm EP, Turner AS, Laurie DA. Photoperiod insensitive *Ppd-A1a* mutations in tetraploid wheat (*Triticum durum* Desf.). *Theor Appl Genet* 2009;118(2):285–94. doi:10.1007/s00122-008-0898-9. PMID:18839130
- [51] Muterko A, Kalendar R, Cockram J, Balashova I. Discovery, evaluation and distribution of haplotypes and new alleles of the *Photoperiod-A1* gene in wheat. *Plant Mol Biol* 2015;88(1–2):149–64. doi:10.1007/s11103-015-0313-2. PMID:25851614

- [52] Díaz A, Zikhali M, Turner AS, Isaac P, Laurie DA. Copy number variation affecting the *Photoperiod-B1* and *Vernalization-A1* genes is associated with altered flowering time in wheat (*Triticum aestivum*). *PLoS One* 2012;7(3):e33234. doi:10.1371/journal.pone.0033234. PMID:22457747
- [53] Cane K, Eagles HA, Laurie DA, Trevaskis B, Vallance N, Eastwood RF, et al. *Ppd-B1* and *Ppd-D1* and their effects in southern Australian wheat. *Crop Pasture Sci* 2013;64(2):100–14. doi:10.1071/CP13086.
- [54] Beales J, Turner A, Griffiths S, Snape JW, Laurie DA. A pseudo-response regulator is misexpressed in the photoperiod insensitive *Ppd-D1a* mutant of wheat (*Triticum aestivum* L.). *Theor Appl Genet* 2007;115(5):721–33. doi:10.1007/s00122-007-0603-4. PMID:17634915
- [55] Guo Z, Song Y, Zhou R, Ren Z, Jia J. Discovery, evaluation and distribution of haplotypes of the wheat *Ppd-D1* gene. *New Phytol* 2010;185(3):841–51. doi:10.1111/j.1469-8137.2009.03099.x. PMID:20002313
- [56] Gotoh T. Genetic studies on growth habit of some important spring wheat cultivars in Japan, with special reference to the identification of the spring genes involved. *Jpn J Breed* 1979;29(2):133–45. Japanese. doi:10.1270/jsbbs1951.29.133.
- [57] Stelmakh A. Geographic distribution of *Vrn*-genes in landraces and improved varieties of spring bread wheat. *Euphytica* 1990;45(2):113–8.
- [58] Goncharov NP. Genetic resources of wheat related species: The *Vrn* genes controlling growth habit (spring vs. winter). *Euphytica* 1998;100(1–3):371–6. doi:10.1023/A:1018323600077.
- [59] Stelmakh AF. Genetic systems regulating flowering response in wheat. *Euphytica* 1998;100(1–3):359–69. doi:10.1023/A:1018374116006.
- [60] Iwaki K, Nakagawa K, Kuno H, Kato K. Ecogeographical differentiation in East Asian wheat, revealed from the geographical variation of growth habit and *Vrn* genotype. *Euphytica* 2000;111(2):137–43. doi:10.1023/A:1003862401570.
- [61] Iwaki K, Haruna S, Niwa T, Kato K. Adaptation and ecological differentiation in wheat with special reference to geographical variation of growth habit and *Vrn* genotype. *Plant Breed* 2001;120(2):107–14. doi:10.1046/j.1439-0523.2001.00574.x.
- [62] Trevaskis B, Bagnall DJ, Ellis MH, Peacock WJ, Dennis ES. MADS-box genes control vernalization-induced flowering in cereals. *Proc Natl Acad Sci USA* 2003;100(22):13099–104. doi:10.1073/pnas.1635053100. PMID:14557548
- [63] Danyluk J, Kane NA, Breton G, Limin AE, Fowler DB, Sarhan F. *TaVRT-1*, a putative transcription factor associated with vegetative to reproductive transition in cereals. *Plant Physiol* 2003;132(4):1849–60. doi:10.1104/pp.103.023523. PMID:12913142
- [64] Szűcs P, Skinner JS, Karsai I, Cuesta-Marcos A, Haggard KG, Corey AE, et al. Validation of the *VRN-H2/VRN-H1* epistatic model in barley reveals that intron length variation in *VRN-H1* may account for a continuum of vernalization sensitivity. *Mol Genet Genomics* 2007;277(3):249–61. doi:10.1007/s00438-006-0195-8. PMID:17151889
- [65] Hemming MN, Fieg S, James Peacock W, Dennis ES, Trevaskis B. Regions associated with repression of the barley (*Hordeum vulgare*) *VERNALIZATION1* gene are not required for cold induction. *Mol Genet Genomics* 2009;282(2):107–17. doi:10.1007/s00438-009-0449-3. PMID:19404679
- [66] McIntosh RA, Hart GE, Devos KM, Gale MD, Rogers WJ. Catalogue of gene symbols for wheat. In: Slinkard AE, editor. Proceedings of the 9th International Wheat Genetics Symposium; 1998 Aug 2–7; Saskatoon, SK, Canada. Saskatoon: University of Saskatchewan; 1998. p. 123–72.
- [67] Goncharov NP. Genetics of growth habit (spring vs winter) in common wheat: confirmation of the existence of dominant gene *Vrn4*. *Theor Appl Genet* 2003;107(4):768–72. doi:10.1007/s00122-003-1317-x. PMID:12838388
- [68] Foulkes MJ, Sylvester-Bradley R, Worland AJ, Snape JW. Effects of a photoperiod-response gene *Ppd-D1* on yield potential and drought resistance in UK winter wheat. *Euphytica* 2004;135(1):63–73. doi:10.1023/B:EUPH.0000009542.06773.13.
- [69] Kumar S, Sharma S, Chaudhary S, Tyagi A, Mishra P, Priyadarshini A, et al. Genetics of flowering time in bread wheat *Triticum aestivum*: complementary interaction between vernalization-insensitive and photoperiod-insensitive mutations imparts very early flowering habit to spring wheat. *J Genet* 2012;91(1):33–47. doi:10.1007/s12041-012-0149-3. PMID:22546824
- [70] Welsh JR, Keim DL, Piratash B, Richards RD. Genetic control of photoperiod response in wheat. In: Sears ER, Sears LMS, editors. Proceedings of the Fourth International Wheat Genetics Symposium; 1973 Aug 6–11; Columbia, MO, USA. Columbia: Agricultural Experiment Station, College of Agriculture, University of Missouri; 1973. p. 879–84.
- [71] Law CN, Sutka J, Worland AJ. A genetic study of day-length response in wheat. *Heredity* 1978;41(2):185–91. doi:10.1038/hdv.1978.87.
- [72] Börner A, Worland AJ, Plaschke J, Schumann E, Law CN. Pleiotropic effects of genes for reduced height (*Rht*) and day-length insensitivity (*Ppd*) on yield and its components for wheat grown in middle Europe. *Plant Breed* 1993;111(3):204–16. doi:10.1111/j.1439-0523.1993.tb00631.x.
- [73] Worland AJ, Börner A, Korzun V, Li WM, Petrović S, Sayers EJ. The influence of photoperiod genes on the adaptability of European winter wheats. *Euphytica* 1998;100(1–3):385–94. doi:10.1023/A:1018327700985.
- [74] Snape JW, Butterworth K, Whitechurch E, Worland AJ. Waiting for fine times: genetics of flowering time in wheat. *Euphytica* 2001;119(1–2):185–90. doi:10.1023/A:1017594422176.
- [75] Takenaka S, Kawahara T. Evolution and dispersal of emmer wheat (*Triticum* sp.) from novel haplotypes of *Ppd-1* (photoperiod response) genes and their surrounding DNA sequences. *Theor Appl Genet* 2012;125(5):999–1014. doi:10.1007/s00122-012-1890-y. PMID:22639190
- [76] Chen Y, Carver BF, Wang S, Cao S, Yan L. Genetic regulation of developmental phases in winter wheat. *Mol Breed* 2010;26(4):573–82. doi:10.1007/s11032-010-9392-6.
- [77] Law CN, Suarez E, Miller TE, Worland AJ. The influence of the group 1 chromosomes of wheat on ear-emergence times and their involvement with vernalization and day length. *Heredity* 1998;80(1):83–91. doi:10.1046/j.1365-2540.1998.00286.x.
- [78] Kulwal PL, Roy JK, Balyan HS, Gupta PK. QTL mapping for growth and leaf characters in bread wheat. *Plant Sci* 2003;164(2):267–77. doi:10.1016/S0168-9452(02)00409-0.
- [79] Tóth B, Galiba G, Fehér E, Sutka J, Snape JW. Mapping genes affecting flowering time and frost resistance on chromosome 5B of wheat. *Theor Appl Genet* 2003;107(3):509–14. doi:10.1007/s00122-003-1275-3. PMID:12734655
- [80] Hanocq E, Laperche A, Jaminon O, Lainé AL, Le Gouis J. Most significant genome regions involved in the control of earliness traits in bread wheat, as revealed by QTL meta-analysis. *Theor Appl Genet* 2007;114(3):569–84. doi:10.1007/s00122-006-0459-z. PMID:17171391
- [81] Griffiths S, Simmonds J, Leverington M, Wang Y, Fish L, Sayers L, et al. Meta-QTL analysis of the genetic control of ear emergence in elite European winter wheat germplasm. *Theor Appl Genet* 2009;119(3):383–95. doi:10.1007/s00122-009-1046-x. PMID:19430758
- [82] Hoogendoorn J. A reciprocal F₁ monosomic analysis of the genetic control of time of ear emergence, number of leaves and number of spikelets in wheat (*Triticum aestivum* L.). *Euphytica* 1985;34(2):545–58. doi:10.1007/BF00022954.
- [83] Chen F, Gao M, Zhang J, Zuo A, Shang X, Cui D. Molecular characterization of vernalization and response genes in bread wheat from the Yellow and Huai Valley of China. *BMC Plant Biol* 2013;13:199. doi:10.1186/1471-2229-13-199. PMID:24314021

- [84] Eagles HA, Cane K, Trevaskis B. Veery wheats carry an allele of *Vrn-A1* that has implications for freezing tolerance in winter wheats. *Plant Breed* 2011;130(4):413–8. doi:10.1111/j.1439-0523.2011.01856.x.
- [85] Eagles HA, Cane K, Vallance N. The flow of alleles of important photoperiod and vernalisation genes through Australian wheat. *Crop Pasture Sci* 2009;60(7):646–57. doi:10.1071/CP09014.
- [86] Iqbal M, Navabi A, Yang RC, Salmon DF, Spaner D. Molecular characterization of vernalization response genes in Canadian spring wheat. *Genome* 2007;50(5):511–6. doi:10.1139/G07-028. PMID:17612620
- [87] Dubcovsky J, Loukoianov A, Fu D, Valarik M, Sanchez A, Yan L. Effect of photoperiod on the regulation of wheat vernalization genes *VRN1* and *VRN2*. *Plant Mol Biol* 2006;60(4):469–80. doi:10.1007/s11103-005-4814-2. PMID:16525885
- [88] Distelfeld A, Dubcovsky J. Characterization of the maintained vegetative phase deletions from diploid wheat and their effect on *VRN2* and *FT* transcript levels. *Mol Genet Genomics* 2010;283(3):223–32. doi:10.1007/s00438-009-0510-2. PMID:20063107
- [89] Distelfeld A, Li C, Dubcovsky J. Regulation of flowering in temperate cereals. *Curr Opin Plant Biol* 2009;12(2):178–84. doi:10.1016/j.pbi.2008.12.010. PMID:19195924
- [90] Shimada S, Ogawa T, Kitagawa S, Suzuki T, Ikari C, Shitsukawa N, et al. A genetic network of flowering-time genes in wheat leaves, in which an *APETALA1/FRUITFULL*-like gene, *VRN1*, is upstream of *FLOWERING LOCUS T*. *Plant J* 2009;58(4):668–81. doi:10.1111/j.1365-3113X.2009.03806.x. PMID:19175767
- [91] Rebetzke GJ, Bonnett DG, Ellis MH. Combining gibberellic acid-sensitive and insensitive dwarfing genes in breeding of higher-yielding, sesqui-dwarf wheats. *Field Crops Res* 2012;127:17–25. doi:10.1016/j.fcr.2011.11.003.
- [92] Waddington SR, Ransom JK, Osmanzai M, Saunders DA. Improvement in the yield potential of bread wheat adapted to northwest Mexico. *Crop Sci* 1986;26(4):698–703.
- [93] Chapman SC, Mathews KL, Trethowan RM, Singh RP. Relationships between height and yield in near-isogenic spring wheats that contrast for major reduced height genes. *Euphytica* 2007;157(3):391–7. doi:10.1007/s10681-006-9304-3.
- [94] Keyes GJ, Paolillo DJ, Sorrells ME. The effects of dwarfing genes *Rht1* and *Rht2* on cellular dimensions and rate of leaf elongation in wheat. *Ann Bot* 1989;64(6):683–90. doi:10.1093/oxfordjournals.aob.a087894.
- [95] Donald CM, Puckridge DW. The ecology of the wheat crop. In: Lazenby A, Matheson EM, editors. *Australian field crops*. Volume 1. Wheat and other temperate cereals. Sydney: Angus and Robertson; 1975. p. 288–303.
- [96] Allan RE. Agronomic comparisons between *Rht1* and *Rht2* semidwarf genes in winter wheat. *Crop Sci* 1989;29(5):1103–8. doi:10.2135/cropsci.1989.0011183X002900050001x.
- [97] Richards RA. The effect of dwarfing genes in spring wheat in dry environments. I. Agronomic characteristics. *Aust J Agric Res* 1992;43(3):517–27. doi:10.1071/AR920517.
- [98] Rebetzke GJ, Botwright TL, Moore CS, Richards RA, Condon AG. Genotypic variation in specific leaf area for genetic improvement of early vigour in wheat. *Field Crops Res* 2004;88(2–3):179–89. doi:10.1016/j.fcr.2004.01.007.
- [99] Botwright TL, Rebetzke GJ, Condon AG, Richards RA. Influence of the gibberellin-sensitive *Rht8* dwarfing gene on leaf epidermal cell dimensions and early vigour in wheat (*Triticum aestivum* L.). *Ann Bot* 2005;95(4):631–9. doi:10.1093/aob/mci069. PMID:15655105
- [100] Peng J, Richards DE, Hartley NM, Murphy GP, Devos KM, Flintham JE, et al. “Green revolution” genes encode mutant gibberellin response modulators. *Nature* 1999;400(6741):256–61. doi:10.1038/22307. PMID:10421366
- [101] Leuning R, Condon AG, Dunin FX, Zegelin S, Denmead OT. Rainfall interception and evaporation from soil below a wheat canopy. *Agric For Meteorol* 1994;67(3–4):221–38. doi:10.1016/0168-1923(94)90004-3.
- [102] Siddique KHM, Tennant D, Perry MW, Belford RK. Water use and water use efficiency of old and modern wheat cultivars in a Mediterranean-type environment. *Aust J Agric Res* 1990;41(3):431–47. doi:10.1071/AR9900431.
- [103] Regan KL, Siddique KHM, Turner NC, Whan BR. Potential for increasing early vigour and total biomass in spring wheat. II. Characteristics associated with early vigour. *Aust J Agric Res* 1992;43(3):541–53. doi:10.1071/AR920541.
- [104] López-Castañeda C, Richards RA. Variation in temperate cereals in rainfed environments III. Water use and water-use efficiency. *Field Crops Res* 1994;39(2–3):85–98. doi:10.1016/0378-4290(94)90011-6.
- [105] Doyle AD, Marcellos H. Time of sowing and wheat yield in northern New South Wales. *Aust J Exp Agric Anim Husband* 1974;14(66):93–102. doi:10.1071/EA9740093.
- [106] Shackley BJ, Anderson WK. Responses of wheat cultivars to time of sowing in the southern wheatbelt of Western Australia. *Aust J Exp Agric* 1995;35(5):579–87. doi:10.1071/EA9950579.
- [107] Hadjichristodoulou A, Della A, Photiades J. Effect of sowing depth on plant establishment, tillering capacity and other agronomic characters of cereals. *J Agric Sci* 1977;89(1):161–7.
- [108] Gan Y, Stobbe EH, Moes J. Relative date of wheat seedling emergence and its impact on grain yield. *Crop Sci* 1992;32(5):1275–81.
- [109] Whan BR. The emergence of semidwarf and standard wheats, and its association with coleoptile length. *Aust J Exp Agric Anim Husband* 1976;16(80):411–6. doi:10.1071/EA9760411.
- [110] Schillinger WF, Donaldson E, Allan RE, Jones SS. Winter wheat seedling emergence from deep sowing depths. *Agron J* 1998;90(5):582–6.
- [111] Richards RA, Rebetzke GJ, Condon AG, van Herwaarden AF. Breeding opportunities for increasing the efficiency of water use and crop yield in temperate cereals. *Crop Sci* 2002;42(1):111–21. PMID:11756261
- [112] Rebetzke GJ, Richards RA. Genetic improvement of early vigour in wheat. *Aust J Agric Res* 1999;50(3):291–302.
- [113] Ellis MH, Rebetzke GJ, Chandler P, Bonnett D, Spielmeier W, Richards RA. The effect of different height reducing genes on the early growth of wheat. *Funct Plant Biol* 2004;31(6):583–9. doi:10.1071/FP03207.
- [114] Rebetzke GJ, Richards RA, Fettel NA, Long M, Condon AG, Forrester RI, et al. Genotypic increases in coleoptile length improves stand establishment, vigour and grain yield of deep-sown wheat. *Field Crops Res* 2007;100(1):10–23. doi:10.1016/j.fcr.2006.05.001.
- [115] Borrell AK, Incoll LD, Simpson RJ, Dalling MJ. Partitioning of dry matter and the deposition and use of stem reserves in a semi-dwarf wheat crop. *Ann Bot* 1989;63(5):527–39. doi:10.1093/oxfordjournals.aob.a087778.
- [116] Flintham JE, Börner A, Worland AJ, Gale MD. Optimizing wheat grain yield: effects of *Rht* (gibberellin-insensitive) dwarfing genes. *J Agric Sci* 1997;128(1):11–25. doi:10.1017/S0021859696003942.
- [117] Rebetzke GJ, Richards RA, Fischer VM, Mickelson BJ. Breeding long coleoptile, reduced height wheats. *Euphytica* 1999;106(2):159–68. doi:10.1023/A:1003518920119.
- [118] Korzun V, Röder MS, Ganal MW, Worland AJ, Law CN. Genetic analysis of the dwarfing gene (*Rht8*) in wheat. Part I. Molecular mapping of *Rht8* on the short arm of chromosome 2D of bread wheat (*Triticum aestivum* L.). *Theor Appl Genet* 1998;96(8):1104–9. doi:10.1007/s001220050845.
- [119] Ellis MH, Bonnett DG, Rebetzke GJ. A 192bp allele at the *Xgwm261* locus is not always associated with the *Rht8* dwarfing gene in wheat (*Triticum aestivum* L.). *Euphytica* 2007;157(1–2):209–14. doi:10.1007/s10681-007-9413-7.

- [120] Slafer GA, Kantolic AG, Appendino ML, Miralles DJ, Savin R. Crop development: genetic control, environmental modulation and relevance for genetic improvement of crop yield. In: Sadras VO, Calderini DF, editors. Crop physiology: applications for genetic improvement and agronomy. Amsterdam: Elsevier Inc.; 2009. p. 277–308.
- [121] Fischer RA. Number of kernels in wheat crops and the influence of solar radiation and temperature. *J Agric Sci* 1985;105(2):447–61.
- [122] Passioura JB. Grain yield, harvest index, and water use of wheat. *J Aust Inst Agric Sci* 1977;43:117–20.
- [123] Jamieson PD, Francis GS, Wilson DR, Martin RJ. Effects of water deficits on evapotranspiration from barley. *Agric For Meteorol* 1995;76(1):41–58. doi:10.1016/0168-1923(94)02214-5.
- [124] Araus JL, Slafer GA, Reynolds MP, Royo C. Plant breeding and drought in C3 cereals: what should we breed for? *Ann Bot* 2002;89(7):925–40. doi:10.1093/aob/mcf049. PMID:12102518
- [125] Roberts DWA. Identification of loci on chromosome 5A of wheat involved in control of cold hardiness, vernalization, leaf length, rosette growth habit, and height of hardened plants. *Genome* 1990;33(2):247–59. doi:10.1139/g90-039.
- [126] Borràs-Geloch G, Rebetzke GJ, Richards RA, Romagosa I. Genetic control of duration of pre-anthesis phases in wheat (*Triticum aestivum* L.) and relationships to leaf appearance, tillering, and dry matter accumulation. *J Exp Bot* 2012;63(1):69–89. doi:10.1093/jxb/err230. PMID:21920907
- [127] Eagles HA, Cane K, Kuchel H, Hollamby GJ, Vallance N, Eastwood RF, et al. Photoperiod and vernalization gene effects in southern Australian wheat. *Crop Pasture Sci* 2010;61(9):721–30. doi:10.1071/CP10121.
- [128] Eagles HA, Cane K, Trevaskis B, Vallance N, Eastwood RF, Gororo NN, et al. *Ppd1*, *Vrn1*, *ALMT1* and *Rht* genes and their effects on grain yield in lower rainfall environments in southern Australia. *Crop Pasture Sci* 2014;65(2):159–70. doi:10.1071/CP13374.
- [129] Rebetzke GJ, Rattey AR, Farquhar GD, Richards RA, Condon ATG. Genomic regions for canopy temperature and their genetic association with stomatal conductance and grain yield in wheat. *Funct Plant Biol* 2013;40(1):14–33. doi:10.1071/FP12184.
- [130] Asana RD. Physiological analysis of yield of wheat in relation to water-stress and temperature. *J Post-Grad Sch Indian Agric. Res Inst* 1966;4:17–31.
- [131] Brooks A, Jenner CF, Aspinnall D. Effects of water deficit on endosperm starch granules and on grain physiology of wheat and barley. *Aust J Plant Physiol* 1982;9(4):423–36.
- [132] Aggarwal PK, Sinha SK. Effect of water stress on grain growth and assimilate partitioning in two cultivars of wheat contrasting in their yield stability in a drought-environment. *Ann Bot* 1984;53(3):329–40. doi:10.1093/oxfordjournals.aob.a086697.
- [133] Blum A. Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress. *Field Crops Res* 2009;112(2–3):119–23. doi:10.1016/j.fcr.2009.03.009.
- [134] Blum A, Shpiler L, Golan G, Mayer J, Sinmena B. Mass selection of wheat for grain filling without transient photosynthesis. *Euphytica* 1991;54(1):111–6. doi:10.1007/BF00145637.
- [135] Kiniry JR. Nonstructural carbohydrate utilization by wheat shaded during grain growth. *Agron J* 1993;85(4):844–9. doi:10.2134/agronj1993.00021962008500040013x.
- [136] Schnyder H. The role of carbohydrate storage and redistribution in the source–sink relations of wheat and barley during grain filling—a review. *New Phytol* 1993;123(2):233–45. doi:10.1111/j.1469-8137.1993.tb03731.x.
- [137] Borrell AK, Incoll LD, Dalling MJ. The influence of the *Rht1* and *Rht2* alleles on the deposition and use of stem reserves in wheat. *Ann Bot* 1993;71(4):317–26. doi:10.1006/anbo.1993.1041.
- [138] IZANLOO A, Condon AG, Langridge P, Tester M, Schnurbusch T. Different mechanisms of adaptation to cyclic water stress in two South Australian bread wheat cultivars. *J Exp Bot* 2008;59(12):3327–46. doi:10.1093/jxb/ern199. PMID:18703496
- [139] Mir RR, Zaman-Allah M, Sreenivasulu N, Trethowan R, Varshney RK. Integrated genomics, physiology and breeding approaches for improving drought tolerance in crops. *Theor Appl Genet* 2012;125(4):625–45. doi:10.1007/s00122-012-1904-9. PMID:22696006
- [140] Reynolds M, Foulkes J, Furbank R, Griffiths S, King J, Murchie E, et al. Achieving yield gains in wheat. *Plant Cell Environ* 2012;35(10):1799–823. doi:10.1111/j.1365-3040.2012.02588.x. PMID:22860982
- [141] Law CN, Worland AJ. Genetic analysis of some flowering time and adaptive traits in wheat. *New Phytol* 1997;137(1):19–28. doi:10.1046/j.1469-8137.1997.00814.x.
- [142] Servin B, Martin OC, Mézard M, Hospital F. Toward a theory of marker-assisted gene pyramiding. *Genetics* 2004;168(1):513–23. doi:10.1534/genetics.103.023358. PMID:15454561
- [143] Mouradov A, Cremer F, Coupland G. Control of flowering time: Interacting pathways as a basis for diversity. *Plant Cell* 2002;14(Suppl 1):S111–30. doi:10.1105/tpc.001362. PMID:12045273