



Murdoch
UNIVERSITY

MURDOCH RESEARCH REPOSITORY

<http://researchrepository.murdoch.edu.au>

This is the author's final version of the work, as accepted for publication following peer review but without the publisher's layout or pagination.

White, N.E. , Mawson, P.R., Dawson, R., Bunce, M.A. and Spencer, P.B.S. (2009) Characterisation and cross-species utility of 20 microsatellite markers for population and forensic applications in the endangered Carnaby's Black-cockatoo, *Calyptorhynchus latirostris*. Conservation Genetics Resources, 1 (1). pp. 341-345.

<http://researchrepository.murdoch.edu.au/4418>

Copyright © Springer Science+Business Media B.V. 2009
It is posted here for your personal use. No further distribution is permitted.

Editorial Manager(tm) for Conservation Genetics Resources
Manuscript Draft

Manuscript Number:

Title: Characterisation and cross-species utility of 20 microsatellite markers for population and forensic applications in the endangered Carnaby's Black-cockatoo, *Calyptorhynchus latirostris*

Article Type: Technical Note

Keywords: Cacatuidae; Carnaby's Black-cockatoo; microsatellite; population genetics; wildlife forensics

Corresponding Author: Dr. Peter B.S. Spencer,

Corresponding Author's Institution: Murdoch University

First Author: Nicole E White

Order of Authors: Nicole E White; Peter R Mawson; Rick Dawson; Michael A Bunce; Peter B.S. Spencer

Abstract: We characterise 20 microsatellite loci identified from the endangered Carnaby's Black-cockatoo (*Calyptorhynchus latirostris*). The primers were tested across 40 individuals from the southwest of Western Australia and displayed between 4 and 11 alleles per locus with expected heterozygosities ranging from 53 to 87% and exclusion probabilities of ≥ 0.999 . These loci will be useful in population genetic studies to facilitate conservation management decisions in addition to wildlife enforcement applications for the endangered Carnaby's Black-cockatoo. We also tested the markers in 12 high profile and smuggled species from five genera, *Cacatua*, *Callocephalon*, *Calyptorhynchus*, *Nymphicus* and *Probosciger*. These species detected between two to 19 alleles per locus with 50 to 100% amplification success.

Submitted as a Technical Note

Characterisation and cross-species utility of 20 microsatellite markers for population and forensic applications in the endangered Carnaby's Black-cockatoo, *Calyptorhynchus latirostris*

N.E. White^{*} P.R. Mawson[†], R. Dawson[§], M.A. Bunce^{*} and P.B.S. Spencer^{*}

^{*}School of Biological Sciences and Biotechnology, Murdoch University, Western Australia, Australia, 6150. [†]Species and Communities Branch & [§]Nature Protection Branch, Department of Environment and Conservation, Locked Bag 104, Bentley, D.C., Western Australia, Australia, 6983.

Abstract

We characterise 20 microsatellite loci identified from the endangered Carnaby's Black-cockatoo (*Calyptorhynchus latirostris*). The primers were tested across 40 individuals from the southwest of Western Australia and displayed between 4 and 11 alleles per locus with expected heterozygosities ranging from 53 to 87% and exclusion probabilities of ≥ 0.999 . These loci will be useful in population genetic studies to facilitate conservation management decisions in addition to wildlife enforcement applications for the endangered Carnaby's Black-cockatoo. We also tested the markers in 12 high profile and smuggled species from five genera, *Cacatua*, *Callocephalon*, *Calyptorhynchus*, *Nymphicus* and *Probosciger*. These species detected between two to 19 alleles per locus with 50 to 100% amplification success.

Keywords: Cacatuidae, Carnaby's Black-cockatoo, microsatellites, population genetics, wildlife forensics

5 Correspondence: Peter Spencer. Fax: +61 0011 89360 6303; Email: P.Spencer@murdoch.edu.au

10 The endangered Carnaby's Black-cockatoo (*Calyptorhynchus latirostris*), also known as the Short-billed Black-cockatoo is one of 19 bird species endemic to the southwest corner of Western Australia, an internationally recognised biodiversity hotspot (Myers et al. 2000). The species has suffered a 50% reduction in its range and dramatic declines in population size since the 1960's. This iconic species is threatened by a combination of illegal poaching and shooting, habitat loss, competition for nest hollows and food resources (Garnett and Crowley 2000; Saunders and Ingram 15 1995). To increase our knowledge and facilitate more informed decision-making by wildlife managers and enforcement officers, this study was undertaken and describes the variability of 20 highly polymorphic microsatellite loci for *C. latirostris* and cross-species utility for a large number of other cockatoo species of conservation concern.

20 A microsatellite library was developed from one deceased individual *C. latirostris* collected by the Department of Environment and Conservation of Western Australia (Western Australian Museum, WAM 37011). We extracted total genomic DNA from liver tissue by standard desalting: tissue digestion in cell lysis buffer (0.1 M Tris-HCl pH 8.0, 0.01 M NaCl, 0.1 M EDTA, 0.5% SDS) for nucleated blood (Longmire et al. 1997) with proteinase K (Invitrogen), 4M ammonium acetate 25 precipitation, and a final precipitation using ethanol prior to resuspending the DNA in TE buffer (10 mM Tris, 0.1 mM EDTA, pH8). Sixty-seven clones from four genomic libraries (enriched for

repetitive regions CA, ATG, CATC and TAGA; GenBank accession numbers GQ358643 – GQ358709) were created by Genetic Identification Services (<http://www.genetic-id-services.com/index.htm>). We used Primer Express 2.0 (Applied Biosystems) to design primers, of which the forward primer was labelled with an M13-tag at its 5'-end as a cost effective approach to screening large numbers of primer sets and described by Schuelke (2000). DNA fragments were separated on an Applied Biosystems 3730 DNA Analyser. Size was determined by co-running a size standard (Genescan™-500 LIZ™; Applied Biosystems, Melbourne) and fragments were scored manually with the aid of GeneMarker™ Software (Soft Genetics). All monomorphic loci were discarded and only the M13-tag primers for polymorphic loci were then exchanged for fluorescently labelled tags (Applied Biosystems).

Each singleplex (25 µL) PCR contained approximately 50 ng DNA, 1 X PCR buffer, 2.5 mM MgCl₂, 400 µM each dNTP, 0.1 mg BSA, 0.16 µM of each primer, and 2.5 units of *Taq* polymerase (Fisher Biotec). Parameters for thermal cycling were as follows: 95 °C for 3 min followed by 40 cycles at 95 °C for 30 s, 60 °C for 45 s and 72 °C for 45 s, followed by 72 °C for 10 min. Genotypic data were initially manipulated using Microsoft Excel, were checked for errors and input files were created for other programs. Descriptive statistics (number of alleles, observed and expected heterozygosities), deviations from Hardy-Weinberg equilibrium (HWE) and total exclusionary powers (*PE1*, *PE2*; Jamieson and Taylor 1997) were generated using GenAlEx 6.2 (Peakall and Smouse 2006). Linkage disequilibrium was assessed using GENEPOP (Raymond and Rousset 1995) and the frequency of null alleles (p_{null}) was estimated within the program ML-Relate (Kalinowski and Taper 2006, Kalinowski et al. 2006). Loci were assigned to a chromosome location in the zebra finch (*Taeniopygia guttata*) genome (compiled by the Washington University School of Medicine Genome Sequencing Centre) based on sequence homology using the program BLASTN 2.2.21 (Altschul et al. 1997; <http://blast.ncbi.nlm.nih.gov/Blast.cgi>) which uses a

heuristic algorithm able to detect relationships among sequences that share only isolated regions of similarity.

55 A total of 20 microsatellite loci were optimised (Table 1) and showed between 4 and 11
(mean= 6.650 ± 0.443) alleles per locus, with observed and expected heterozygosities ranging from
0.400 - 0.825 (mean= 0.635 ± 0.032) and 0.535 - 0.865 (mean= 0.699 ± 0.021), respectively.
Eighteen of 20 loci were assigned to a chromosome location (Table 1) and locus pCIB11 was found
to be Z-linked therefore only males were described, as females were homozygous. Two loci
60 pCID108 and pCID118 did not conform to Hardy-Weinberg equilibrium and were found to have a
frequency of null alleles (p_{null}) greater than >0.20 , although there may be other reasons for
observing a heterozygote deficiency at these loci such as inbreeding or Wahlund effect (Dakin and
Avisé 2004). Linkage disequilibrium was detected with loci pCID8 and pCID119, which were
subsequently removed from further analysis, and were not assessed for cross-species utility. Total
65 exclusionary powers ≥ 0.999 was reached at nine locus combination, irrespective of when only one
parent or both parents were known (Jamieson and Taylor 1997). These markers demonstrate their
utility for population genetic studies, making them suitable for detecting illegal trade (Manel et al.
2002) and providing extremely useful genetic resolution to contribute to the overall conservation,
management and protection of an internationally recognised, endangered and iconic cockatoo
70 species.

A total of 180 individuals from five genera were assessed for cross-species utility and presented in
Table 2. The number of alleles per locus ranged from two to 19 for the *Calyptrorhynchus* genus, two
to 11 alleles per locus for *Callocephalon* and *Cacatua*, two to nine alleles for *Nymphicus* and two to
75 four alleles were identified for *Probosciger* (Table 2). The amplification success rate for
polymorphic loci identified in related cockatoo species ranged from 50 to 100 per cent (Table 2),

therefore these microsatellite markers may facilitate future conservation genetic studies while reducing the associated time and costs of development.

80

Acknowledgements

This work was supported by funding from the Robert Hammond Research Scholarship (to NW), Murdoch University, Department of Environment and Conservation, Department of the Environment, Water, Heritage and the Arts and Birds Australia. The authors wish to thank Perth Zoo, Western Australian Museum, Black-cockatoo Rehabilitation Centre, Western Australian State Agricultural Biotechnology Centre (particularly Francis Brigg) and George Wildlife Park for samples and technical assistance.

85

References

- 90 Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucl Acid Res* **25**:3389-3402.
- Dakin EE, Avise JC (2004) Microsatellite null alleles in parentage analysis. *Hered* **93**:504-509.
- Garnett S, Crowley GM (2000) *The Action Plan for Australian Birds 2000*, Environment Australia, Canberra.
- 95 Jamieson A, Taylor S (1997) Comparisons of three probability formulae for parentage exclusion. *Anim Genet* **28**:397.
- Kalinowski ST, Taper ML (2006) Maximum likelihood estimation of the frequency of null alleles at microsatellite loci. *Cons Genet* **7**:991-995.
- 100 Kalinowski ST, Wagner AP, Taper ML (2006) ml-relate: a computer program for maximum likelihood estimation of relatedness and relationship. *Mol Ecol Notes* **6**:576-579.

- Longmire JL, Maltbie M, Baker RJ (1997) Use of "Lysis Buffer" in DNA isolation and its implication for museum collections. *Occasional Papers, Museum Texas Tech Uni*, 1-4.
- 105 Manel S, Berthier P, Luikart G (2002) Detecting wildlife poaching: identifying the origin of individuals with Bayesian assignment tests and multilocus genotypes. *Cons Biol* **16**:650-659.
- McInnes, LM, Dadour, IR, Stewart, ME, Ditcham, WGF, Mawson, P, Spencer, PBS (2005) Characterization of polymorphic microsatellite markers for the Carnaby's cockatoo (*Calyptorhynchus latirostris*) and related black cockatoo species. *Mol Ecol Notes* **5**:504-506.
- 110 Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* **403**:853-858.
- Peakall R, Smouse PE (2006) Genalex 6: genetic analysis in Excel: Population genetic software for teaching and research. *Mol Ecol Notes* **6**:288-295.
- Raymond M, Rousset F (1995) GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *J Hered* **86**:248-249.
- 115 Saunders DA, Ingram JA (1995) *Birds of southwestern Australia: an atlas of changes in the distribution and abundance of the wheatbelt avifauna* Surrey Beatty & Sons, Chipping Norton.
- Schuelke M (2000) An economic method for the fluorescent labelling of PCR fragments. *Nature Biotech* **18**:233-234.

Table 1 Description of 20 variable microsatellite loci isolated from the Carnaby's Black-cockatoo (*Calyptrorhynchus latirostris*)

Locus	Repeat motif	Chromosome and location (bp)	Primer sequence 5'-3' (including the fluorescent label)	N	N_A	Size range (bp)	H_O	H_E
pCIA119	(GT) ₂₁	Tgu7 (8743875)	F: ^{FAM} TGACACTTTCCTGTGGCTGC R: ATTACTTTGTTATTTCCACTGCTTGC	39	11	89 – 117	0.538	0.554
pCIA125	(CA) ₂₀	NSH	F: ^{VIC} GCTTAGCGAACATTAATCTGCAC R: TCAGGTTTCCTGAAGAGAAACCAG	40	11	115 – 139	0.775	0.865
pCIA139	(CA) ₁₈	Tgu6 (8681202)	F: GTTGCTAAGATTGGATAACACCAGATT R: ^{FAM} ATAAGTTGCAGTTTGTACGCGC	40	8	167 – 185	0.700	0.724
pCIA105	(CA) ₁₅	Tgu14 (1626425)	F: ^{PET} TCCCATCCACCCCATGC R: TCATGTGTTCTTGCCAGTTTG	40	8	107 – 123	0.725	0.745
pCIA9	(GT) ₁₅	Tgu7 (1105022)	F: ^{NED} GCTGCAGAACATGGTCACATTC R: ACTCTGATAACTCAACATTGCCCA	40	7	74 – 92	0.650	0.580
pCIA118	(GT) ₁₅	Tgu1 (9293668)	F: ^{VIC} GAGTTAAAAAAGACCCAAACCCAA R: TGTATTGTGTTAAGATGATGCATGACA	40	5	119 – 127	0.750	0.722
pCIA128	(AC) ₁₃	Tgu1 (9454118)	F: ^{FAM} TGGAATATACATAAAAATTTGTTACAGGG R: TTGTTTGCTTTTATAACAGAGAGAATG	19	4	104 – 110	0.632	0.719
pCIA138	(GT) ₁₂	Tgu3 (17204322)	F: ^{FAM} ATGAGACCAAAGGTCTTGGGTG R: TGCAATCAGATGCTGGAATCA	40	4	174 – 188	0.650	0.568
pCIB11	(ATG) ₁₆	TguZ (7531463)	F: ^{FAM} CTGCTTTTGTGCTCTAAC R: ATGCCCTATGATGAGATGC	20	7	199 – 230	0.850	0.749
pCID108‡	(TAGA) ₁₅	Tgu11 (1883671)	F: ^{FAM} ATCCTGCTGTGTTTAGAAAATACTTTGTAAT R: TCAGATATTTACATAACAGCCAGACTTG	39	7	214 – 242	0.385	0.819
pCID122	(TAGA) ₁₄	Tgu3 (1470384)	F: ^{NED} AAAGTCACTTGGACTGAGGAGCTC R: CATTTCTGAATAGTTTTAGTAGCTCTTCCTG	40	8	114 – 142	0.800	0.748

(Males)

pCID118‡	(TAGA) ₁₃	NSH	F: ^{VIC} CTATTGACTAGGAGATCTACAAGCCGT R: TGCACCCCTCTTCTGCATC	38	7	187 – 211	0.421	0.792
pCID7	(TAGA) ₁₂	Tgu2 (27449558)	F: ^{PET} AGATCTCATTACTGCTTTGCATGATAG R: TTGCAACTGCTTCTGAAACTGC	40	7	88 – 120	0.750	0.759
pCID105	(ATAG) ₁₀	Tgu5 (22806466)	F: ^{FAM} TCCCCACAACCACAGGACA R: CATGAAACATCCAAGAGCAATACAA	39	6	83 – 115	0.744	0.772
pCID109	(TCTA) ₁₀	Tgu10 (5810806)	F: ^{VIC} TGAGAGCATAAATCAAGAAGGCAT R: AGCAGAGCTCCTGCATATTGG	40	6	168 – 188	0.775	0.675
pCID114	(CTAT) ₉	Tgu3 (4409470)	F: GTCTTGTATTAATCACTGCAGAGAACTGT R: ^{FAM} ATGATACACTGGGAGCTGGGAG	40	5	120 – 136	0.550	0.570
pCID112	(ATCT) ₈	Tgu20 (1346575)	F: ^{NED} GCAGATGCAAATGTTACAGCAAC R: AACTAAATGGATATATTAATGGAGAAATAGATG	40	5	161 – 179	0.600	0.683
pCIB*	(AC) ₁₁	Tgu3 (5594880)	F: ^{NED} TGCAAATAAACTTGTCCCAGATG R: GAGACGTGAGAAATGGGTTCTGT	40	6	118 – 134	0.825	0.714
pCI5*	(CA) ₁₃	Tgu17 (2552480)	F: ^{NED} GTGCAGGGACCACTTGTATG R: CACATGGAGCATGAGCCC	40	5	164 – 190	0.400	0.535
pCI8*	(GGAT) ₁₃	Tgu18 (2246152)	F: ^{PET} GTCCTCCGAATAACCTGTGCA R: TGCTAAAAAGGTGGTAACACAGTGTC	38	5	128 – 144	0.605	0.660

For each locus (GenBank accession numbers: GQ358643 - 661), we list the core repeat motif (from the original clone), loci that could be assigned a chromosome location in the zebra finch genome, locus pCIB11 is Z-linked and only males described, no strong hits (NSH) for loci pCIA125 and pCID118, primer pair (including the labeled fluorophore), number of individuals genotypes (N), number of alleles at each locus (N_A), size range (bp), observed (H_O) and expected (H_E) heterozygosity, ‡loci showed deviation from Hardy-Weinberg equilibrium, *primers (GenBank accession numbers AY775326 – 29) redesigned for this study McInnes et al. 2005.

Table 2 Cross-species amplification of microsatellite loci developed for the Carnaby's Black-cockatoo (*Calyptorhynchus latirostris*)
[Click here to download table: Table2.doc](#)

Locus	Baudin's Black-cockatoo (<i>Calyptorhynchus baudinii</i>) (n = 24)	Yellow-tailed Black-cockatoo (<i>Calyptorhynchus funereus</i>) (n = 17)	Red-tailed Black-cockatoo (<i>Calyptorhynchus banksii</i>) (n = 45)	Glossy Black-cockatoo (<i>Calyptorhynchus lathami</i>) (n = 13)	Gang-gang cockatoo (<i>Callocephalon fimbriatum</i>) (n = 14)	Palm cockatoo (<i>Probosciger aterrimus</i>) (n = 3)	Sulphur-crested cockatoo (<i>Cacatua. galerita</i>) (n = 14)	Little corella (<i>Cacatua sanguinea</i>) (n = 10)	Western Long-billed corella (<i>Cacatua pastinator</i>) (n = 10)	Major Mitchell's cockatoo (<i>Cacatua leadbeateri</i>) (n = 12)	Galah (<i>Cacatua roseicapilla</i>) (n = 6)	Cockatiel (<i>Nymphicus hollandicus</i>) (n = 12)
pCIA119	7 (95-117)	7 (93-117)	10 (89-115)	1 (105)	1 (99)	1 (85)	3 (89 – 93)	5 (87 – 109)	4 (87 – 109)	2 (89 – 91)	1 (85)	2 (105 – 115)
pCIA125	7 (115-133)	10 (111-137)	8 (111-139)	3 (117-121)	3 (119-135)	1 (107)	3 (105 – 113)	8 (107 – 125)	7 (111 – 125)	2 (109 – 113)	4 (111 – 119)	1 (107)
pCIA139	4 (169-181)	10 (161-191)	7 (169-185)	1 (173)	1 (150)	2 (169-171)	1 (150)	-	1 (150)	-	1 (150)	3 (161 – 165)
pCIA105	5 (109-121)	9 (113-139)	9 (107-123)	6 (99-143)	10 (109-139)	4 (99-115)	7 (99 – 121)	6 (109 – 129)	5 (105 – 123)	1 (103)	7 (105 – 121)	3 (98 – 102)
pCIA9	5 (76-84)	8 (76-96)	-	2 (66-82)	-	-	-	-	-	2 (58 – 64)	2 (64 – 66)	1 (62)
pCIA118	5 (117-125)	6 (109-123)	5 (107-127)	3 (107-119)	1 (97)	2 (111-113)	2 (81 – 91)	4 (79 – 99)	4 (79 – 99)	2 (93 – 99)	1 (95)	4 (99 – 107)
pCIA128	4 (100-108)	4 (104-110)	6 (92-108)	1 (107)	4 (107-117)	2 (97-111)	2 (97 – 105)	4 (97 – 107)	4 (97 – 107)	2 (93 – 99)	-	-
pCIA138	5 (174-186)	6 (178-192)	5 (166-188)	1 (176)	1 (170)	-	3 (180 – 202)	8 (180 – 200)	6 (178 – 194)	3 (178 – 182)	4 (174 – 182)	3 (176 – 182)
pCIB11	6 (119-214)	4 (187-205)	-	-	7 (193-238)	-	3 (199 – 226)	-	-	-	-	1 (205)
pCID108	4 (218-232)	6 (210-234)	11 (214-254)	10 (234-278)	-	2 (242-250)	8 (198 – 234)	-	-	3 (222 – 246)	-	9 (222 – 266)
pCID122	7 (114-148)	7 (22-146)	10 (94-142)	-	-	3 (150-154)	-	3 (98 – 122)	3 (106 – 130)	-	-	-
pCID118	8 (187-215)	6 (191-211)	19 (187-259)	9 (249-293)	11 (191-239)	2 (169-173)	11 (201 – 257)	6 (183 – 211)	-	3 (173 – 181)	1 (177)	2 (171 – 175)
pCID7	7 (96-124)	7 (92-116)	8 (80-116)	4 (92-104)	4 (99-131)	3 (152-160)	2 (116 – 120)	4 (116 – 128)	4 (116 – 128)	4 (130 – 142)	4 (104 – 120)	-
pCID105	4 (83-111)	5 (83-115)	8 (83-115)	4 (115-127)	3 (129-137)	-	3 (117 – 125)	6 (109 – 131)	4 (115 – 127)	3 (111 – 119)	5 (115 – 135)	8 (83 – 119)
pCID109	5 (168-184)	5 (168-188)	4 (144-180)	1 (144)	4 (202-214)	3 (156-164)	4 (192 – 204)	3 (188 – 196)	3 (188 – 196)	2 (168 – 176)	-	1 (160)
pCID114	4 (124-136)	4 (120-132)	10 (116-156)	7 (140-162)	6 (124-144)	-	5 (124 – 140)	6 (120 – 144)	5 (120 – 148)	5 (108 – 128)	5 (120 – 144)	5 (120 – 140)
pCID112	4 (167-179)	8 (151-191)	-	8 (179-227)	2 (171-191)	-	-	3 (163 – 175)	-	6 (169 – 237)	-	-
pCI5*	5 (164-190)	3 (164-172)	6 (154-170)	3 (160-172)	4 (152-170)	3 (152-160)	5 (160 – 176)	-	-	-	6 (158 – 214)	-
pCI13*	6 (118-132)	7 (116-130)	6 (118-134)	2 (116-120)	2 (112-114)	2 (112-122)	3 (118 – 122)	4 (122 – 128)	4 (120 – 128)	6 (120 – 134)	2 (120 – 124)	5 (110 – 120)
pCI8*	5 (128-144)	5 (128-144)	9 (132-164)	8 (128-164)	6 (120-144)	2 (136-140)	5 (112 – 140)	6 (112 – 212)	5 (112 – 132)	2 (108 – 116)	3 (120 – 140)	-
% of loci variable	100	100	85	65	65	60	80	75	65	75	50	50

For each locus (GenBank accession numbers: GQ358643- GQ358661), we list the number of alleles and size range, (). “ – “ indicates unsuccessful amplification of PCR product, *primers (GenBank accession numbers AY775326 – 29) redesigned for this study McInnes *et al.* 2005.

1st of August, 2009

The Editor,
Conservation Genetics Resources

Dear Sir/Madam,

Please find attached a manuscript entitled "**Characterisation and cross-species utility of 20 microsatellite markers for population and forensic applications in the endangered Carnaby's Black-cockatoo, *Calyptorhynchus latirostris***" which we would like to submit as a technical note for consideration in *Conservation Genetics Resources*.

As a short communication, we believe the manuscript will add to the increased interest that cockatoos and parrots are contributing to our understanding of conservation and as animals that have a high profile in the illegal trade in wildlife.

This manuscript has never been previously submitted, published, nor is it being considered for publication in any journal other than *Conservation Genetics Resources*.

Kind regards

P. Spencer

(on behalf of both authors)