



Murdoch
UNIVERSITY

MURDOCH RESEARCH REPOSITORY

This is the author's final version of the work, as accepted for publication following peer review. The definitive version is available at <http://dx.doi.org/10.1016/j.aquaculture.2011.05.039>

Partridge, G.J., Benetti, D.D., Stieglitz, J.D., Hutapea, J., McIntyre, A., Chen, B., Hutchinson, W. and Scholey, V.P. (2011) *The effect of a 24-hour photoperiod on the survival, growth and swim bladder inflation of pre-flexion yellowfin tuna (*Thunnus albacares*) larvae. Aquaculture, 318 (3-4). pp. 471-474.*

<http://researchrepository.murdoch.edu.au/4858/>

Copyright © 2011 Elsevier B.V.

It is posted here for your personal use. No further distribution is permitted.

Accepted Manuscript

The effect of a 24-hour photoperiod on the survival, growth and swim bladder inflation of pre-flexion yellowfin tuna (*Thunnus albacares*) larvae

G.J. Partridge, D.D. Benetti, J.D. Stieglitz, J. Hutapea, A. McIntyre, B. Chen, W. Hutchinson, V.P. Scholey

PII: S0044-8486(11)00444-3
DOI: doi: [10.1016/j.aquaculture.2011.05.039](https://doi.org/10.1016/j.aquaculture.2011.05.039)
Reference: AQUA 629716

To appear in: *Aquaculture*

Received date: 31 August 2010
Revised date: 23 May 2011
Accepted date: 24 May 2011



Please cite this article as: Partridge, G.J., Benetti, D.D., Stieglitz, J.D., Hutapea, J., McIntyre, A., Chen, B., Hutchinson, W., Scholey, V.P., The effect of a 24-hour photoperiod on the survival, growth and swim bladder inflation of pre-flexion yellowfin tuna (*Thunnus albacares*) larvae, *Aquaculture* (2011), doi: [10.1016/j.aquaculture.2011.05.039](https://doi.org/10.1016/j.aquaculture.2011.05.039)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**The effect of a 24-hour photoperiod on the survival, growth and swim bladder
inflation of pre-flexion yellowfin tuna (*Thunnus albacares*) larvae.**

Partridge, G. J.^{a,b*} Benetti, D. D.^c Stieglitz, J. D.^c Hutapea, J.^d McIntyre, A.^e, Chen, B.^f
Hutchinson, W.^f Scholey, V. P.^g

- a. Australian Centre for Applied Aquaculture Research, Challenger Institute of Technology, Fremantle, Australia.
- b. Fish Health Unit, Centre for Fish and Fisheries Research, Murdoch University, Murdoch, Australia.
- c. Rosenstiel School Of Marine and Atmospheric Science, University of Miami, Florida, USA.
- d. Gondol Research Institute for Mariculture, Bali, Indonesia.
- e. Clean Seas Tuna, Port Lincoln, Australia.
- f. South Australian Research and Development Institute, Adelaide, Australia.
- g. Inter-American Tropical Tuna Commission, Los Santos, Panama.

* Corresponding author: Telephone: +61 8 9239 8032
Facsimile: +61 8 9239 8081
Email: gavin.partridge@challenger.wa.edu.au

Abstract

The effects of two different continuous photoperiod regimes on survival, growth and swim bladder inflation of pre-flexion yellowfin tuna (*Thunnus albacares*) larvae were investigated. Each photoperiod regime was tested twice with a different larval cohort to confirm the observed results. Trials 1 and 2 tested the effect of a reduced night-time light intensity ($10 \mu\text{moles m}^{-2}\text{sec}^{-1} = 30\%$ of the daytime intensity) and found that those larvae reared for 8 days under the 24 hour lighting (24-L) photoperiod exhibited a slight improvement in survival compared to those reared under the control photoperiod of 12 hours light (12-L); however these improvements were not significant. In addition, those larvae reared under this photoperiod regime were equal in length to those in the control. Trials 3 and 4 compared the same variables in larvae reared under a continuous photoperiod (24-L) with a constant light intensity of $30 \mu\text{moles m}^{-2}\text{sec}^{-1}$, against those reared under the aforementioned 12-L photoperiod. Survival of larvae under the continuous photoperiods were $9 \pm 1\%$ ($n = 2$) and $10 \pm 2\%$ ($n = 3$) for Trials 3 & 4, respectively, compared to less than 1% in both control treatments; differences that in both cases were highly significant. In addition, in both trials larvae cultured under the 24-L photoperiod were significantly larger and exhibited more advanced development than those reared under the 12-L photoperiod, however swim bladder inflation was significantly lower. We suggest that the improved survival and growth achieved under a continuous photoperiod is due to the extended foraging time combined with the prevention of mortality caused by night-time sinking.

Keywords: early larval rearing; early mortality; hatchery; buoyancy; swim bladder inflation.

1. Introduction

The global demand and subsequent production of tuna has increased exponentially over the last 50 years (Miyake et al., 2004). Approximately 32,000 MT of tuna marketed worldwide are now ranched; a process that involves the capture and fattening of wild caught juveniles or sub-adults (Lioka et al., 2000; Farwell, 2001). The consequent pressure on wild stocks is now threatening tuna fisheries worldwide, with some populations considered to be at the limits of sustainability, or overfished (Williams, 2007; Safina and Klinger, 2008; FAO, 2009). The total yield of tuna able to be produced via ranching is also limited via the quota restrictions placed on wild caught fish. In order to alleviate the pressures on wild tuna stocks; to overcome the constraints associated with ranching and to satisfy the growing demand for tuna, hatchery techniques for tuna must be developed. Major progress has recently been reported regarding reproduction, larval rearing and fingerling production of Pacific bluefin tuna (*Thunnus orientalis*) in Japan (Normile, 2009). In addition, progress has been reported regarding spawning of wild bluefin tuna broodstock as well as the rearing of their larvae in Australia and Europe (Mylonas et al., 2007; Hutchinson, 2009). Despite research efforts on tuna hatchery production for three decades in many countries, commercial scale production is yet to become reality, with survival rates of <0.5% to weaned juveniles typical for most tuna species (Sawada et al., 2005; Margulies et al., 2007).

The first feeding stage of all marine fish larvae is one of the most critical and high mortality is often experienced during this stage. In their review of the constraints to tuna larval rearing, Sawada et al. (2005) listed mortality during the first 10 days as one of the major limiting factors to mass culture and recommended focused research on this issue as a high priority.

Mortality of first feeding marine fish larvae has been attributed to factors including a poor transition from lecithotrophy to exotrophy (Fyhn, 1989); an inappropriate size and/or nutritional composition of feeds (Koven et al., 1992) or inappropriate environmental conditions (Takashi et al., 2006). Key environmental parameters to be considered for effective larval rearing include temperature and salinity (Fielder et al., 2005), the microbial environment (Harboe et al., 1994) and factors including turbulence, light intensity, light quality and photoperiod (Boeuf and Le Bail, 1999; Utne-Palm and Stiansen, 2002). Optimising photoperiod, for example, has been demonstrated to improve the performance of many species of marine fish larvae (Tandler and Helps, 1985; Duray and Kohno, 1988; Trotter et al., 2003); however, there appear to be no such published studies on tuna larvae. Lighting has, however, been shown to be of major importance in preventing high night-time mortalities caused by poor scotopic vision in early juvenile (30 DPH, 5 cm total length) Pacific bluefin tuna (Ishibashi et al., 2009).

These trials were conducted to determine if a continuous photoperiod would benefit the survival and growth of early stage yellowfin tuna (*Thunnus albacares*) larvae. In order to ensure the results of the experiment were not confounded by the physiologically

stressful event of flexion, the trial was conducted on pre-flexion larvae. Two sets of trials were conducted. The first investigated a continuous photoperiod, but with a reduced intensity during the 12 hour night phase, whilst the second set investigated a continuous photoperiod with equal light intensity during both night and day phases. To confirm the observed results, each trial was conducted twice with separate larval cohorts.

2. Materials and Methods

Larvae for each trial were sourced from a single spawning of eggs from naturally spawning yellowfin tuna broodstock at the Inter-American Tropical Tuna Commission's (IATTC) Achotines Laboratory in Los Santos Province, Republic of Panama. For details on the broodstock system and the management of these fish, refer to Wexler et al. (2003). Harvested eggs were incubated in 280 litre conical bottom tanks and hatching occurred after ca. 24 hours. Consistent with other published data on yellowfin tuna development, the day of hatching in this paper is referred to as Day 1. On Day 2, pre-feeding yolk-sac larvae were randomly stocked into 1000-litre, cylindrical, flat-bottom tanks (1 m diameter and 1 meter deep) with a number of replicated tanks per treatment (see Table 1). Larvae (2.47 ± 0.05 mm notochord length; NL) were stocked at a density of 15 larvae litre⁻¹ and reared using the standard larval rearing protocol employed by the IATTC (Margulies et al., 2007). The photoperiod regimes tested in each trial are outlined in Table 1. All control tanks operated under the standard photoperiod of 12 hours per day (6 am to 6 pm), while those in the treatment tanks received light for 24 hours per day. Treatment tanks were screened from control tanks to

ensure no incident light fell on the control tanks during their dark phase. Each tank was illuminated with 4 x 40 W fluorescent lamps (General Electric, Daylight F40D-EX), suspended 300 mm from the water surface. Surface light intensity was $30 \mu\text{moles m}^{-2}\text{sec}^{-1}$ (Li-Cor, LI250) (ca. 2,300 lux). The reduced light intensities employed during the night phase in the treatment tanks of Trials 1 & 2 were achieved by covering the lights with shade cloth, which reduced the light intensity to $10 \mu\text{moles m}^{-2}\text{sec}^{-1}$ (ca. 760 lux). Aeration in all tanks was provided by four, fine-bubble diffusers equally spaced across the tank base. The rate of aeration was set to achieve the optimum microturbulence for early feeding yellowfin tuna larvae ($1 - 2.4 \times 10^{-8} \text{ m}^{-2}\text{sec}^{-3}$ as an energy dissipation rate) by setting the mean horizontal velocity of a surface drogue to $5-7 \text{ cm sec}^{-1}$ (Kimura et al., 2004). Water temperature, dissolved oxygen, pH and salinity were measured in each tank three times per day.

Rotifers enriched on Algamac 3050 (Aquafauna Bio-Marine, Hawthorne, California) were maintained at a density of 5 mL^{-1} in all tanks throughout the trials. All tanks (including controls) were sampled every 6 hours and enriched rotifers added as necessary to maintain this density. Water flowed through the tanks via a $200 \mu\text{m}$ screen at the turn-over rate of $200\% \text{ day}^{-1}$. At this larval stocking density and flow rate, the removal of rotifers from the tank was dominated by flushing rather than ingestion and residual rotifer numbers were subsequently similar in all tanks at $1-2 \text{ mL}^{-1}$ prior to the next feed. Sufficient *Nannochloropsis oculata* was added by continuous siphon to maintain a cell density between 0.5 and $1.0 \times 10^6 \text{ cells mL}^{-1}$.

Trials 1 and 2 were terminated on Day 8 and Trials 3 and 4 on Day 9. At the conclusion of each trial, all tanks were drained and all remaining larvae counted. Twenty larvae from each tank were sampled. Notochord length was measured to 0.05 mm under a stereo microscope using a calibrated eye piece graticule and the presence or absence of a swim bladder was noted. Flexion was deemed to be commencing if signs of a developing caudal fin were evident under the notochord.

Survival, notochord length, percentage incidence of swim bladder inflation, percentage incidence of the commencement of flexion and the average daily values of each measured water quality parameter were compared between treatments with one-way analyses of variance. Percentage data (survival, swim bladder inflation and pre-flexion rates) were arcsine transformed prior to analysis. All reported errors are standard errors around the mean values and statements of significance refer to the 0.05 level, unless otherwise stated.

3. Results

In each trial, there were no significant differences in any of the measured water quality parameters between the control and photoperiod treatments (Table 2).

In Trials 1 and 2, larvae reared under continuous photoperiods displayed survival rates that were slightly higher than those of their respective controls (Table 3), however in both cases these improvements were not significant ($P > 0.05$). Those reared in Trial 1 achieved $2.3 \pm 0.6\%$ survival in the treatments and $1.0 \pm 0.6\%$ in the controls, whilst in

Trial 2 survival was $0.2 \pm 0.1\%$ and $0.1 \pm 0.1\%$ in the treatments and controls, respectively. Final notochord length of larvae in both experiments was equal between treatments and controls (4.59 ± 0.05 and 4.59 ± 0.16 mm, respectively for Trial 1 and 4.00 ± 0.27 and 3.97 ± 0.07 mm, respectively for Trial 2) (Table 3). No measurements of the onset of flexion or swim bladder inflation were made during these two trials.

Increasing the night-time light intensity to match that of the daytime intensity in Trials 3 and 4, resulted in significant improvements in larval survival. Larvae in the treatment tanks of Trial 3 had a survival of $8.9 \pm 0.8\%$ compared with $1.0 \pm 0.3\%$ in the control (Table 3); a difference that was highly significant ($P = 0.01$). Similarly, survival of larvae reared under the continuous photoperiod in Trial 4 was $10.4 \pm 1.9\%$, whilst the mean survival in the control treatment of this experiment averaged $0.2 \pm 0.1\%$; a difference that was again highly significant ($P = 0.005$) (Table 3). In contrast to Trials 1 and 2, larvae reared under the continuous photoperiod of Trials 3 and 4 also experienced significantly faster growth. Larvae cultured in the 24-L treatment from Trial 3 had a mean final notochord length of 5.1 ± 0.1 mm compared with 4.2 ± 0.1 mm in the control ($P = 0.03$) (Table 3). In Trial 4, treatment larvae averaged 4.5 ± 0.2 mm, compared with 3.9 ± 0.1 mm in the control ($P = 0.03$) (Table 3). Coincident with improved growth, larvae reared under continuous light were developmentally more advanced. In Trial 3, $90 \pm 0\%$ of those larvae reared under continuous light had commenced flexion by Day 9, compared with only $5 \pm 5\%$ in the control tanks ($P = 0.0006$). Similarly, in Trial 4, $90 \pm 3\%$ of larvae in the treatment tanks had commenced flexion by Day 9 compared with $52 \pm 7\%$ in the controls ($P = 0.006$).

The continuous photoperiod employed in Trial 4 had a significant detrimental impact on swim bladder inflation ($P=0.001$). On Day 9, $10 \pm 0\%$ of larvae in the treatment tanks had inflated swim bladders, compared with $50 \pm 5\%$ in the controls.

4. Discussion

The results of this study demonstrate that providing continuous light with an intensity of $30 \mu\text{moles m}^{-2}\text{sec}^{-1}$ significantly improves both survival and growth of yellowfin tuna larvae. During food selectivity studies, Margulies et al. (2007) reported that larval yellowfin tuna feed only during daylight hours, confirming that like most species they are visual predators (Blaxter, 1986). We therefore attribute the improved growth and advanced development of larvae reared under continuous light in Trials 3 and 4 to a longer foraging time and subsequently greater food intake. Larvae reared under the continuous photoperiod regimes of Trials 1 & 2 showed no improvement in growth, suggesting that either the reduced night-time light of $10 \mu\text{moles m}^{-2}\text{sec}^{-1}$ was below the threshold of light required for yellowfin tuna larvae to see and capture their prey or that the level of ingestion of prey at this light level was insufficient to offset any prolonged activity of the larvae at this light intensity.

The positive correlation between larval growth and continuous light and feeding appears to be common amongst many species of marine fish, including sole (*Solea solea*) (Fuchs, 1978), gilthead seabream (*Sparus auratus*) (Tandler and Helps, 1985), rabbitfish (*Siganus guttatus*) (Duray and Kohno, 1988), barramundi (*Lates calcarifer*) (Barlow et al., 1995), greenback flounder (*Rhombosolea tapirina*) (Hart et al., 1996),

Australian snapper (*Pagrus auratus*) (Fielder et al., 2002) and Atlantic cod (*Gadus morhua*) (Puvanendran and Brown, 2002). Downing and Litvak (2000) reported no benefit to haddock (*Melanogrammus aeglefinus*) larvae reared under continuous light, however their control photoperiod of 15L:9D was considerably longer than most other studies. There appears to be only one study in which an extended photoperiod was detrimental to the growth of a marine fish larvae, the sea bream (*Archosargus rhomboidalis*) (Dowd and Houde, 1980).

Despite the consistent, positive relationship between photoperiod and larval growth described above for a wide range of species, the effect on survival is more equivocal. The majority of studies report no difference in survival between natural and extended photoperiods (Fuchs, 1978; Barlow et al., 1995; Hart et al., 1996; Fielder et al., 2002; Moustakas et al., 2004); however, a 24 hour photoperiod had a detrimental effect on the survival of European seabass *Dicentrarchus labrax* (Barahona-Fernandes, 1979; Ronzani Cerqueira and Chatain, 1991). Tandler and Helps (1985), Duray and Kohno (1988) and Trotter et al. (2003) all reported significantly higher survival under continuous light for gilthead seabream, rabbitfish and striped trumpeter, respectively, although the gains in survival were not as great as those reported here for yellowfin tuna larvae.

The improved survival obtained in the 24 hour light treatments with continuous intensity in this study may be attributable to a greater number of larvae successfully commencing feeding. Fielder et al. (2002) pointed out that prey capture is a learned response, and extended photoperiods may therefore provide a greater opportunity for first feeding larvae to learn this vital lesson.

Alternatively, continuous lighting may have improved survival by preventing negatively buoyant larvae from sinking to the bottom of the tank at night. Takashi et al. (2006) pointed out that a major cause of mortality of early stage Pacific bluefin tuna larvae is their tendency to sink to the tank bottom at night and showed that the density of larvae increases with increasing age. Between Days 4 and 8 these larvae are denser than seawater, even with an inflated swim bladder, and at night when swimming ceases they sink to the bottom of the tank and die. From Day 9, swim bladder volume of Pacific bluefin tuna larvae greatly increases, which assists in keeping them suspended at night. In addition, once the process of flexion is complete, the larvae become more efficient swimmers and perhaps more adept at swimming away from the tank bottom. Regardless of the mode, mortality due to sinking becomes less problematic from Day 8-9 (Takashi et al., 2006). We believe that yellowfin tuna are similar to Pacific bluefin in this regard, as mortalities are commonly seen on the tank bottom in the early morning when reared under the standard photoperiod.

The results of this study demonstrate that the mode of action for the improved survival of larval tuna from the provision of 24 hour light is different to that for juvenile tuna. Ishibashi et al (2009) reported that the improved survival obtained in juvenile Pacific bluefin tuna in the 23 days post transfer from tanks to seacages was due to the prevention of collisions with the walls of the net pen. Due to the inability of early larvae to burst swim, lethal impact collisions with tank walls is not believed to be a cause of larval mortality.

A continuous photoperiod had a negative impact on swim bladder inflation in the current study. As discussed by Trotter et al. (2003), many physostomic fish larvae inflate their swim bladder at the water surface in darkness and continuous lighting can therefore inhibit this process. This appears to be the case with yellowfin tuna larvae, however swim bladder inflation in the control treatment was also relatively poor. Munday et al. (2003) pointed out that it is common practice to place a layer of oil on the surface of bluefin tuna larval rearing tanks during the first 7 days to prevent larvae getting trapped in the surface tension. This suggests it may be possible for tuna larvae to inflate their swim bladder after this time. It may therefore be necessary to shift from a 24 hour photoperiod, to one with at least some darkness once the larvae have passed the negatively buoyant stage in order to allow swim bladder inflation to occur.

Although Takashi et al. (2006) pointed out that photoperiod manipulation may be effective in preventing mortality of tuna larvae due to sinking, this appears to be the first study to have investigated the effects of a continuous photoperiod on the larvae of any tuna species. Previous trials conducted by the IATTC at the Ashotines Laboratory have indicated variable patterns of larval growth and survival in response to extended photoperiods (D. Margulies, pers. comm.). Given the highly significant benefits to growth and survival reported here using a 24 hour period, further detailed investigations into the use of extended photoperiods for tuna hatchery production are clearly warranted. Additional research should confirm the mode of action of the improved survival and whether continuing a 24 photoperiod beyond the pre-flexion stage remains beneficial. More data are required to determine the window of opportunity for swim

bladder inflation and to optimise photoperiod strategies to ensure the process of swim bladder inflation is not compromised.

Acknowledgements

These experiments were conducted during the 6th, 7th and 8th Annual Tuna Workshop held at the Achotines Laboratory; organized and run by the University of Miami Rosenstiel School of Marine and Atmospheric Science (RSMAS) and the Inter-American Tropical Tuna Commission (IATTC). We would like to thank all staff at the Achotines Laboratory as well as the following participants of these workshops for their assistance; Todd Glodek, Kevin Polk, Francesca Forrestal, Alex Muhlholzl, Pollyanna Hilder, Luke Cheviot, Luke Vandenberg, Zac Daugherty and Patrick Dunaway. We are especially grateful for Dr. Daniel Margulies of the IATTC for his valuable comments and suggestions. Gavin Partridge acknowledges funding from the Aquaculture Development Council of Western Australia and a Fellowship from the International Specialised Skills Institute, supported by The Department of Education, Employment and Workplace Relations. Daniel Benetti and John Stieglitz acknowledge funding from NOAA-SK Grant # NA09NMF4270080. Jhon Hutapea acknowledges funding from ACIAR Grant # FIS/2006/140.

References

- Barahona-Fernandes, M.H., 1979. Some effects of light intensity and photoperiod on the sea bass larvae (*Dicentrarchus labrax* (L.)) reared at the Centre Oceanologique de Bretagne. *Aquaculture* 17, 311-321.
- Barlow, C.G., Pearce, M.G., Rodgers, L.J., Clayton, P., 1995. Effects of photoperiod on growth, survival and feeding periodicity of larval and juvenile barramundi *Lates calcarifer* (Bloch). *Aquaculture* 138, 159-168.
- Blaxter, J.H.S., 1986. Development of sense organs and behaviour of teleost larvae with special reference to feeding and predator avoidance. *Trans. Am. Fish. Soc.* 115, 98-114.
- Boeuf, G., Le Bail, P.-Y., 1999. Does light have an influence on fish growth? *Aquaculture* 177, 129-152.
- Dowd, C.E., Houde, E.D., 1980. Combined effects of prey concentration and photoperiod on survival and growth of larval sea bream, *Archosargus rhomboidalis* (Sparidae). *Mar. Ecol. Prog. Ser.* 3, 181-185.
- Downing, G., Litvak, M.K., 2000. The effect of photoperiod, tank colour and light intensity on growth of larval haddock. *Aquac. Int.* 7, 369-382.
- Duray, M., Kohno, H., 1988. Effects of continuous lighting on growth and survival of first-feeding larval rabbitfish, *Siganus guttatus*. *Aquaculture* 72, 73-79.
- FAO, 2009. State of world fisheries and aquaculture 2008. Food and Agriculture Organization of the United Nations, Fisheries and Aquaculture Department, Rome, pp. 196.

- Farwell, C., 2001. Tunas in captivity. In: Block, B.A., Stevens, E.D. (Eds.), Tuna: Physiology, Ecology, and Evolution. Academic Press, London, pp. 391-412.
- Fielder, D.S., Bardsley, W.J., Allan, G.L., Pankhurst, P.M., 2002. Effect of photoperiod on growth and survival of snapper *Pagrus auratus* larvae. *Aquaculture* 211, 135-150.
- Fielder, D.S., Bardsley, W.J., Allan, G.L., Pankhurst, P.M., 2005. The effects of salinity and temperature on growth and survival of Australian snapper, *Pagrus auratus* larvae. *Aquaculture* 250, 201-214.
- Fuchs, J., 1978. Effect of photoperiod on growth and survival during rearing of larvae and juveniles of sole (*Solea solea*). *Aquaculture* 15, 63-74.
- Fyhn, H., 1989. First feeding of marine fish larvae: Are free amino acids the source of energy? *Aquaculture* 80, 111-120.
- Harboe, T., Huse, I., Øie, G., 1994. Effects of egg disinfection on yolk sac and first feeding stages of halibut (*Hippoglossus hippoglossus* L.) larvae. *Aquaculture* 119, 157-165.
- Hart, P.R., Hutchinson, W.G., Purser, G.J., 1996. Effects of photoperiod, temperature and salinity on hatchery-reared larvae of the greenback flounder (*Rhombosolea tapirina* Günther, 1862). *Aquaculture* 144, 303-311.
- Hutchinson, W.G., 2009. Southern bluefin tuna (*Thunnus maccoyii*) larval rearing advances at the South Australian Research and Development Institute and collaborating institutions. In: Allan, G., Booth, M., Mair, G., Clarke, S., Biswas, A. (Eds.), Proceedings of the 2nd Global COE Program Symposium of Kinki University, Sustainable Aquaculture of the Bluefin and Yellowfin Tuna -

- Closing the Life Cycle for Commercial Production. Kinki University, SARDI, Adelaide South Australia.
- Ishibashi, Y., Honryo, T., Saida, K., Hagiwara, A., Miyashita, S., Sawada, Y., Okada, T., Kurata, M., 2009. Artificial lighting prevents high night-time mortality of juvenile Pacific bluefin tuna, *Thunnus orientalis*, caused by poor scotopic vision. *Aquaculture* 293, 157-163.
- Kimura, S., Nakata, H., Margulies, D., Suter, J., Hunt, S., 2004. Effect of oceanic turbulence on the survival of yellowfin tuna larvae. *Nippon Suisan Gakkaishi* 70, 175-178.
- Koven, W.M., Tandler, A., Kissil, G.W., Sklan, D., 1992. The importance of n-3 highly unsaturated fatty acids for growth in larval *Sparus aurata* and their effect on survival, lipid composition and size distribution. *Aquaculture* 104, 91-104.
- Lioka, C., Kani, K., Nhhala, H., 2000. Present status and prospects of technical development of tuna sea-farming. *Cahiers Options Méditerranéennes* 47, 275-285.
- Margulies, D., Scholey, V.P., Wexler, J.B., Olson, R.J., Suter, J.M., Hunt, S.L., 2007. A Review of IATTC Research on the Early Life History and Reproductive Biology of Scombrids Conducted at the Achotines Laboratory from 1985 to 2005. Inter-American Tropical Tuna Commission, La Jolla, California, pp. 67.
- Miyake, M.P., Miyabe, N., Nakano, H., 2004. Historical trends of tuna catches in the world. FAO, Rome, pp. 74.
- Moustakas, C.T., Watanabe, W.O., Copeland, K.A., 2004. Combined effects of photoperiod and salinity on growth, survival, and osmoregulatory ability of larval southern flounder *Paralichthys lethostigma*. *Aquaculture* 229, 159-179.

- Munday, B.L., Sawada, Y., Cribb, T., Hayward, C.J., 2003. Diseases of tunas, *Thunnus spp.* J. Fish Dis. 26, 187-206.
- Mylonas, C.C., Bridges, C.R., Gordin, H., Belmonte, A., Garcia-Gomez, A., De La Gandara, F., Fauvel, C., Suquet, M., Medina, A., Papadaki, M., Heinisch, G., De Metrio, G., Corriero, A., Vassallo-Agius, R., Guzman, J.M., Mañanos, E., Zohar, Y., 2007. Preparation and administration of gonadotropin-releasing hormone agonist (GnRH α) implants for the artificial control of reproductive maturation in captive-reared Atlantic bluefin tuna (*Thunnus thynnus thynnus*). Rev. Fish. Sci. 15, 183-210.
- Normile, D., 2009. Persevering researchers make a splash with farm-bred tuna. Science 324, 1260-1261.
- Puvanendran, V., Brown, J.A., 2002. Foraging, growth and survival of Atlantic cod larvae reared in different light intensities and photoperiods. Aquaculture 214, 131-151.
- Ronzani Cerqueira, V., Chatain, B., 1991. Photoperiodic effects on the growth and feeding rhythm of European seabass, *Dicentrarchus labrax*, larvae in intensive rearing. In: Lavens, P., Sorgeloos, P., Jaspers, E., Ollevier, F. (Eds.), Larvi '91 - Fish and Crustacean Larviculture Symposium. European Aquaculture Society, Gent, Belgium.
- Safina, C., Klinger, D.H., 2008. Collapse of bluefin tuna in the western Atlantic. Conserv. Biol. 22, 243-246.
- Sawada, Y., Okada, T., Miyashita, S., Murata, O., Kumai, H., 2005. Completion of the Pacific bluefin tuna *Thunnus orientalis* (Temminck and Schlegel) life cycle. Aquac. Res. 36, 413-421.

- Takashi, T., Hirotohi Kohno, Wataru Sakamoto, Shigeru Miyashita, Osamu Murata, Yoshifumi Sawada, 2006. Diel and ontogenetic body density change in Pacific bluefin tuna, *Thunnus orientalis* (Temminck and Schlegel), larvae. *Aquac. Res.* 37, 1172-1179.
- Tandler, A., Helps, S., 1985. The effects of photoperiod and water exchange rate on growth and survival of gilthead sea bream (*Sparus aurata*, Linnaeus; Sparidae) from hatching to metamorphosis in mass rearing systems. *Aquaculture* 48, 71-82.
- Trotter, A.J., Battaglione, S.C., Pankhurst, P.M., 2003. Effects of photoperiod and light intensity on initial swim bladder inflation, growth and post-inflation viability in cultured striped trumpeter (*Latris lineata*) larvae. *Aquaculture* 224, 141-158.
- Utne-Palm, A.C., Stiansen, J.E., 2002. Effect of larval ontogeny, turbulence and light on prey attack rate and swimming activity in herring larvae. *J. Exp. Mar. Biol. Ecol.* 268, 147-170.
- Wexler, J.B., Scholey, V.P., Olson, R.J., Margulies, D., Nakazawa, A., Suter, J.M., 2003. Tank culture of yellowfin tuna, *Thunnus albacares*: developing a spawning population for research purposes. *Aquaculture* 220, 327-353.
- Williams, N., 2007. Tuna crisis looms. *Curr. Biol.* 17, R108-R109.

Table 1. Photoperiod and light intensity investigated in each trial.

Trial	Photoperiod (hours)	Day Phase Intensity ($\mu\text{moles m}^{-2}\text{sec}^{-1}$)	Night Phase Intensity ($\mu\text{moles m}^{-2}\text{sec}^{-1}$)	<i>n</i>
1	24-L	30	10	3
	12-L	30	0	
2	24-L	30	10	3
	12-L	30	0	
3	24-L	30	30	2
	12-L	30	0	
4	24-L	30	30	3
	12-L	30	0	

Table 2. Average daily water quality parameters (\pm S.E.) measured in each trial.

Trial	Photoperiod (hours)	Temperature ($^{\circ}$ C)	Dissolved Oxygen (mg/L)	pH	Salinity (g/L)
1	24-L	28.38 ± 0.69	6.25 ± 0.01	8.20 ± 0.05	31.36 ± 0.01
	12-L	27.73 ± 0.02	6.28 ± 0.01	8.25 ± 0.05	31.35 ± 0.01
2	24-L	27.11 ± 0.05	6.58 ± 0.02	8.19 ± 0.05	30.94 ± 0.01
	12-L	27.12 ± 0.02	6.56 ± 0.01	8.20 ± 0.05	30.94 ± 0.01
3	24-L	27.34 ± 0.02	6.36 ± 0.01	7.93 ± 0.01	32.10 ± 0.01
	12-L	27.44 ± 0.01	6.32 ± 0.07	7.93 ± 0.01	32.10 ± 0.01
4	24-L	27.59 ± 0.01	5.99 ± 0.06	8.04 ± 0.02	31.34 ± 0.01
	12-L	27.45 ± 0.01	5.80 ± 0.05	8.07 ± 0.01	31.34 ± 0.01

ACCEPTED

Table 3. Final notochord length (mm), % survival, % swim bladder inflation and % of yellowfin tuna larvae commencing flexion from Trials 1 to 4. All data are mean \pm S.E. Within each trial, parameters sharing the same letter are not significantly different ($P > 0.05$). Refer to Table 1 for light intensities within each photoperiod.

Trial	1		2		3		4	
	24-L	12-L	24-L	12-L	24-L	12-L	24-L	12-L
Survival (%)	2.3 \pm 0.6 ^a	1.0 \pm 0.6 ^a	0.2 \pm 0.1 ^a	0.1 \pm 0.1 ^a	8.9 \pm 0.8 ^b	1.0 \pm 0.3 ^a	10.4 \pm 1.9 ^b	0.2 \pm 0.1 ^a
Final notochord length (mm)	4.59 \pm 0.05 ^a	4.59 \pm 0.16 ^a	4.00 \pm 0.27 ^a	3.97 \pm 0.07 ^a	5.1 \pm 0.1 ^b	4.2 \pm 0.1 ^a	4.5 \pm 0.2 ^b	3.9 \pm 0.1 ^a
Initiating flexion (%)	-	-	-	-	90 \pm 0 ^b	5 \pm 5 ^a	90 \pm 3 ^b	52 \pm 7 ^a
Swim bladder inflation (%)	-	-	-	-	-	-	10 \pm 0 ^b	50 \pm 5 ^a