



Corrigendum: Hypocoagulability and Platelet Dysfunction Are Exacerbated by Synthetic Colloids in a Canine Hemorrhagic Shock Model

Corrin J. Boyd*, Melissa A. Claus, Anthea L. Raisis, Giselle Hosgood, Claire R. Sharp and Lisa Smart

School of Veterinary and Life Sciences, College of Veterinary Medicine, Murdoch University, Perth, WA, Australia

Keywords: hydroxyethyl starch, succinylated gelatin, crystalloid, fresh whole blood, platelet closure time, PFA-100, rotational thromboelastometry (ROTEM), viscoelastic coagulation tests

A Corrigendum on

Hypocoagulability and Platelet Dysfunction Are Exacerbated by Synthetic Colloids in a Canine Hemorrhagic Shock Model

by Boyd, C. J., Claus, M. A., Raisis, A. L., Hosgood, G., Sharp, C. R., and Smart, L. (2018). *Front. Vet. Sci.* 5:279. doi: 10.3389/fvets.2018.00279

In the original article, there was a mistake in Table 2 as published. The EXTEM CFT baseline mean (95% confidence interval) for the HES group was slightly incorrect due to a data entry error. The corrected Table 2 appears below.

In the original article, there was a mistake in Table 5 as published. The EXTEM CFT baseline mean (95% confidence interval) was slightly incorrect due to a data entry error. The corrected Table 5 appears below.

In the original article, there was an error. A conversion factor was inadvertently omitted from the equation for thrombodynamic potential index (TPI).

A correction has been made to *Materials and Methods, Coagulation Parameters, Paragraph 3:*

Rotational thromboelastometry (ROTEM[®] delta, Tem International GmbH) was performed according to the manufacturer's instructions and PROVETS guidelines (28) using the INTEM (star-TEM and in-TEM reagents), EXTEM (star-TEM and r ex-TEM reagents), FIBTEM (r ex-TEM and fib-TEM reagents), and APTEM (r ex-TEM and ap-TEM reagents) profiles. Measurement was started 30 min after sample collection. Each profile was run for at least 1 h following initiation. Data recorded for the INTEM and EXTEM profiles included clotting time (CT), clot formation time (CFT), alpha angle (α), maximum clot firmness (MCF), and lysis index at 60 min (LI60). Thrombodynamic potential index (TPI) was recorded for the EXTEM profile as a measure of global coagulation (29), calculated using the equations:

$$\begin{aligned} \text{TPI} &= \text{EMX} * 30 / \text{CFT}, \\ \text{EMX} &= (100 * \text{MCF}) / (100 - \text{MCF}). \end{aligned}$$

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2020 Boyd, Claus, Raisis, Hosgood, Sharp and Smart. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

OPEN ACCESS

Edited and reviewed by:

Katja-Nicole Adamik,
University of Bern, Switzerland

*Correspondence:

Corrin J. Boyd
c.boyd@murdoch.edu.au

Specialty section:

This article was submitted to
Veterinary Emergency and Critical
Care Medicine,
a section of the journal
Frontiers in Veterinary Science

Received: 29 June 2020

Accepted: 06 August 2020

Published: 08 September 2020

Citation:

Boyd CJ, Claus MA, Raisis AL,
Hosgood G, Sharp CR and Smart L
(2020) Corrigendum:
Hypocoagulability and Platelet
Dysfunction Are Exacerbated by
Synthetic Colloids in a Canine
Hemorrhagic Shock Model.
Front. Vet. Sci. 7:641.
doi: 10.3389/fvets.2020.00641

TABLE 2 | Rotational thromboelastometry parameters (mean, 95% confidence interval) in dogs ($n = 6$ per group) with hemorrhagic shock given 20 mL kg⁻¹ of either fresh whole blood (FWB), hydroxyethyl starch 130/0.4 (HES), 4% succinylated gelatin (GELO), or 80 mL kg⁻¹ of balanced isotonic crystalloid (CRYST).

Parameter	Baseline	Shock	T60	T180
INTEM CT (sec)				
FWB	279.3 (137.5–421.2)	204.8 (123.1–286.5)	330.5 (17.2–643.8)	269.0 (150.7–387.3)
CRYST	171.3 (149.1–193.5)	160.7 (129.2–192.1)	162.2 (134.0–190.3)	159.8 (142.8–176.9)
HES	201.2 (129.1–273.2)	145.8 (72.2–219.5)	231.8 (130.4–333.3)	199.0 (113.0–285.0)
GELO	221.5 (146.0–297.0)	161.0 (117.0–205.0)	224.3 (108.6–340.1)	193.5 (146.5–240.5)
INTEM CFT (sec)				
FWB	224.2 (143.1–305.2)	260.3 (180.6–340.0)	303.8 (102.0–505.7)	235.3 (178.9–291.8)
CRYST	143.5 (119.1–167.9)	227.8 (190.9–264.8)	298.8 (269.5–328.1)	234.5 (207.6–261.4)
HES	202.8 (106.9–298.8)	264.0 (168.0–360.0)	431.5 (287.3–575.7)	309.2 (221.0–397.3)
GELO	178.2 (128.4–227.9)	186.2 (114.2–258.1)	305.5 (211.1–399.9)	278.7 (178.1–379.2)
INTEM α (°)				
FWB	54.2 (44.9–63.4)	54.0 (46.3–61.7)	52.7 (38.5–66.8)	56.2 (49.8–62.6)
CRYST	68.0 (65.4–70.6)	62.7 (57.2–68.1)	56.8 (53.7–60.0)	60.7 (59.8–61.5)
HES	60.5 (49.4–71.6)	56.0 (46.5–65.5)	50.7 (40.2–61.1)	53.5 (46.4–60.6)
GELO	61.2 (56.5–65.9)	61.7 (53.8–69.6)	53.0 (46.7–59.3)	56.2 (49.8–62.6)
INTEM MCF (mm)				
FWB	51.2 (46.9–55.4)	47.7 (43.0–52.3)	45.3 (38.3–52.3) ⁺	48.2 (44.5–51.8) ⁺
CRYST	54.2 (50.8–57.5)	46.3 (43.0–49.7)	42.2 (39.9–44.4)	44.0 (41.4–46.6)
HES	50.3 (45.3–55.4)	44.5 (40.3–48.7)	37.0 (31.6–42.4) ^{&c}	41.0 (36.2–45.8) ^{&c}
GELO	54.0 (48.3–59.7)	51.2 (45.8–56.5)	41.5 (36.0–47.0)	42.8 (37.2–48.5)
INTEM LI60 (%)				
FWB	99.0 (97.4–100.6)	99.7 (98.8–100.5)	99.7 (98.8–100.5)	99.2 (97.9–100.4)
CRYST	99.7 (98.8–100.5)	98.7 (97.6–99.8)	96.8 (94.7–99.0)	97.2 (94.4–99.9)
HES	99.2 (97.9–100.4)	98.5 (96.4–100.6)	87.8 (67.3–108.4)	90.0 (74.1–105.9)
GELO	99.5 (98.6–100.4)	99.7 (99.1–100.2)	98.3 (94.0–102.6)	96.5 (87.5–105.5)
EXTEM CT (sec)				
FWB	52.5 (40.8–64.2)	75.2 (57.2–93.1)	64.7 (45.3–84.1) ⁺	62.0 (53.1–70.9)
CRYST	54.8 (37.8–71.9)	66.2 (59.4–72.9)	72.8 (62.9–82.8) ⁺	71.5 (58.7–84.3)
HES	61.8 (40.6–83.0)	69.0 (56.1–81.9)	119.3 (57.4–181.3) ^{&c,*:#}	82.7 (50.6–114.8)
GELO	47.8 (28.6–67.0)	55.8 (36.1–75.6)	76.2 (27.5–124.8) ⁺	75.2 (50.6–99.8)
EXTEM CFT (sec)				
FWB	178.2 (121.5–234.8)	217.3 (149.5–285.2)	251.0 (127.6–374.4)	202.2 (148.5–255.9)
CRYST	129.3 (115.4–143.3)	184.2 (152.6–215.7)	246.0 (198.3–293.7)	221.0 (177.7–264.3)
HES	187.2 (106.0–268.3)	208.0 (137.8–278.2)	373.4 (179.9–566.9)	437.3 (0.0–918.6)
GELO	139.3 (102.6–176.1)	165.0 (115.5–214.5)	230.4 (167.7–293.1)	263.3 (174.5–352.1)
EXTEM α (°)				
FWB	59.8 (53.5–66.1)	60.0 (55.9–64.1)	55.8 (48.8–62.9)	58.3 (54.4–62.3)
CRYST	66.5 (64.4–68.6)	63.3 (60.9–65.8)	59.3 (54.2–64.5)	60.8 (59.0–62.6)
HES	61.7 (55.1–68.2)	61.8 (56.4–67.2)	46.2 (28.6–63.8)	53.0 (39.8–66.2)
GELO	65.7 (62.8–68.5)	63.5 (59.9–67.1)	49.5 (32.7–66.3)	56.0 (52.5–59.5)
EXTEM MCF (mm)				
FWB	50.5 (46.3–54.7)	44.7 (39.6–49.7)	45.3 (38.6–52.1) ⁺	47.0 (43.4–50.6)
CRYST	54.7 (52.1–57.2)	45.7 (42.1–49.3)	39.0 (33.3–44.7)	39.3 (34.7–44.0)
HES	49.3 (42.7–55.9)	42.5 (33.4–51.6)	31.2 (20.5–41.9) ^{&c}	37.8 (28.6–47.1)
GELO	54.3 (47.8–60.8)	50.0 (43.4–56.6) ^β	39.0 (26.6–51.4)	40.3 (31.9–48.8)
EXTEM TPI				
FWB	18.8 (12.7–25.0)	12.5 (6.7–18.3)	12.3 (5.6–19.0)	14.0 (9.6–18.4)
CRYST	28.5 (22.8–34.2)	14.3 (10.5–18.2)	8.2 (5.6–10.8)	9.3 (6.1–12.6)
HES	20.2 (7.4–32.9)	14.2 (1.1–27.3)	4.8 (2.0–7.6)	8.2 (1.4–14.9)
GELO	28.8 (15.3–42.4)	21.0 (10.3–31.7)	11.2 (5.3–17.1)	9.5 (4.2–14.8)

(Continued)

TABLE 2 | Continued

Parameter	Baseline	Shock	T60	T180
EXTEM LI60 (%)				
FWB	89.3 (66.9–111.7)	78.7 (48.1–109.3)	90.7 (70.7–110.7)	91.0 (78.2–103.8)
CRYST	99.0 (97.9–100.1)	87.7 (73.0–102.3)	84.7 (52.9–116.5)	78.8 (58.5–99.2)
HES	98.3 (95.5–101.2)	83.3 (56.1–110.6)	74.3 (36.5–112.2)	84.3 (56.2–112.5)
GELO	97.5 (93.8–101.2)	86.7 (65.3–108.1)	75.8 (49.2–102.5)	72.5 (42.5–102.5)
FIBTEM MCF (mm)				
FWB	4.3 (3.5–5.2)	3.5 (2.4–4.6)	3.3 (1.5–5.2)	2.8 (1.0–4.6)
CRYST	6.0 (3.1–8.9)	4.7 (1.9–7.5)	2.8 (1.0–4.6)	3.2 (1.2–5.1)
HES	5.0 (2.6–7.4)	2.0 (0.0–4.9)	0.7 (0.0–2.4)	2.2 (0.0–5.2)
GELO	4.8 (3.3–6.4)	3.8 (2.3–5.4)	2.0 (0.2–3.8)	2.7 (1.1–4.2)

Fluid was delivered directly after Shock time point.

& Significantly different to FWB; *Significantly different to CRYST; †Significantly different to HES; #Significantly different to GELO (Bonferroni–Holm corrected P -value < 0.05). α , alpha angle; CFT, clot formation time; CT, clotting time; LI60, lysis index at 60 min; MCF, maximum clot firmness; TPI, thrombodynamic potential index.

TABLE 5 | Platelet closure time and count, rotational thromboelastometry, and plasma coagulation assay parameters (mean, 95% confidence interval) in dogs ($n = 24$) with hemorrhagic shock.

Parameter	Baseline	Shock	P value*
Platelet Closure Time (sec)	83.4 (77.6–89.1)	69.0 (65.2–72.8)	<0.001
Estimated Platelet Count ($\times 10^9 \text{ L}^{-1}$)	134.3 (124.7–143.9)	119.4 (106.3–132.5)	0.057
INTEM CT (sec)	218.3 (180.9–255.7)	168.1 (143.2–193.0)	<0.001
INTEM CFT (sec)	158.5 (135.9–181.1)	234.6 (203.6–265.6)	<0.001
INTEM α (°)	61.0 (57.4–64.5)	58.6 (55.3–61.9)	0.067
INTEM MCF (mm)	52.4 (50.5–54.3)	47.4 (45.4–49.4)	<0.001
INTEM LI60 (%)	99.3 (98.9–99.8)	99.1 (98.6–99.7)	0.458
EXTEM CT (sec)	54.3 (47.3–61.2)	66.5 (60.1–73.0)	<0.001
EXTEM CFT (sec)	148.0 (125.3–170.7)	193.6 (170.5–216.8)	<0.001
EXTEM α (°)	63.4 (61.2–65.6)	62.2 (60.5–63.8)	0.117
EXTEM MCF (mm)	52.2 (50.0–54.4)	45.7 (43.0–48.4)	<0.001
EXTEM LI60 (%)	96.0 (91.4–100.7)	84.1 (74.9–93.3)	0.002
EXTEM TPI	24.1 (19.8–28.4)	15.5 (11.8–19.2)	<0.001
FIBTEM MCF (mm)	5.0 (4.2–5.9)	3.5 (2.6–4.4)	<0.001
Prothrombin Time (sec)	7.5 (7.3–7.7)	8.5 (8.2–8.8)	<0.001
Activated Partial Thromboplastin Time (sec)	15.4 (14.8–16.0)	14.4 (13.6–15.1)	<0.001
Fibrinogen Concentration (g L^{-1})	1.41 (1.29–1.54)	0.97 (0.87–1.08)	<0.001
Factor VII Activity (%)	99.1 (92.5–105.7)	77.8 (71.7–84.0)	<0.001
Factor VIII Activity (%)	109.8 (98.0–121.6)	155.4 (139.9–171.0)	<0.001
von Willebrand Factor Antigen (%)	100.6 (90.0–111.3)	74.3 (60.2–88.4)	<0.001

*Paired t -test, bold indicates $P < 0.05$. α , alpha angle; CFT, clot formation time; CT, clotting time; LI60, lysis index at 60 min; MCF, maximum clot firmness; TPI, thrombodynamic potential index.